MATH 313 HMWK 3

Daniel Xu

February 9, 2018

Exercise 2.2.1 An example of a vercongent sequence would be $x_n = \sin x$. Let $\epsilon = \frac{101}{100}$ and let x = 0. Then x_n converges to 0, since $|\sin x - 0| < \frac{101}{100}$. Since x_n does not converge to any number, it is a vercongent sequence that is also divergent. What a vercongent sequence describes is a sequence for whom all values lie in a certain bound, namely $(x - \epsilon, x + \epsilon)$.

Excercise 2.2.2

1. Let $\epsilon > 0$ be arbitrary. Now choose N such that

$$N > \frac{3}{25\epsilon} - \frac{4}{5}$$

To verify that our choice of N is appropiate, let $n \in \mathbb{N}$ satisfy $n \geq N$. Then, $n \geq N$ implies that

$$n > \frac{3}{25\epsilon} - \frac{4}{5}$$

$$5n > \frac{3}{5\epsilon} - 4$$

$$5n + 4 > \frac{3}{5\epsilon}$$

$$\epsilon > \frac{3}{5 \cdot (5n+4)}$$

$$\epsilon > \frac{2 \cdot (5n+4) - 5 \cdot (2n+1)}{5 \cdot (5n+4)}$$

$$\epsilon > \frac{2 \cdot (5n+4) - \frac{5 \cdot (2n+1)}{5 \cdot (5n+4)}$$

$$\epsilon > \frac{2}{5} - \frac{2n+1}{5n+4}$$

$$\epsilon > \left| \frac{2}{5} - \frac{2n+1}{5n+4} \right|$$

$$\left| \frac{2}{5} - \frac{2n+1}{5n+4} \right| < \epsilon$$

$$\left| \frac{2n+1}{5n+4} - \frac{2}{5} \right| < \epsilon$$

2.

2.3.1

1. If $x_n \to 0$, then there exists $\epsilon_0 > 0$ such that there exists $N \in \mathbb{N}$ for all $n \geq \mathbb{N}, x_n - 0 < \epsilon_0$.

2.3.1

1. If $x_n\to 0$, then it must be true that for all ϵ_0 , there exists a $N\in\mathbb{N}$ such that for all $n\ge N,\, |x_n-0|<\epsilon_0.$ Then

$$|x_n - 0| < \epsilon_0$$

 $x_n < \epsilon_0$ since $x_n \ge 0$
 $\sqrt{x_n} < \sqrt{\epsilon_0}$

Exercise 2.3.7

- 1. Let $x_n = n$ and $y_n = -n$. Both x_n and y_n diverge. However, $x_n + y_n$ converge to zero.
- 2. Let $b_n = \frac{1}{n}$ with $b_n \neq 0$ for all $n \in \mathbb{N}$. However, $(1/b_n)$ diverges.
- 3. Let $a_n = \frac{1}{n}$ and $b_n = n$. a_n converges to 0 and b_n diverges. $(a_n b_n)$ converges to 1.