

Lecture 21: Equilibrium figures

David Al-Attar

Michaelmas Term, 2026

Outline and motivation

In the previous lecture we obtained the equations of motion for a self-gravitating and rotating earth model, and showed that these effects are relevant at sufficiently long periods. A particular conclusion was that the equilibrium state of such an earth model **cannot be stress-free** as has been assumed in earlier parts of the course. In this lecture we investigate the constraints imposed by the equilibrium equations in further detail, and show that they are key to understanding the **observed shapes of planets**.

The equilibrium equations

In the previous lecture we obtained the following equilibrium equations

$$\begin{aligned} \rho \epsilon_{ijk} \epsilon_{klm} \Omega_j \Omega_l x_m - \frac{\partial T_{ij}^0}{\partial x_j} - \rho \gamma_i^0 &= 0, \\ \rho \epsilon_{ijk} \epsilon_{klm} \Omega_j \Omega_l x_m - \frac{\partial T_{ij}^0}{\partial x_j} - \rho \gamma_i^0 &= 0, \end{aligned} \tag{1}$$

for a self-gravitating and steadily rotating planet¹. Here ρ is the density, Ω_i the angular velocity, T_{ij}^0 the stress, and γ_i^0 the gravitational acceleration. To simplify ~~notations~~notation, let us write σ_{ij} for the equilibrium stress, and g_i for the gravitational acceleration. The equilibrium equations then become

$$\begin{aligned} \rho \epsilon_{ijk} \epsilon_{klm} \Omega_j \Omega_l x_m - \frac{\partial \sigma_{ij}}{\partial x_j} - \rho g_i &= 0, \\ \rho \epsilon_{ijk} \epsilon_{klm} \Omega_j \Omega_l x_m - \frac{\partial \sigma_{ij}}{\partial x_j} - \rho g_i &= 0, \end{aligned} \tag{2}$$

¹ ~~Though our focus in this course is on the Earth, the material of this lecture applies so naturally to other planets it seems sensible to generalise the discussion somewhat.~~

¹ Though our focus in this course is on the Earth, the material of this lecture applies so naturally to other planets it seems sensible to generalise the discussion somewhat.

which are subject to the boundary conditions $\sigma_{ij}\hat{n}_j = 0$ on ∂M . Though these equations were derived from elasticity, the force balance expressed is valid in more general materials. This is important because over long ~~time-scales~~ time scales the Earth or Earth-like planets are not elastic.

If we knew the shape, density distribution, and angular velocity of the planet, then all terms in the equilibrium equations ~~are~~ would be fixed except for the stress tensor. Recalling from Lecture ~~13-14~~ 13-14 that equilibrium stress tensors are necessarily symmetric, we see that we have only three equations for six unknown functions. The problem is, therefore, ~~under-determined~~ underdetermined, and we cannot expect to uniquely determine ~~stress-tensor~~² the stress tensor.²

Hydrostatic equilibrium

You have seen in the first part of this course that over thousands of years the interior of the Earth flows like a fluid, this being due to solid-state creep. Considering for simplicity a Newtonian fluid, we know that the stress tensor³ takes the form

$$\sigma_{ij} = -p \delta_{ij} + \eta \left(\frac{\partial v_i}{\partial x_j} + \frac{\partial v_j}{\partial x_i} - \frac{2}{3} \frac{\partial v_k}{\partial x_k} \delta_{ij} \right)$$

$$\sigma_{ij} = -p \delta_{ij} + \eta \left(\frac{\partial v_i}{\partial x_j} + \frac{\partial v_j}{\partial x_i} - \frac{2}{3} \frac{\partial v_k}{\partial x_k} \delta_{ij} \right), \quad (3)$$

where p is the pressure, η the viscosity, and v_i the velocity. Note that the given expression is valid in a **compressible fluid**, but it reduces to that for an incompressible one by requiring the velocity to be divergence-free. If the fluid is not flowing, then the stress tensor takes the simple hydrostatic form

$$\sigma_{ij} = -p \delta_{ij}$$

$$\sigma_{ij} = -p \delta_{ij}, \quad (4)$$

and the same is true for non-Newtonian fluids. Such an expression must also be approximately valid if the fluid is flowing sufficiently slowly. To get a sense of this approximation for the Earth, we note that the size of the non-hydrostatic stress associated with mantle convection scales like $\eta V/L$ with $\eta \sim 10^{21} \text{ Pa s}$ $\eta \sim 10^{21} \text{ Pa s}$ a typical

² ~~This counting argument is suggestive but not entirely correct. Rather, the relevant point is that there exist non-zero symmetric tensor fields having vanishing divergence. This can be compared to divergence-free vector fields with which you will be very familiar (i.e. divergence of a curl is zero). In fact, a method for generating divergence-free tensor fields exists using an extension of the curl operator that generates divergence-free symmetric tensor fields.~~

² This counting argument is suggestive but not entirely correct. Rather, the relevant point is that there exist non-zero symmetric tensor fields having vanishing divergence. This can be compared to divergence-free vector fields with which you will be very familiar (i.e. the divergence of a curl is zero). In fact, a method for generating divergence-free tensor fields exists using an extension of the curl operator that generates divergence-free symmetric tensor fields.

³ This is the Cauchy stress that occurs within spatial formulations of continuum mechanics. But note that at equilibrium the referential and spatial pictures coincide due to our choice of reference body.

viscosity, $V \sim 10^{-8} \text{ m s}^{-1}$ $V \sim 10^{-8} \text{ m s}^{-1}$ a typical speed of a few centimetres per year, and $L \sim 10^6 \text{ m}$ $L \sim 10^6 \text{ m}$ the length scale. From this we find that deviatoric stresses are around $\eta V/L \sim 10^7 \text{ Pa}$ $\eta V/L \sim 10^7 \text{ Pa}$. Pressures in the mantle are, however, known to be of order 10^{10} Pa 10^{10} Pa ,⁴ and hence the stress is quite close to being hydrostatic.

This motivates us to look for **hydrostatic** solutions of the equilibrium equations. Putting eq.(4) into eq.(2) we obtain

$$\rho \epsilon_{ijk} \epsilon_{klm} \Omega_j \Omega_l x_m + \frac{\partial p}{\partial x_i} - \rho g_i = 0.$$

$$\rho \epsilon_{ijk} \epsilon_{klm} \Omega_j \Omega_l x_m + \frac{\partial p}{\partial x_i} - \rho g_i = 0, \quad (5)$$

along with the boundary condition $p = 0$ on ∂M . In assuming the stress is hydrostatic we have moved from an **under-determined** **underdetermined** problem to an **over-determined** **overdetermined** one, with three equations for a single unknown, p . The form of this hydrostatic equilibrium equation can be usefully simplified by noting first that

$$g_i = - \frac{\partial \phi}{\partial x_i}$$

$$g_i = - \frac{\partial \phi}{\partial x_i}, \quad (6)$$

with ϕ the equilibrium gravitational potential. Similarly, the centrifugal term can also be written as the gradient of a potential

$$\epsilon_{ijk} \epsilon_{klm} \Omega_j \Omega_l x_m = \frac{\partial \psi}{\partial x_i}$$

$$\epsilon_{ijk} \epsilon_{klm} \Omega_j \Omega_l x_m = \frac{\partial \psi}{\partial x_i}, \quad (7)$$

where we have defined

$$\psi = \frac{1}{2} (\Omega_i \Omega_j - \Omega_k \Omega_k \delta_{ij}) x_i x_j$$

$$\psi = \frac{1}{2} (\Omega_i \Omega_j - \Omega_k \Omega_k \delta_{ij}) x_i x_j. \quad (8)$$

Moving to vector notation, the hydrostatic equilibrium equation takes the simple form

$$\nabla p + \rho \nabla \gamma = \mathbf{0},$$

$$\nabla p + \rho \nabla \gamma = \mathbf{0}, \quad (9)$$

⁴ Values for the pressure can be calculated once the density profile is known.

⁴ Values for the pressure can be calculated once the density profile is known.

where we have introduced the **gravity potential**, $\gamma = \phi + \psi$. Note that “gravity” is used here for the **combined effect of gravitational and centrifugal forces**.

An immediate consequence of eq.(9) is that ∇p and $\nabla \gamma$ are parallel, and hence the level surfaces of these fields are coincident. In particular, pressure vanishes on the surface of the planet, and so this must also be an equipotential of gravity. If we take the curl of eq.(9), recalling $\nabla \times \nabla f = 0$ for any scalar field, we find

$$\begin{aligned} \nabla \rho \times \nabla \gamma &= 0, \\ \nabla \rho \times \nabla \gamma &= 0, \end{aligned} \quad (10)$$

and so ~~note that~~ ρ and γ also have coincident level surfaces. In conclusion we have seen that for ~~an a~~ planet to be in hydrostatic equilibrium the level surfaces of p , ρ , and γ must all be the same.

Suppose we arbitrarily specified the shape, density variations, and angular velocity of ~~an a~~ planet. We could then, in principle, determine its gravitational potential by solving Poisson’s equation, and hence find the gravity potential. There would be no reason, however, that this field would have level surfaces that agree with those for the chosen density. This suggests that in requiring a planet to be in hydrostatic equilibrium we are placing rather strong restrictions on, and interrelations between, its shape, density distribution, and angular velocity.

Non-rotating hydrostatic planets

As a starting point, suppose that the planet does not rotate, and hence $\psi = 0$. Assume also that the planet is spherical in shape, and that density varies only with radius. We then know that ϕ is also spherically symmetric, and hence eq.(9) reduces to

$$\begin{aligned} \nabla p + \rho \frac{d\phi}{dr} \hat{r} &= 0, \\ \nabla p + \rho \frac{d\phi}{dr} \hat{r} &= 0, \end{aligned} \quad (11)$$

with $\hat{r}$ a radial unit vector. To solve this equation we can clearly take p to be a function of r alone, and hence need only integrate

$$\begin{aligned} \frac{dp}{dr} + \rho \frac{d\phi}{dr} &= 0, \\ \frac{dp}{dr} + \rho \frac{d\phi}{dr} &= 0, \end{aligned} \quad (12)$$

using the boundary condition $p = 0$ on the surface $r = b$. This argument shows that for a non-rotating ~~planets planet~~ it is possible to find at least one class of hydrostatic equilibrium states, namely those that are spherically symmetric. From a physical perspective, it seems reasonable that these are the only hydrostatic equilibria possible. This is indeed true, but a mathematical proof of this result is not easy⁵.

⁵ ~~In fact, I am aware of a proof only in the case of a homogeneous planet, with the argument summarised in Chandrasekhar’s book “Ellipsoidal figures of equilibrium”.~~

⁵ In fact, I am aware of a proof only in the case of a homogeneous planet, with the argument summarised in the book by Chandrasekhar (1969).

Slowly rotating hydrostatic planets

Given that a non-rotating and hydrostatic planet must be spherically symmetric, we would expect that a slowly rotating planet should have an equilibrium figure that is close to spherical. In this section, we show that this is indeed the case. First, however, we should clarify what is meant by **slowly rotating**.

The essential idea is that the ratio of the centrifugal to gravitational potentials should be small. Looking at these quantities on the planet's surface we have approximately

$$\begin{aligned} \psi &\sim \frac{1}{2}\Omega^2 b^2 \\ \psi &\sim \frac{1}{2}\Omega^2 b^2, \end{aligned} \quad (13)$$

where Ω is the angular frequency and b the mean radius, while also

$$\begin{aligned} \phi &\sim \frac{4}{3}\pi G \bar{\rho} b^2 \\ \phi &\sim \frac{4}{3}\pi G \bar{\rho} b^2, \end{aligned} \quad (14)$$

with $\bar{\rho}$ the mean density. The ratio of these terms is given by

$$\begin{aligned} \frac{\text{centrifugal potential}}{\text{gravitational potential}} &\sim \frac{3\Omega^2}{8\pi G \bar{\rho}} \\ \frac{\text{centrifugal potential}}{\text{gravitational potential}} &\sim \frac{3\Omega^2}{8\pi G \bar{\rho}}. \end{aligned} \quad (15)$$

For the Earth, we have $\bar{\rho} \sim 5000 \text{ kg m}^{-3}$ and $\Omega \sim 7 \times 10^{-5} \text{ s}^{-1}$, and hence we find

$$\begin{aligned} \frac{\text{centrifugal potential}}{\text{gravitational potential}} &\sim 2 \times 10^{-3}, \\ \frac{\text{centrifugal potential}}{\text{gravitational potential}} &\sim 2 \times 10^{-3}, \end{aligned} \quad (16)$$

which is indeed a fairly small quantity.

We model the rotating planet through a first-order perturbation of a spherical reference model. The density in the reference model will be written ρ_0 , and similarly for other quantities. Within the rotating planet we then have

$$\begin{aligned} \rho &= \rho_0 + s \rho_1 + \dots, \\ p &= p_0 + s p_1 + \dots, \\ \phi &= \phi_0 + s \phi_1 + \dots, \\ \psi &= s \psi_1, \end{aligned}$$

$$\underline{\rho = \rho_0 + s \rho_1 + \dots}, \quad (17)$$

$$\underline{p = p_0 + s p_1 + \dots}, \quad (18)$$

$$\underline{\phi = \phi_0 + s \phi_1 + \dots}, \quad (19)$$

$$\underline{\psi = s \psi_1}, \quad (20)$$

with s a perturbation parameter introduced for book-keeping purposes. We must also consider the aspherical shape of the planet. For simplicity we neglect internal boundaries^{6,6} and so need only parameterise the outer surface in the form

$$\underline{r(\theta, \varphi) = b + s h_1(\theta, \varphi) + \dots}, \quad (21)$$

with (θ, φ) spherical polar co-ordinates.

Putting the above expansions into the equilibrium equations we find at ~~first-order~~

$$\underline{\nabla p_1 + \rho_0 \nabla \gamma_1 + \rho_1 g_0 \hat{\mathbf{r}} = \mathbf{0}},$$

first order

$$\underline{\nabla p_1 + \rho_0 \nabla \gamma_1 + \rho_1 g_0 \hat{\mathbf{r}} = \mathbf{0}}, \quad (22)$$

where $\gamma_1 = \phi_1 + \psi_1$, and we have written $g_0 = \frac{d\phi_0}{dr}$ for convenience. The boundary condition on pressure is applied on the perturbed surface, and hence

$$\underline{p[b + s h_1(\theta, \varphi) + \dots, \theta, \varphi] = 0.}$$

$$\underline{p[b + s h_1(\theta, \varphi) + \dots, \theta, \varphi] = 0.} \quad (23)$$

Expanding this out to ~~first-order~~ first order in s , we arrive at

$$\underline{p_1 + \frac{dp_0}{dr} h_1 = 0}$$

$$\underline{p_1 + \frac{dp_0}{dr} h_1 = 0} \quad (24)$$

on the reference surface $r = b$. The zeroth-order equilibrium equation tells us that

$$\underline{\frac{dp_0}{dr} + \rho_0 g_0 = 0},$$

$$\underline{\frac{dp_0}{dr} + \rho_0 g_0 = 0}, \quad (25)$$

⁶ ~~But these can be easily included into the theory.~~

⁶ But these can be easily included in the theory.

and hence the boundary ~~conditions simplify to~~

$$\underline{p_1 - \rho_0 g_0 h_1 = 0.}$$

condition simplifies to

$$\underline{p_1 - \rho_0 g_0 h_1 = 0.} \quad (26)$$

We will show that ~~eq.(22)~~eqs. (22) and ~~(26)~~(26) allow the density, pressure, and boundary perturbations to be expressed in terms of that for the gravity potential. First, we take the cross product of ~~eq.(22)~~eq. (22) with the unit vector $\hat{\mathbf{r}}$ to obtain

$$\underline{\hat{\mathbf{r}} \times \nabla p_1 + \hat{\mathbf{r}} \times \rho_0 \nabla \gamma_1 = \mathbf{0}.$$

$$\underline{\hat{\mathbf{r}} \times \nabla p_1 + \hat{\mathbf{r}} \times \rho_0 \nabla \gamma_1 = \mathbf{0}. \quad (27)$$

Noting that

$$\underline{\hat{\mathbf{r}} \times \rho_0 \nabla \gamma_1 = \hat{\mathbf{r}} \times \left[\nabla(\rho_0 \gamma_1) - \gamma_1 \frac{d\rho_0}{dr} \hat{\mathbf{r}} \right] = \hat{\mathbf{r}} \times \nabla(\rho_0 \gamma_1),}$$

$$\underline{\hat{\mathbf{r}} \times \rho_0 \nabla \gamma_1 = \hat{\mathbf{r}} \times \left[\nabla(\rho_0 \gamma_1) - \gamma_1 \frac{d\rho_0}{dr} \hat{\mathbf{r}} \right] = \hat{\mathbf{r}} \times \nabla(\rho_0 \gamma_1), \quad (28)$$

eq.~~(27)~~(27) can be written

$$\underline{\hat{\mathbf{r}} \times \nabla(p_1 + \rho_0 \gamma_1) = \mathbf{0}.$$

$$\underline{\hat{\mathbf{r}} \times \nabla(p_1 + \rho_0 \gamma_1) = \mathbf{0}. \quad (29)$$

This relation tells us that the gradient of $p_1 + \rho_0 \gamma_1$ is parallel to $\hat{\mathbf{r}}$, and hence this function depends on radius alone. We are free to assume that the perturbations all average to zero over spherical surfaces^{7,7} and hence we arrive at

$$\underline{p_1 = -\rho_0 \gamma_1}$$

$$\underline{p_1 = -\rho_0 \gamma_1,} \quad (30)$$

which expresses the pressure perturbation in terms of that to the gravity potential. From the perturbed boundary condition we then immediately obtain

$$\underline{h_1 = -\frac{\gamma_1}{g_0}}$$

$$\underline{h_1 = -\frac{\gamma_1}{g_0}.} \quad (31)$$

⁷ ~~If not, these terms could be combined into the spherical reference model.~~

⁷ If not, these terms could be combined into the spherical reference model.

For the second stage of the argument, we take the curl of eq. (22) to obtain

$$\begin{aligned} \frac{d\rho_0}{dr} \hat{\mathbf{r}} \times \nabla \gamma_1 + g_0 \nabla \rho_1 \times \hat{\mathbf{r}} &= \mathbf{0}. \\ \frac{d\rho_0}{dr} \hat{\mathbf{r}} \times \nabla \gamma_1 + g_0 \nabla \rho_1 \times \hat{\mathbf{r}} &= \mathbf{0}. \end{aligned} \quad (32)$$

Using the same trick as before, this is equivalent to

$$\begin{aligned} \hat{\mathbf{r}} \times \nabla \left(\frac{d\rho_0}{dr} \gamma_1 - g_0 \rho_1 \right) &= \mathbf{0}, \\ \hat{\mathbf{r}} \times \nabla \left(\frac{d\rho_0}{dr} \gamma_1 - g_0 \rho_1 \right) &= \mathbf{0}, \end{aligned} \quad (33)$$

from which we see that the density perturbation is given by

$$\begin{aligned} \rho_1 &= \frac{1}{g_0} \frac{d\rho_0}{dr} \gamma_1 \\ \rho_1 &= \frac{1}{g_0} \frac{d\rho_0}{dr} \gamma_1. \end{aligned} \quad (34)$$

To summarise, we started from the hydrostatic equilibrium equations in a slowly rotating planet, and have obtained the following identities

$$\begin{aligned} p_1 &= -\rho_0 \gamma_1, \quad h_1 = -\frac{\gamma_1}{g_0}, \quad \rho_1 = \frac{1}{g_0} \frac{d\rho_0}{dr} \gamma_1. \\ p_1 &= -\rho_0 \gamma_1, \quad h_1 = -\frac{\gamma_1}{g_0}, \quad \rho_1 = \frac{1}{g_0} \frac{d\rho_0}{dr} \gamma_1. \end{aligned} \quad (35)$$

To complete the discussion, we recall that ϕ_1 is also related to ρ_1 and h_1 through solution of Poisson's equation. Indeed, we have

$$\begin{aligned} \nabla^2 \phi_1 &= \begin{cases} 4\pi G \rho_1 & r < b \\ 0 & r > b \end{cases} \\ \nabla^2 \phi_1 &= \begin{cases} 4\pi G \rho_1 & r < b \\ 0 & r > b \end{cases} \end{aligned} \quad (36)$$

while the appropriate continuity conditions at $r = b$ are

$$\begin{aligned} [\phi_1]_{-}^{+} &= 0, \quad [\hat{\mathbf{r}} \cdot \nabla \phi_1 + 4\pi G \rho_0 h_1]_{-}^{+} = 0 \\ [\phi_1]_{-}^{+} &= 0, \quad [\hat{\mathbf{r}} \cdot \nabla \phi_1 + 4\pi G \rho_0 h_1]_{-}^{+} = 0, \end{aligned} \quad (37)$$

whose derivation is left as an exercise in the second problem set. Using eq.(35 (35)), we see that Poisson's equation is transformed into

$$\nabla^2 \phi_1 = \begin{cases} 4\pi G \frac{1}{g_0} \frac{d\rho_0}{dr} (\phi_1 + \psi_1) & r < b \\ 0 & r > b \end{cases}$$

$$\nabla^2 \phi_1 = \begin{cases} 4\pi G \frac{1}{g_0} \frac{d\rho_0}{dr} (\phi_1 + \psi_1) & r < b \\ 0 & r > b \end{cases} \quad (38)$$

along with the continuity conditions

$$[\phi_1]_{-}^{+} = 0, \quad \left[\hat{\mathbf{r}} \cdot \nabla \phi_1 - 4\pi G \rho_0 \frac{(\phi_1 + \psi_1)}{g_0} \right]_{-}^{+} = 0.$$

$$[\phi_1]_{-}^{+} = 0, \quad \left[\hat{\mathbf{r}} \cdot \nabla \phi_1 - 4\pi G \rho_0 \frac{(\phi_1 + \psi_1)}{g_0} \right]_{-}^{+} = 0. \quad (39)$$

Eq.(38 Eqs. (38) and (39 39) constitute a boundary value problem that can be solved for ϕ_1 in terms of ψ_1 . Having done this we can then use eq.(35 (35)) to determine the other parameters for the perturbed planet. These equations, albeit in a less general form, were first obtained by Clairaut (1713-1765 1713-1765), this work building on Newton's original discussion of equilibrium figures in the "*Principia*" *Principia* (Newton 1687). To implement these equations in practice we must know the variation of spherically averaged density with radius, and in later lectures we will see how this was determined for the Earth.

It is worth noting these arguments apply to more general potentials, ψ_1 , and not only that linked to the centrifugal force. The theory can, therefore, also be applied to model things like equilibrium tides within gaseous planets. Another point to make is that the equations are linear in ϕ_1 and ψ_1 , and hence symmetries of ψ_1 are directly mirrored in ϕ_1 and the other perturbed quantities. In terms of spherical harmonics, the aspherical part of the centrifugal potential perturbation can be shown to be proportional to Y_{20} . It is for this reason that equilibrium figures of slowly rotating planets are found to be oblate spheroids, and characterised in terms of a single radial function, ϵ , known as the ellipticity and defined such that

$$\phi_1 + \psi_1 = \sqrt{\frac{4\pi}{5}} \frac{2}{3} r g_0 \epsilon Y_{20}.$$

$$\phi_1 + \psi_1 = \sqrt{\frac{4\pi}{5}} \frac{2}{3} r g_0 \epsilon Y_{20}. \quad (40)$$

At the mean surface $r = b$ it can be shown that the ellipticity is such that

$$\epsilon(b) = \frac{\text{equatorial radius} - \text{polar radius}}{\text{mean radius}}$$




Fig. 1: The upper panel shows variations in the Earth's spherically averaged density structure as determined from free oscillation data along with mass and moment of inertia constraints. The lower figure panel then plots the reciprocal of the ellipticity as determined through numerical solution of Clairaut's equation.

$$\epsilon(b) = \frac{\text{equatorial radius} - \text{polar radius}}{\text{mean radius}}, \quad (41)$$

and for the Earth this value is around 1/300. Using properties of spherical harmonics, it can be shown that eq.(38) reduces to

$$\frac{d^2\epsilon}{dr^2} + 8\pi G\rho_0 g_0^{-1} \left(\frac{d\epsilon}{dr} + \frac{\epsilon}{r} \right) - \frac{6\epsilon}{r^2} = 0. \quad (42)$$

which is known as Clairaut's equation and is subject to the boundary conditions

$$\left. \frac{d\epsilon}{dr} \right|_{r=0} = 0, \quad \left. \frac{d\epsilon}{dr} \right|_{r=b} = \frac{1}{b} \left[\frac{5\Omega^2 b^3}{2GM} - 2\epsilon(b) \right]$$

$$\left. \frac{d\epsilon}{dr} \right|_{r=0} = 0, \quad \left. \frac{d\epsilon}{dr} \right|_{r=b} = \frac{1}{b} \left[\frac{5\Omega^2 b^3}{2GM} - 2\epsilon(b) \right], \quad (43)$$

with M the planet's mass. This problem can be solved numerically using a shooting method, whereby the initial conditions at $r = 0$ are progressively adjusted so that the required condition at $r = b$ is met.

What you need to know and be able to do

- (i) Know the form of the equilibrium equations, and that they hold in a general planet and not just an elastic one.
- (ii) That the equilibrium equations are ~~under-determined, but they admit non-unique solutions subject to the two compatibility conditions~~ underdetermined, and why.
- (iii) What a hydrostatic equilibrium state is, and be able to derive the constraints imposed on the level surfaces of p , ρ , and γ .
- (iv) Derive the hydrostatic equilibrium state in a non-rotating planet.
- (v) Know in outline how solutions can be obtained within a slowly rotating planet. Here you will not need to reproduce the full derivation, but may be asked to perform related calculations with appropriate guidance.
- (vi) ~~How minimum stress fields can be defined and their relation to slow viscous flow. Again, you do not need to learn this derivation in full, but may be asked to perform related calculations given the necessary information.~~