

Lecture 0.3: Review of Trigonometry and Graphing Trigonometric Functions

Professor Leonard

December 22, 2022

1 Angles

Plotting a vector and making an angle relative to the x-axis. The x-axis is called the "initial side", the vector line is called the "terminal side".

Counter clockwise rotation from the x-axis gives positive angles. Clockwise gives negative angles.

1.1 Degrees vs. radians

Radians involve π .

Covertng radians to degrees:

$$2\pi = 360^\circ$$
$$f(x^\circ) = x^\circ \cdot \frac{\pi}{180^\circ}$$
$$g(r) = r \cdot \frac{180^\circ}{\pi}$$

$f(x^\circ)$ converts degrees to radians. $g(r)$ converts radians to degrees.

He's going through some plotting of radian angles... not sure how to take notes on that. A bit tricky, but if you have a radian (eg, $5\pi/2$), then you take the four cartesian quadrants, chop up the top and bottom halves equally by the denominator. The terminals of the four axes end points are as follows (+x: 0 or 2π , +y: $\pi/2$, -x: π , -y: $3\pi/2$). And you then take the numerator coefficient and count that many either counter clockwise (for positive numerator) or clockwise (for negative numerator).

2 Trigonometric Functions

Sine, cosine, and tangent are *functions*. They need an input (usually an angle). Sine, cosine, and tangent can be used to find the angles of a right triangle

(remember SohCahToa: sine, opposite over adjacent; cosine, adjacent over hypotenuse, tangent, opposite over adjacent).

There's also the reciprocals of the three trig functions.

$$\begin{aligned}\text{Secant: } \sec(\theta) &= \frac{1}{\cos(\theta)} = \frac{Hyp}{Adj} \\ \text{Cosecant: } \csc(\theta) &= \frac{1}{\sin(\theta)} = \frac{Hyp}{Opp} \\ \text{Cotangent: } \cot(\theta) &= \frac{1}{\tan(\theta)} = \frac{Adj}{Opp}\end{aligned}$$

For the unit circle (ie, circle with radius of 1), for any point (x, y) along the circumference of the circle, $\cos(\theta) = x$, $\sin(\theta) = y$, $\tan(\theta) = y/x = \sin(\theta)/\cos(\theta)$. And the reciprocal functions are just the reciprocals of these outcomes.

2.1 Trigonometric functions across the quadrants

Whether the trig functions are positive or negative depends on the quadrant. The mnemonic "All Students Take Calculus" → each letter iterates through cartesian quadrants I:IV, A=All functions are positive in quadrant I, S=Sine (and cosecant) is positive in quadrant II, T=Tangent (and cotangent) is positive in quadrant III, C=Cosine (secant) is positive in quadrant IV.

2.2 Reference angles

The idea is to make an acute angle with the x-axis, find the trig function of that, then use the ASTC idea.

If the angle is already acute relative to x-axis, just take theta. Otherwise, find the angle with x that makes it acute. So for example, if $\pi > \theta > 0$, then you calculate the acute angle by $\pi - \theta$. Similar idea for quadrants III & IV.

2.3 Trig function for any angle

One can use the properties of the functions across the quadrants and the reference angles, along with knowledge of the unit circle, to determine for any input angle the output of any trig function.

Example: find sin, cos, tan of $5\pi/3$.

Steps:

1. Locate the quadrant.
I think it's Quadrant IV

- Find the reference angle.

$$2\pi - 5\pi/3 = -3\pi/3$$

Oh, fractions need a common denominator when adding or subtracting...


$$6\pi/3 - 5\pi/3 = \pi/3$$

- Find all trig functions of the reference angle.

I guess this is where we use the unit circle.¹

3 Graphic Trigonometric Functions

We're going to graph things like this: $y = A \cdot \sin(Bx)$ or $y = A \cdot \cos(Bx)$. To appreciate the impact of the coefficients, we'll compare $y = \sin(x)$ against $y = 2 \cdot \sin(4x)$.



The period of a wave function is the stretch over which it's unique (ie, before it repeats).

Left off at 47:39.

¹I had to take a little detour into the unit circle. Apparently the x y coordinates of the so-called "special angles/trianles" (30-60-90 and 45-45-90) are just givens and expressed as fractions of roots (so unsatisfying). So if want to find $\sin(\pi/3)$ I have to go look up that the coordinates for the unit circle at $\theta = \pi/3$ are $(1/2, \sqrt{3}/2)$. Then obviously $\sin(\pi/3) = y = \sqrt{3}/2$. And then you determine the sign of that by the quadrant.