Quantum Algorithms Using Clebsch-Gordan Transforms



Dave Bacon
University of Washington

|QW⟩ group

Quantum Computers

188198812920607963838697239461650439807163563379417382700763356422988859715234665485319060606504743045317388011303396716199692321205734031879550656996221305168759307650257059

=

3980750864240649373971255005503864911990643623425267084063851895759463889572617685833174727721461074353025362230719730482246329146953020971164598521711305207112563636590397527

×



Peter Shor

1994

- Quantum computers can efficiently factor whole numbers (and compute discrete log.)
- No known classical efficient algorithm.
- Breaks RSA, Diffie-Hellman cryptosystems.


Dr. Quantum Love

“Only time will tell if and when the problems of building a quantum computer can be overcome…. the development of a fully operational quantum computer would imperil our personal privacy, destroy electronic commerce and demolish the concept of national security. **A quantum computer would jeopardize the stability of the world.** Whichever country gets there first will have the ability to monitor the communications of its citizens, read the minds of its commercial rivals and eavesdrop on the plans of its enemies. Although it is still in its infancy, quantum computing presents a potential threat to the individual, to international business and to global security.”



-Simon Singh


Physicists and Computer Scientists



A Golden Era

Email I received from Wim van Dam about FOCS 2006:

Leslie Valiant in his great talk on 'accidental algorithms': "We have all these complexity classes, and we don't know at all if they re if they turn out to and so on." <comment frc lly be so bad> "Yes it possible nightmare wc hysicist* who proved i







On Quantum Algorithms



Ronald de Wolf
(CWI)

From a 2003 review of three quantum computing books:

"Finally, to all those currently working on yet another book on quantum computing: what this field needs most is more algorithms, not more books."



My Nightmare

HELP WANTED Quantum programmer. Only skill necessary is ability to type in large RSA keys. Pay inversely commensurate with years since Ph.D. completed.

HELP WANTED Real estate agent. Best market since bubble popped in 07. Join a dedicated

The State of Quantum Algorithms

Grover-Type

Needle in Haystack Search
[Grover 96]

Quantum amplitude amplification
[Brassard, Hoyer, Mosca, Tapp 00]

Adiabatic quantum algorithms
[Farhi, Goldstone Gutman, Sipser 00]

NAND tree algorithm
[Farhi, Goldstone Gutman 07]

Quantum simulated annealing
[Somma, Boixo, Barnum 07]

BQP completeness of Jones, Tutte, and other polynomials, Exponential speedups in quantum random walks, Simulation of Schrodinger's equation, Physical chemistry, etc.

Shor-Type

Factoring, Discrete Logarithm
[Shor 94]

Pell's Equation
[Hallgren 02]

Some Hidden Shift Problems
[van Dam, Hallgren, Ip 02]
[van Dam, Childs 05]

Gauss Sums
[van Dam, Seroussi 02]

Hidden Subgroup Problems
Abelian groups, almost abelian groups, near Hamiltonian groups, normal subgroups, small commutator subgroup, solvable groups of bounded exponent and of bounded derived series extraspecial groups, groups of nilpotency class at most 2, more

The Power of Quantum Computers


Clichéd Question: Where does the power of quantum computers come from?

?????

Retort: Where does the power of classical computers come from?



= powerful exactly why?



Outline

1. Lessons from a simple quantum algorithm
2. Lessons from the symmetric and unitary groups
3. When at first you don't succeed…..symmeterize
4. Quantum algorithms from Clebsch-Gordan transforms

Simon's Problem

(is that nobody does what Simon says)

Given: A function with n bit strings as input and one bit as output


$$f : x \in \{0,1\}^n \rightarrow \{0,1\}^n$$

Promise: The function is guaranteed to satisfy

$$f(x) = f(y) \Leftrightarrow y = x \oplus s, s \neq 0$$


$$x \oplus s = (x_1 \oplus s_1, x_2 \oplus s_2, \ldots, x_n \oplus s_n)$$

Problem: Find the n bit string $s \neq 0$



Classical Simon's Problem

query pairs $(x_\alpha, f(x_\alpha))$



If we find $x_\alpha \neq x_\beta$ such that $f(x_\alpha) = f(x_\beta)$ then

$$f(x_\alpha) = f(x_\beta) \Rightarrow x_\alpha = s \oplus x_\beta$$


$$s = x_\alpha \oplus x_\beta$$

Querying the function *m* times….

Probability of getting a matching pair $\approx \frac{\binom{m}{2}}{2^n} = \frac{m(m-1)}{2^{n+1}}$

Bounded error query complexity: $m = O\left(2^{\frac{n}{2}}\right)$

Quantum Simon's Problem



$$H = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$$

$$\frac{1}{\sqrt{2^n}} \sum_{x=0}^{2^n-1} |x\rangle \otimes |f(x)\rangle$$

Quantum Simon's Problem

For random uniformly distributed $x \in \{0,1\}^n$

$$\frac{1}{\sqrt{2^n}} \sum_{x=0}^{2^n-1} |x\rangle \otimes |f(x)\rangle \longrightarrow |\psi_s(x)\rangle = \frac{1}{\sqrt{2}}(|x\rangle + |x \oplus s\rangle)$$

$$\rho_s = \frac{1}{2^n} \sum_x |\psi_s(x)\rangle\langle\psi_s(x)|$$

Symmetry:

$$R = X^{a_1} \otimes X^{a_2} \otimes \cdots X^{a_n} \qquad a_i \in \{0,1\}$$

$$R\rho_s R^\dagger = \rho_s \quad X = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

Symmetry → Simultaneously diagonalize symmetry operators


and symmetri[illegible]

$$HZH^\dagger = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$$

Quantum Simon's Problem

n qubits

n qubits



$$H^{\otimes n}\frac{1}{\sqrt{2}}(|x\rangle+|x\oplus s\rangle) = \frac{1}{\sqrt{2^{n+1}}}\sum_{y=0}^{2^n-1}((-1)^{y\cdot x}+(-1)^{y\cdot(x\oplus s)})|y\rangle$$

$$y\cdot x = y_1x_1 \oplus y_2x_2 \oplus \cdots \oplus y_nx_n$$

Quantum Simon's Problem

$$H^{\otimes n}\frac{1}{\sqrt{2}}(|x\rangle+|x\oplus s\rangle) = \frac{1}{\sqrt{2^{n+1}}}\sum_{y=0}^{2^n-1}((-1)^{y\cdot x}+(-1)^{y\cdot(x\oplus s)})|y\rangle$$

$$= \frac{1}{\sqrt{2^{n+1}}}\sum_{y=0}^{2^n-1}(-1)^{y\cdot x}(1+(-1)^{y\cdot s})|y\rangle$$

Measuring this state, uniformly distributed random y such that

$$y \cdot s = 0 \mod 2$$


Multiple runs:

$$\begin{aligned} y_1 \cdot s &= 0 \\ y_2 \cdot s &= 0 \\ &\vdots \\ y_k \cdot s &= 0 \end{aligned}$$

random uniform y_i

If we obtain $n-1$ linearly independent equations of this form,
we win (Gaussian elimination)

Moral



Symmetries

↓

Diagonalize with respect to symmetry

↓

Quantum algorithm

Quantum

$$S\rho S^\dagger = \rho$$

ρ and S simultaneously diagonalized

Classical

$$S\vec{p} = \vec{p}$$

What do we do?

But what about that Gaussian Elimination?

Representation Theory in 1 Slide

Group $\mathcal{G}$

Representation of group $T : \mathcal{G} \rightarrow U(d)$

Irreducible representations of a group (irreps):

T_γ has no invariant subspace except full space and empty subspace

Decomposition into irreps

$$T = \bigoplus_i T_{\gamma_i} = \bigoplus I_{n_\gamma} \otimes T_\gamma$$

Schur's lemma:

$$T\rho = \rho T \Rightarrow \rho = \bigoplus_\gamma q_\gamma \otimes I_{d_\gamma}$$

Simon's Problem Redux

Group $\mathcal{G} = \mathbb{Z}_2^n$ addition bitwise mod *2*

$$(x_1, \ldots, x_n) + (y_1, \ldots, y_n) = (x_1 + y_1, \ldots, x_n + y_n)$$

$$x_i, y_i \in \{0, 1\}$$

Representation of group

$$T((x_1, \ldots, x_n)) = X^{x_1} \otimes \cdots \otimes X^{x_2}$$

Irreps of group:

$$T_{(k_1,\ldots,k_n)}((x_1, \ldots, x_n)) = (-1)^{x_1 k_1 + \cdots + x_n k_n}$$

Unitary transform to direct product of irreps:

$$H^{\otimes n} T((x_1, \ldots, x_n)) H^{\otimes n\dagger} = \bigoplus_k T_{k_1,\ldots,k_n}((x_1, \ldots, x_n))$$

More Simon's Problem Redux

$$
\begin{aligned}
y_1 \cdot s &= 0 \\
y_2 \cdot s &= 0 \\
&\vdots \\
y_k \cdot s &= 0
\end{aligned}
$$

random uniform y_i

are irrep labels

Gaussian elimination : adding y_i bitwise mod *2*

Symmetry interpretation?

Symmetry Interpretation

Clebsch-Gordan transform:

$$T_\alpha(g) \otimes T_\beta(g) = \bigoplus_\gamma I_{n_\gamma} \otimes T_\gamma(g)$$

$$T_{(k_1,\ldots,k_n)}((x_1,\ldots,x_n)) \otimes T_{(l_1,\ldots,l_n)}((x_1,\ldots,x_n))$$

$$= (-1)^{x_1k_1+\cdots+x_nk_n}(-1)^{x_1l_1+\cdots+x_nl_n}$$

$$= T_{(k_1+l_1,\ldots,k_n+l_n)}((x_1,\ldots,x_n))$$

So the “addition” of Gaussian elimination in Simon’s algorithm can be interpreted as a Clebsch-Gordan transform….

The Second Basis Change We Learn In Quantum Mechanics:

Clebsch-Gordan Transform

Singlet:

$$|S=0,m=0\rangle = \frac{1}{\sqrt{2}}(|\uparrow\rangle \otimes |\downarrow\rangle - |\downarrow\rangle \otimes |\uparrow\rangle)$$

Triplet:

$$|S=1,m=1\rangle = |\uparrow\rangle \otimes |\uparrow\rangle$$

$$|S=1,m=0\rangle = \frac{1}{\sqrt{2}}(|\uparrow\rangle \otimes |\downarrow\rangle + |\downarrow\rangle \otimes |\uparrow\rangle)$$

$$|S=1,m=-1\rangle = |\downarrow\rangle \otimes |\downarrow\rangle$$

What Use?

Collective Unitaries


$$|S=0,m=0\rangle = \frac{1}{\sqrt{2}}(|\uparrow\rangle \otimes |\downarrow\rangle - |\downarrow\rangle \otimes |\uparrow\rangle)$$

$$|S=1,m=1\rangle = |\uparrow\rangle \otimes |\uparrow\rangle$$

$$|S=1,m=0\rangle = \frac{1}{\sqrt{2}}(|\uparrow\rangle \otimes |\downarrow\rangle + |\downarrow\rangle \otimes |\uparrow\rangle)$$


$$|S=1,m=-1\rangle = |\downarrow\rangle \otimes |\downarrow\rangle$$

$$\left[\begin{array}{c|c} 1 & 0 \\ \hline 0 & U \end{array}\right]$$




$U \otimes U$

Symmetry Under n Collective Unitary Rotations



$U^{\otimes n}$



The Symmetric Group

Transform Under SWAP

Singlet:

$$|S=0,m=0\rangle = \frac{1}{\sqrt{2}}(|\uparrow\rangle\otimes|\downarrow\rangle - |\downarrow\rangle\otimes|\uparrow\rangle) \qquad -1$$


Triplet:

$$|S=1,m=1\rangle = |\uparrow\rangle\otimes|\uparrow\rangle$$


$$|S=1,m=0\rangle = \frac{1}{\sqrt{2}}(|\uparrow\rangle\otimes|\downarrow\rangle + |\downarrow\rangle\otimes|\uparrow\rangle) \qquad +1$$

$$|S=1,m=-1\rangle = |\downarrow\rangle\otimes|\downarrow\rangle$$

Symmetry Under Permutation of n Subsystems




Quantum Schur Transform



Schur duality

$$[\mathbb{C}^d]^{\otimes n} \overbrace{\cong}^{\text{Schur}} \bigoplus_{\mathcal{Y}} S_{\mathcal{Y}} \otimes U_{\mathcal{Y}}$$

Young diagram




$$k_1 \geq k_2 \geq \cdots \geq k_d$$

$$\sum_{i=1}^{d} k_i = n$$

Uses of

Young diagram




$$k_1 \geq k_2 \geq \cdots \geq k_d$$

$$\sum_{i=1}^{d} k_i = n$$

What does this remind you of?

$\frac{k_i}{n}$ like probabilities like the spectrum of a density operator



[Keyl and Werner. PRA, 2001]

The Plethora of Uses of Schur

Optimal spectrum estimation

[M. Keyl and R. F. Werner. Phys. Rev. A, 2001]

Universal distortion-free entanglement concentration

[M. Hayashi and K. Matsumoto. Information Theory, 2004]

Universal compression with optimal overflow exponent

[M. Hayashi and K. Matsumoto. Phys. Rev. A, 2002]

Optimal quantum hypothesis testing

[M. Hayashi, J. Phys. A, 2002]

Communicating without a shared reference frame

[S. Bartlett, T. Rudolph, and R. Spekkens, Phys. Rev. Lett., 2003]

Encoding into decoherence-free (noiseless) subsystems

[J. Kempe, D. Bacon, D. A. Lidar, and K. B. Whaley, Phys. Rev. A, 2001]


Simulating lattice gauge theories on a quantum computer

[T. Byrnes and Y. Yoshihisa, Phys. Rev. A 2006]

….

But Efficient?

QFT



Schu
r



Who would need to efficiently calculate Clebsch-Gordan coefficients for SU(d)?

[L. C. Biedenharn and J. D. Louck, Commun. Math. Phys. 1968]

Efficient Schur

[D. Bacon, I.L. Chuang, A. Harrow, Phys. Rev. Lett. 2006]
[D. Bacon, I.L. Chuang, A. Harrow, SODA 2007]




FIG. 4: The $\mathcal{U}_d$ CG transform, $\mathbf{U}_{\mathrm{CG}}^{[d]}$, is decomposed into a $\mathcal{U}_{d-1}$ CG transform $\tilde{U}_{\mathrm{CG}}^{[d-1]}$ (see Eq. (54)) and a reduced Wigner operator $\hat{T}^{[d]}$. In Fig. 5 we show how to reduce the reduced Wigner operator to a $d \times d$ matrix conditioned on μ and $\mu' + e_{j'}$.




FIG. 5: The reduced Wigner transform $\hat{T}^{[d]}$ can be expressed as a $d \times d$ rotation whose coefficients are controlled by μ and $\mu' + e_{j'}$.

Schur transform to accuracy ϵ : time $poly\left(log\frac{1}{\epsilon}, n, d\right)$

The Plethora of Uses of Schur

Optimal spectrum estimation

[M. Keyl and R. F. Werner. Phys. Rev. A, 2001]

Universal distortion-free entanglement concentration

[M. Hayashi and K. Matsumoto. Information Theory, 2004]

Universal compression with optimal overflow exponent

[M. Hayashi and K. Matsumoto. Phys. Rev. A, 2002]

Optimal quantum hypothesis testing

[M. Hayashi, J. Phys. A, 2002]

Communicating without a shared reference frame

[S. Bartlett, T. Rudolph, and R. Spekkens, Phys. Rev. Lett., 2003]

Encoding into decoherence-free (noiseless) subsystems

[J. Kempe, D. Bacon, D. A. Lidar, and K. B. Whaley, Phys. Rev. A, 2001]

Simulating lattice gauge theories on a quantum computer

[T. Byrnes and Y. Yoshihisa, Phys. Rev. A 2006]

….

Tr

ac

ta

bl


But Quantum Algorithms?

The Hidden Subgroup Problem

Setup: Fix a group $\mathcal{G}$(known) and a subgroup $\mathcal{H} \subset \mathcal{G}$(unknown). Given query access to a function $f : \mathcal{G} \to S$ that is constant and distinct on left (right) cosets of $\mathcal{H}$ in $\mathcal{G}$ We say that f hides $\mathcal{H}$

Problem: Find a generating set for $\mathcal{H}$ ("identify $\mathcal{H}$").

Efficient: If runs in time polylog($|\mathcal{G}|$)





Hidden Subgroup Disease

“The quantum measurement problem refers to a set of people.”
- Hideo Mabuchi

“The hidden subgroup problem refers to a subgroup of people.”

Why the Fuss?





Solved Problems (Shor 1994)

efficient algorithm for HSP over $\mathbb{Z}_N$ → efficient algorithm for factoring whole numbers


efficient algorithm for HSP over $\mathbb{Z}_N \times \mathbb{Z}_N$ → efficient algorithm for the discrete logarithm problem

Open Problems

efficient algorithm for HSP over $\mathcal{S}_n \wr \mathbb{Z}_2$ (or $\mathcal{S}_{2n}$) → efficient algorithm for the graph isomorphism problem


efficient "standard" quantum algorithm for HSP over dihedral group $\mathcal{D}_N$ → efficient algorithm for the poly(*n*) unique shortest vector in a lattice problem

Graph Isomorphism



adjacency matrix A_1

$G_1 = (V_1, E_1)$



adjacency matrix A_2

$G_2 = (V_2, E_2)$

Problem: Is there a 1-to-1 mapping p from the vertices V_1 to the vertices V_2 such that the edges are respected, $p(E_1) = E_2$?


Mapping to the HSP:

$$f : \mathcal{S}_{2n} \rightarrow \mathbb{Z}_{n(2n-1)}$$

$$f(\pi) = \pi(A_1 \oplus A_2)$$

If graphs are not isomorphic, hidden subgroup will not mix the graphs. If they are isomorphic, they will mix the graphs.

Standard Query Model



Query in equal superposition over group elements

$$\frac{1}{\sqrt{|\mathcal{G}|}} \sum_{g \in \mathcal{G}} |g\rangle|0\rangle \rightarrow \sum_{g \in \mathcal{G}} \frac{1}{\sqrt{|\mathcal{G}|}} |g\rangle|f(g)\rangle$$

$$\rho_{\mathcal{H}} = \frac{|\mathcal{H}|}{|\mathcal{G}|} \sum_{g \in \hat{\mathcal{G}}} |g\mathcal{H}\rangle\langle g\mathcal{H}|$$ hidden subgroup state


You Down With Symmetry?

$$\rho_{\mathcal{H}} = \frac{|\mathcal{H}|}{|\mathcal{G}|} \sum_{g \in \hat{\mathcal{G}}} |g\mathcal{H}\rangle\langle g\mathcal{H}|$$ hidden subgroup state

Symmetry under left regular representation:

$$D_L(g)|g'\rangle = |gg'\rangle \qquad D_L(g)\rho_{\mathcal{H}} D_L(g^{-1}) = \rho_{\mathcal{H}}$$


Fourier transform over finite group $\mathcal{G}$



Can we solve HSP this way?

Multicopy






$\rho_{\mathcal{H}}^{\otimes m}$

Can we make a measurement on $m < O(\log |\mathcal{G}|)$ copies and learn the subgroup?

In general (and for important case of symmetric group) **NO**

[Moore, Russell, Hallgren, Roetteler, Sen 05]

Required:



You Down With Symmetry?

$$\rho_{\mathcal{H}} = \frac{|\mathcal{H}|}{|\mathcal{G}|} \sum_{g \in \hat{\mathcal{G}}} |g\mathcal{H}\rangle\langle g\mathcal{H}|$$ hidden subgroup state

Symmetry under left regular representation:

$$D_L(g)|g'\rangle = |gg'\rangle \qquad D_L(g)\rho_{\mathcal{H}} D_L(g^{-1}) = \rho_{\mathcal{H}}$$

Symmetry of two copies? $\rho_{\mathcal{H}} \otimes \rho_{\mathcal{H}}$

Permutation symmetry….measure the Young Diagram?

Nope [Childs, Harrow, Wocjan 06]

Heisenberg Group

Semidirect product group $\mathbb{Z}_p^2 \rtimes \mathbb{Z}_p$



Elements of the group:

$$\left(\begin{pmatrix} x \\ y \end{pmatrix}, b \right) \quad x, y, b \in \mathbb{Z}_p$$

Multiplication rule:

$$\left(\begin{pmatrix} x_1 \\ y_1 \end{pmatrix}, b_1 \right) \left(\begin{pmatrix} x_2 \\ y_2 \end{pmatrix}, b_2 \right) =$$

$$\left(\begin{pmatrix} x_1 \\ y_1 \end{pmatrix} + \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}^{b_1} \begin{pmatrix} x_2 \\ y_2 \end{pmatrix}, b_1 + b_2 \right)$$

From the play “Copenhagen” Margrethe Bohr to Heisenberg:

“Your talent is for ski-ing too fast for anyone to see where you are. For always being in more than one position at a time, like one of your particles.”

Multicopy Success!

Efficient algorithm for the Heisenberg HSP: $\mathbb{Z}_p^2 \rtimes \mathbb{Z}_p$

[D. Bacon, A.M. Childs, and W. van Dam FOCS 2005]

PGM/optimal measurement approach:

$$\rho_{\mathcal{H}} \qquad Pr(succ) = O\left(\frac{1}{p}\right)$$

$$\rho_{\mathcal{H}} \otimes \rho_{\mathcal{H}} \qquad Pr(succ) = O(1)$$

Efficient algorithm for the Heisenberg HSP:

1. PGM is optimal and succeed w.h.p.
2. How to efficiently implement the PGM?

But Why Does It Work?


Clebsch-Gordan Transform To The Rescue?

Problem:

$\rho_{\mathcal{H}} \otimes \rho_{\mathcal{H}}$ does not have a Clebsch-Gordan related symmetry




Graph Isomorphism



adjacency matrix A_1

$G_1 = (V_1, E_1)$



adjacency matrix A_2

$G_2 = (V_2, E_2)$

Problem: Is there a 1-to-1 mapping p from the vertices V_1 to the V_2 vertices such that the edges are respected, $p(E_1) = E_2$?

Mapping to the HSP:

$$f : \mathcal{S}_n \wr \mathcal{S}_2 \rightarrow \mathbb{Z}_{n(2n-1)}$$

$$f((\pi_1, \pi_2, \tau)) = \tau(\pi_1(A_1) \oplus \pi_2(A_2))$$

Notice now we are only worried if hidden subgroup has a nontrivial τ

Conjugate Subgroups

Two subgroups, $\mathcal{H}_1$ and $\mathcal{H}_2$, are conjugate if there exists an element of the group $g \in \mathcal{G}$ such that

$$\mathcal{H}_1 = \{ghg^{-1}, h \in \mathcal{H}_2\}$$

This is an equivalence relation among the subgroups of $\mathcal{G}$

$$\mathcal{H}_i \equiv_{\mathcal{G}} \mathcal{H}_j$$


ir: voy, vas, va, vamos,

Hidden Subgroup Conjugacy Problem

Setup: Fix a group $\mathcal{G}$(known) and a subgroup $\mathcal{H} \subset \mathcal{G}$(unknown). Given query access to a function $f : \mathcal{G} \to S$ that is constant and distinct on left (right) cosets of $\mathcal{H}$ in $\mathcal{G}$ We say that f hides $\mathcal{H}$

Problem: identify the conjugate subgroup class of $\mathcal{H}$.

Efficient: If runs in time polylog($|\mathcal{G}|$)



Symmetries of HSCP States

Hidden subgroup conjugate state

$$\rho_{[\mathcal{H}]} = \frac{1}{|\mathcal{G}|} \sum_{g \in \mathcal{G}} \rho_{g\mathcal{H}g^{-1}}$$

Hidden subgroup conjugate states

$$\rho_{[\mathcal{H}]}^{\otimes m} = \frac{1}{|\mathcal{G}|} \sum_{g \in \mathcal{G}} (\rho_{g\mathcal{H}g^{-1}})^{\otimes m}$$

In context of optimal measurement method (PGM)
[e.g. Ip 03, Bacon, Childs, and van Dam 05]
this translates into extra symmetry of the optimal measurement

$$\rho_{[\mathcal{H}]}^{\otimes m} = \bigoplus_{\mu_1,\ldots,\mu_m} \left(I_D \otimes \bigoplus_{\mu} \left(R_{[\mathcal{H}]} \otimes I_{d_\mu} \right) \right)$$

Symmetries of HSCP States

What is the extra symmetry of the multiple HSCP states?

Clebsch-Gordan transform for the regular representation of finite groups:

$$D_R(g)^{\otimes m} = \left[\bigoplus_{\mu}(I_{d_\mu} \otimes D_\mu(g))\right]^{\otimes m} \quad D_R(g)|g'\rangle = |g'g^{-1}\rangle$$

$$D_{\mu_1}(g) \otimes \cdots \otimes D_{\mu_m}(g) = \bigoplus_{\mu} \left(I_{n_\mu(\mu_1,\ldots,\mu_m)} \otimes D_\mu(g)\right)$$

Relationships

How do the hidden subgroup problem (HSP) and the hidden subgroup conjugacy problem (HSCP) differ?

1. If group is abelian, HSCP=HSP

All subgroups are only conjugate to themselves

$$g\mathcal{H}g^{-1} = \mathcal{H}$$

2. If subgroups are normal, HSCP=HSP

All subgroups are only conjugate to themselves


$$g\mathcal{H}g^{-1} = \mathcal{H}$$

These are two cases where we know quantum computers can efficiently solve the HSP. Coincidence?



General Circuit






Heisenberg HSCP and HSP

[Bacon, arXiv:quant-ph/0612107]

Heisenberg HSCP ⇒ Heisenberg HSP




Conjugate subgroups are all normal subgroups of a subgroup of Heisenberg group

Identifying which conjugate subgroup identifies this subgroup

Thus solving HSCP implies solving HSP (use normal subgroup algorithm)

Heisenberg HSCP, CG Way

[Bacon, arXiv:quant-ph/0612107]



i labels set of conjugate subgroups

Heisenberg HSCP, CG Way

[Bacon, arXiv:quant-ph/0612107]

$$\frac{1}{\sqrt{p}} \sum_{s \in \mathbb{Z}_p} \omega^{i 2^{-1} \frac{k_1 k_2}{k_1 + k_2} s^2} |s\rangle$$

i can be found by performing

$$|t^2\rangle \rightarrow \frac{1}{\sqrt{2}}(|t\rangle + |-t\rangle)$$

and performing a
QFT

Clebsch-Gordan transform over Heisenberg group is related to “symmetry” which allows for efficient quantum algorithm for the Heisenberg HSP.

Heisenberg HS State

Restricted subgroups:

$$\mathcal{H}_{d_1,d_2} = \left\{ \left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, 0 \right), \left(\begin{pmatrix} d_1 \\ d_2 \end{pmatrix}, 1 \right), \left(\begin{pmatrix} 2d_1 + d_2 \\ 2d_2 \end{pmatrix}, 2 \right), \ldots \right\}$$

Random coset states:

$$|l_1, l_2, d_1, d_2\rangle = \frac{1}{\sqrt{p}} \sum_{b \in \mathbb{Z}_p} \left| \left(\left(\begin{pmatrix} l_1 \\ l_2 \end{pmatrix} + M^{(b)} \begin{pmatrix} d_1 \\ d_2 \end{pmatrix} \right), b \right) \right\rangle$$

$$M^{(b)} = \sum_{i=0}^{b-1} \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}^i = \begin{bmatrix} b & \frac{b(b-1)}{2} \\ 0 & b \end{bmatrix}$$

Coset labels: $l_1, l_2 \in_U \mathbb{Z}_p$

Hidden Subgroup states:

$$\rho_{d_1,d_2} = \frac{1}{p^2} \sum_{l_1,l_2 \in \mathbb{Z}_p} |l_1, l_2, d_1, d_2\rangle\langle l_1, l_2, d_1, d_2|$$

HS

$$|l_1, l_2, d_1, d_2\rangle = \frac{1}{\sqrt{p}} \sum_{b \in \mathbb{Z}_p} \left| \left(\left(\begin{pmatrix} l_1 \\ l_2 \end{pmatrix} + M^{(b)} \begin{pmatrix} d_1 \\ d_2 \end{pmatrix} \right), b \right) \right\rangle$$

$$|l_1, l_2, d_1, d_2\rangle = \frac{1}{\sqrt{p}} \sum_{b \in \mathbb{Z}_p} \left| l_1 + bd_1 + \frac{b(1-b)}{2} d_2, l_2 + bd_2; b \right\rangle$$

Perform the QFT over $(\mathbb{Z}_p)^2$ to the first two registers:

$$|l_1, l_2, d_1, d_2\rangle = \frac{1}{\sqrt{p^3}} \sum_{b, x_1, y_1 \in \mathbb{Z}_p} \omega_p^{(l_1 + bd_1 + \frac{b(1-b)}{2} d_2) x_1 + (l_2 + bd_2) y_1} |x_1, y_1, b\rangle$$

Measure these two registers:

$$x_1, y_1 \in_U \mathbb{Z}_p$$

$$|x_1, y_1, d_1, d_2\rangle = \frac{1}{\sqrt{p}} \sum_{b \in \mathbb{Z}_p} \omega_p^{(bd_1 + \frac{b(1-b)}{2} d_2) x_1 + bd_2 y_1} |b\rangle$$

HS

Two copies: $x_1, y_1, x_2, y_2 \in_U \mathbb{Z}_p$

$$\frac{1}{p} \sum_{b_1, b_2 \in \mathbb{Z}_p} \omega_p^{(b_1 x_1 + b_2 x_2) d_1 + \left(\frac{b_1(1-b_1)}{2} x_1 + b_1 y_1 + \frac{b_2(1-b_2)}{2} x_2 + b_2 y_2 \right) d_2} |b_1, b_2\rangle$$

$$v = b_1 x_1 + b_2 x_2$$

$$w = \frac{b_1(1-b_1)}{2} x_1 + b_1 y_1 + \frac{b_2(1-b_2)}{2} x_2 + b_2 y_2$$

$$\frac{1}{p} \sum_{b_1, b_2 \in \mathbb{Z}_p} \omega_p^{v d_1 + w d_2} |b_1, b_2\rangle$$

Compute v and w:

$$\frac{1}{p} \sum_{b_1, b_2 \in \mathbb{Z}_p} \omega_p^{v d_1 + w d_2} |b_1, b_2\rangle |v, w\rangle$$

Uncompute

If we could erase the b1,b2 values:

$$\frac{1}{p}\sum_{b_1,b_2\in\mathbb{Z}_p}\omega_p^{vd_1+wd_2}|b_1,b_2\rangle|v,w\rangle \rightarrow \frac{1}{p}|0,0\rangle\sum_{v,w}\omega_p^{vd_1+wd_2}|v,w\rangle$$

Then a QFT over $(\mathbb{Z}_p)^2$ would find d1 and d2.

Remember v and w are a function of b1 and b2.

Solving Equations

Compute v and w:

$$\frac{1}{p}\sum_{b_1,b_2\in\mathbb{Z}_p}\omega_p^{vd_1+wd_2}|b_1,b_2\rangle|v,w\rangle$$

$$v = b_1x_1 + b_2x_2$$

$$w = \frac{b_1(1-b_1)}{2}x_1 + b_1y_1 + \frac{b_2(1-b_2)}{2}x_2 + b_2y_2$$

For fixed x1,x2,y1,y2 let $S_{v,w}$ be the set of b1,b2

solutions to these equations

$$\frac{1}{p}\sum_{v,w\in\mathbb{Z}_p}\omega_p^{vd_1+wd_2}\left(\sum_{b_1,b_2\in S_{v,w}}|b_1,b_2\rangle\right)|v,w\rangle$$

We want to implement $\frac{1}{\sqrt{|S_{v,w}|}}\sum_{b_1,b_2\in S_{v,w}}|b_1,b_2\rangle|v,w\rangle \rightarrow |0,0\rangle|v,w\rangle$

Solving Equations

$$\frac{1}{\sqrt{|S_{v,w}|}} \sum_{b_1,b_2 \in S_{v,w}} |b_1, b_2\rangle |v, w\rangle \rightarrow |0,0\rangle |v,w\rangle$$

Easier to think about the inverse (do it unitarily):

$$|0,0\rangle |v,w\rangle \rightarrow \frac{1}{\sqrt{|S_{v,w}|}} \sum_{b_1,b_2 \in S_{v,w}} |b_1, b_2\rangle |v, w\rangle$$

i.e. given v and w, transform to superposition over solutions to the equations

$$v = b_1 x_1 + b_2 x_2$$

$$w = \frac{b_1(1-b_1)}{2} x_1 + b_1 y_1 + \frac{b_2(1-b_2)}{2} x_2 + b_2 y_2$$

Solving Equations

$$v = b_1 x_1 + b_2 x_2$$

$$w = \frac{b_1(1-b_1)}{2} x_1 + b_1 y_1 + \frac{b_2(1-b_2)}{2} x_2 + b_2 y_2$$

Assuming y1,y2,y1+y2 do not equal 0 (prob O(1/p)),
then

$\frac{p(p-1)}{2}$ solutions have $|S_{v,w}| = 0$

p solutions have $|S_{v,w}| = 1$

$\frac{p(p-1)}{2}$ solutions have $|S_{v,w}| = 2$

Solving Equations

$$v = b_1 x_1 + b_2 x_2$$

$$w = \frac{b_1(1-b_1)}{2} x_1 + b_1 y_1 + \frac{b_2(1-b_2)}{2} x_2 + b_2 y_2$$

Focus on when there are two solutions.

We can use standard finite arithmetic to find these solutions. Use to create labeled superposition of solutions:

$$\frac{1}{\sqrt{2}} \sum_{j=0,1} |j, (b_1)_j, (b_2)_j\rangle$$

Hadamard first qubit, measure. If 0, you have

$$\frac{1}{\sqrt{2}} \sum_{j=0,1} |(b_1)_j, (b_2)_j\rangle$$

Heisenberg HSP Solved

This allows us to perform:

$$\frac{1}{p}\sum_{b_1,b_2\in\mathbb{Z}_p}\omega_p^{vd_1+wd_2}|b_1,b_2\rangle|v,w\rangle \rightarrow \frac{1}{p}|0,0\rangle\sqrt{|S_{v,w}|}\sum_{v,w}\omega_p^{vd_1+wd_2}|v,w\rangle$$

Which has a O(1/2) fidelity squared with the state we desire.

QFTing the second register will let us find d1 and d2

Efficient algorithm for the Heisenberg HSP!

Heisenberg HSP Lesson

Clebsch-Gordan transform can be used to help solve HSPs

Other groups where this works?

$(\mathbb{Z}_4 \rtimes \mathbb{Z}_2)^n$ [Ivanyos, Magniez, Sen 01]

Some Extraspecial groups [Ivanyos, Sanselme, Santha 07]

But for important groups like dihedral and symmetric?

Hidden Subgroup Divide

Successes
- abelian groups
- almost abelian groups
- near Hamiltonian groups
- normal subgroups
- small commutator subgroup
- solvable groups of bounded exponent and of bounded derived series
- extraspecial groups
- groups of nilpotency class at most 2

Specific groups

$\mathbb{Z}_2^n \rtimes \mathbb{Z}_2$

$\mathbb{Z}_{p^k}^n \rtimes \mathbb{Z}_2$

some $\mathbb{Z}_p^k \rtimes \mathbb{Z}_p$

including

the Heisenberg group

$\mathbb{Z}_N \rtimes \mathbb{Z}_p, N/p = \text{polylog} N$

Failures
- symmetric group **(graph ismorphism!)**
- dihedral group **(shortest vector!)**
- Gn, fixed G. Even for G=S3!

Pessimism
- in standard approach, multiregister measurements required
- optimal measurement for dihedral HSP leads naturally to subset sum problem
- Clebsch-Gordan sieves cannot solve symmetric group HSP

Subexponential algorithms
- Dihedral
- Gn, fixed G

Fin

Clebsch-Gordan transform reveals essential symmetry of efficiently solvable nonabelian hidden subgroup problems….Can they lead us all the way, the $109 question



Quick advertisement: Scirate.com

