

Measurement of $H \rightarrow b\bar{b}$ in Associated Production with the CMS Detector

by

Daniel Robert Abercrombie

B.S., Pennsylvania State University (2014)

Submitted to the Department of Physics
in partial fulfillment of the requirements for the degree of

Doctor of Philosophy

at the

MASSACHUSETTS INSTITUTE OF TECHNOLOGY

June 2021

© Massachusetts Institute of Technology 2021. All rights reserved.

Author

Department of Physics
March 27, 2021

Certified by

Christoph M. E. Paus
Professor of Physics
Thesis Supervisor

Accepted by

Deepto Chakrabarty
Associate Department Head, Physics

Measurement of $H \rightarrow b\bar{b}$ in Associated Production with the CMS Detector

by

Daniel Robert Abercrombie

Submitted to the Department of Physics
on March 27, 2021, in partial fulfillment of the
requirements for the degree of
Doctor of Philosophy

Abstract

The differential cross section of $VH \rightarrow b\bar{b}$ is measured with the CMS Detector. The Simplified Template Cross Section framework is used. The inclusive strength of the measured signal relative to the Standard Model is $0.568^{+0.154}_{-0.147}(\text{stat})^{+0.134}_{-0.133}(\text{sys})$, which agrees with the Standard Model within 2.1 standard deviations. The measured spectrum of the recoiling vector boson transverse momentum has a p -value of 9.3%, assuming Standard Model predictions at the measured signal strength.

Thesis Supervisor: Christoph M. E. Paus
Title: Professor of Physics

Acknowledgments

Thanks.

Contents

| | | |
|----------|---|-----------|
| 1 | Introduction | 17 |
| 1.1 | Measurement of the Higgs Cross Section | 18 |
| 1.2 | Motivation for the Measurement | 19 |
| 1.3 | Using the CMS Detector at the LHC | 20 |
| 2 | Theory | 21 |
| 2.1 | Electroweak Symmetry Breaking | 23 |
| 2.2 | Associated Production | 26 |
| 2.2.1 | Coupling Between Vector Bosons and Fermions | 27 |
| 2.2.2 | Decay Channels of Vector Bosons | 32 |
| 2.3 | Decay Channels of the Higgs | 33 |
| 3 | The CMS Detector | 37 |
| 3.1 | The Large Hadron Collider | 37 |
| 3.2 | Detector Requirements | 39 |
| 3.3 | Detector Design | 40 |
| 3.3.1 | Solenoid Magnet | 42 |
| 3.3.2 | Silicon Tracker | 42 |
| 3.3.3 | Electromagnetic Calorimeter | 43 |
| 3.3.4 | Hadronic Calorimeter | 44 |
| 3.3.5 | Muon Chambers | 45 |
| 3.4 | Event Reconstruction | 46 |
| 3.4.1 | Charged Particle Tracks | 46 |

| | | |
|----------|--|-----------|
| 3.4.2 | Calorimeters | 47 |
| 3.4.3 | Linking and Particle Identification | 48 |
| 3.5 | Trigger | 49 |
| 3.6 | Simulation | 51 |
| 3.6.1 | Short-Scale Simulation | 52 |
| 3.6.2 | Parton Showers | 53 |
| 3.6.3 | Detector Simulation | 53 |
| 3.7 | Accessing Data | 54 |
| 4 | Event Selection | 55 |
| 4.1 | Object Definitions | 55 |
| 4.1.1 | Variable Definitions | 56 |
| 4.1.2 | Isolated Muons | 57 |
| 4.1.3 | Isolated Electrons | 58 |
| 4.1.4 | Jets | 59 |
| 4.1.5 | Identification of b Jets and Energy Regression | 60 |
| 4.1.6 | Fat Jets | 62 |
| 4.1.7 | Missing Transverse Momentum | 63 |
| 4.1.8 | Soft Hadronic Activity | 63 |
| 4.1.9 | Kinematic Fit | 64 |
| 4.2 | Simplified Template Cross Section Bins | 65 |
| 4.3 | Treatment of Background Yields | 66 |
| 4.3.1 | Identification of Background Processes | 66 |
| 4.3.2 | Goals for Control Regions | 68 |
| 4.4 | Resolved Analysis Selection | 68 |
| 4.4.1 | 0 Leptons | 71 |
| 4.4.2 | 1 Lepton | 74 |
| 4.4.3 | 2 Leptons | 75 |
| 4.5 | Boosted Analysis Selection | 76 |
| 4.5.1 | 0 Leptons | 76 |

| | | |
|----------|---|------------|
| 4.5.2 | 1 Lepton | 77 |
| 4.5.3 | 2 Leptons | 77 |
| 4.6 | Overlap in Resolved and Boosted Selections | 78 |
| 5 | Corrections and Uncertainties | 79 |
| 5.1 | Efficiency Scale Factors | 80 |
| 5.1.1 | Muons | 80 |
| 5.1.2 | Electrons | 81 |
| 5.1.3 | MET Trigger Scale Factors | 82 |
| 5.2 | Energy Corrections of b -Jets | 82 |
| 5.3 | Theoretical Corrections | 87 |
| 5.3.1 | LO to NLO Reweighting | 87 |
| 5.3.2 | Electroweak and QCD Corrections to Background | 90 |
| 5.3.3 | Corrections to Signal Process | 90 |
| 5.4 | Application of Uncertainties | 92 |
| 5.4.1 | Normalization Uncertainties | 92 |
| 5.4.2 | Shape Uncertainties | 93 |
| 5.4.3 | Migration Uncertainties | 94 |
| 6 | Analysis Results | 95 |
| 6.1 | Run 2 Data Collection | 95 |
| 6.2 | Control Region Plots | 95 |
| 6.3 | Multivariate Discriminator | 102 |
| 6.3.1 | Resolved DNN | 102 |
| 6.3.2 | Boosted BDT | 102 |
| 6.4 | Combination Fit | 104 |
| 6.4.1 | VZ Cross Check Analysis | 105 |
| 6.4.2 | VH Combination Fit Results | 108 |
| 7 | Conclusions | 119 |

| | | |
|----------|-----------------------------------|------------|
| A | Detector Projects | 121 |
| A.1 | Dynamo Consistency | 121 |
| A.1.1 | Installation | 123 |
| A.1.2 | Inventory Listing | 123 |
| A.1.3 | Remote Listing | 125 |
| A.1.4 | Executables | 126 |
| A.1.5 | Configuration | 130 |
| A.1.6 | Comparison Script | 133 |
| A.2 | Workflow Web Tools | 134 |
| A.2.1 | Error Information | 135 |
| A.2.2 | Possible Actions | 135 |
| A.2.3 | Automatic Actions | 136 |
| B | Simulation Details | 137 |
| C | Data Card | 141 |
| D | Nuisance Parameter Impacts | 173 |

List of Figures

| | | |
|-----|--|----|
| 2-1 | Feynman diagram of associated production | 27 |
| 2-2 | Feynman diagram of generating W^\pm | 28 |
| 2-3 | Feynman diagram of generating Z | 30 |
| 2-4 | Parton Distribution Function for protons | 32 |
| 2-5 | Feynman diagrams of other production mechanisms | 33 |
| 2-6 | Tau decay | 34 |
| 2-7 | Vector Boson decays in the analysis | 34 |
| 2-8 | Full Feynman diagram for the two lepton process | 36 |
| 3-1 | CMS detector slice | 41 |
| 4-1 | Higgs di-jet mass fit with kinematic fit | 65 |
| 4-2 | Feynman diagram for $DY + \text{jets}$ background | 67 |
| 4-3 | Feynman diagram for $t\bar{t}$ background | 67 |
| 4-4 | Efficiency for MET trigger in data | 72 |
| 4-5 | MET ϕ distribution before and after shutting off HCAL modules . . . | 72 |
| 5-1 | Tag and probe fits | 81 |
| 5-2 | MET trigger scale factors | 83 |
| 5-3 | Distribution of α used for b jet energy smearing | 85 |
| 5-4 | Response to evaluate jet smearing | 86 |
| 5-5 | Resolution fits for jet smearing | 87 |
| 5-6 | LO to NLO reweighting shape comparisons | 89 |
| 5-7 | Electroweak corrections to background | 90 |

| | | |
|------|---|-----|
| 5-8 | VH in NNLO QCD | 91 |
| 6-1 | Control region plots for two leptons in 2016 | 96 |
| 6-2 | Control region plots for one lepton in 2016 | 97 |
| 6-3 | Control region plots for zero lepton in 2017 | 98 |
| 6-4 | Control region plots for two electrons in 2017 | 99 |
| 6-5 | Control region plots for one lepton in 2018 | 100 |
| 6-6 | Control region plots for two muons in 2018 | 101 |
| 6-7 | Goodness of Fit for VZ | 106 |
| 6-8 | Example impacts for WH $250 < p_T(V) < 400$ GeV | 107 |
| 6-9 | $Z \rightarrow \mu\mu$ VZ distributions for low p_T in 2017 | 109 |
| 6-10 | $W \rightarrow e\nu$ VZ distributions for the boosted selection in 2018 | 110 |
| 6-11 | Inclusive likelihood scan of VZ | 111 |
| 6-12 | Measured STXS values of VZ | 112 |
| 6-13 | Goodness of Fit for VH | 112 |
| 6-14 | $Z \rightarrow \nu\nu$ VH distributions for high p_T in 2016 | 114 |
| 6-15 | $W \rightarrow \mu\nu$ VH distributions for medium p_T in 2017 | 115 |
| 6-16 | $Z \rightarrow ee$ VH distributions for medium p_T in 2018 | 116 |
| 6-17 | Inclusive likelihood scan of VH | 117 |
| 6-18 | Measured values of WH and ZH | 117 |
| 6-19 | Measured STXS values of VH | 118 |
| 6-20 | Measured values of κ_V and κ_b | 118 |
| A-1 | Comparison algorithm | 125 |
| A-2 | Listing algorithm with retries | 126 |
| A-3 | Example error information | 136 |
| D-1 | Impacts for WH $p_T(V) > 400$ GeV | 174 |
| D-2 | Impacts for WH $250 < p_T(V) < 400$ GeV | 175 |
| D-3 | Impacts for WH $150 < p_T(V) < 250$ GeV | 176 |
| D-4 | Impacts for ZH $p_T(V) > 400$ GeV | 177 |

| | | |
|-----|--|-----|
| D-5 | Impacts for ZH $250 < p_T(V) < 400$ GeV | 178 |
| D-6 | Impacts for ZH $150 < p_T(V) < 250$ GeV with jets | 179 |
| D-7 | Impacts for ZH $150 < p_T(V) < 250$ GeV without jets | 180 |
| D-8 | Impacts for ZH $75 < p_T(V) < 150$ GeV | 181 |

List of Tables

| | | |
|-----|---|-----|
| 2.1 | Fermion charges | 22 |
| 4.1 | DeepCSV working points | 61 |
| 4.2 | Summary of resolved selection cuts | 70 |
| 4.3 | Triggers for the 0 lepton selection | 71 |
| 4.4 | Triggers for the 1 lepton selections | 74 |
| 4.5 | Triggers for the 2 lepton selections | 75 |
| 5.1 | b -jet energy correction parameters | 87 |
| 6.1 | Resolved DNN inputs | 103 |
| A.1 | dynamo-consistency site statuses | 128 |
| B.1 | 2016 MC samples | 138 |
| B.2 | 2017 MC samples | 139 |
| B.3 | 2018 MC samples | 140 |
| C.1 | 2016 0-lepton signal selection yields | 142 |
| C.2 | 2016 0-lepton control region yields | 143 |
| C.3 | 2016 1-lepton (e) signal selection yields | 144 |
| C.4 | 2016 1-lepton (e) control region yields | 145 |
| C.5 | 2016 1-lepton (μ) signal selection yields | 146 |
| C.6 | 2016 1-lepton (μ) control region yields | 147 |
| C.7 | 2016 2-lepton (e) signal selection yields | 148 |
| C.8 | 2016 2-lepton (e) control region yields | 149 |

| | | |
|------|---|-----|
| C.9 | 2016 2-lepton (μ) signal selection yields | 150 |
| C.10 | 2016 2-lepton (μ) control region yields | 151 |
| C.11 | 2017 0-lepton signal selection yields | 152 |
| C.12 | 2017 0-lepton control region yields | 153 |
| C.13 | 2017 1-lepton (e) signal selection yields | 154 |
| C.14 | 2017 1-lepton (e) control region yields | 155 |
| C.15 | 2017 1-lepton (μ) signal selection yields | 156 |
| C.16 | 2017 1-lepton (μ) control region yields | 157 |
| C.17 | 2017 2-lepton (e) signal selection yields | 158 |
| C.18 | 2017 2-lepton (e) control region yields | 159 |
| C.19 | 2017 2-lepton (μ) signal selection yields | 160 |
| C.20 | 2017 2-lepton (μ) control region yields | 161 |
| C.21 | 2018 0-lepton signal selection yields | 162 |
| C.22 | 2018 0-lepton control region yields | 163 |
| C.23 | 2018 1-lepton (e) signal selection yields | 164 |
| C.24 | 2018 1-lepton (e) control region yields | 165 |
| C.25 | 2018 1-lepton (μ) signal selection yields | 166 |
| C.26 | 2018 1-lepton (μ) control region yields | 167 |
| C.27 | 2018 2-lepton (e) signal selection yields | 168 |
| C.28 | 2018 2-lepton (e) control region yields | 169 |
| C.29 | 2018 2-lepton (μ) signal selection yields | 170 |
| C.30 | 2018 2-lepton (μ) control region yields | 171 |

Chapter 1

Introduction

One of the most curious features of physics at small scales, which will likely frustrate students for the rest of time, is that certain events are not deterministic and only have a probability of happening. There is no guarantee that the initial state of an electron and a positron approaching each other will annihilate and produce the final state of a muon and an anti-muon, even if their energies in the center-of-mass frame are adequate for muon production. However, trying long enough with the same initial conditions will eventually produce a muon and anti-muon pair. Furthermore, the observation of resonances, where this is more likely to happen when the electron and positron approach each other at particular energies, does not mean that a Z boson was present in a given interaction. It just means that the weak component of the electroweak force significantly increases the probability of the muonic final state, when the total energy of the system is close to the mass of the Z boson. The sum of probabilities from different possible field interactions leading from a specific initial condition to a particular final state is the only thing that can be measured. This measurement is only done accurately when observing many events with the same initial conditions.

This point is difficult to convey concisely, so many laypeople, as well as some practicing physicists, are confused by the terminology adopted by the field. But this distinction is relevant to the topic of this work. This document presents a measurement of a cross section. Cross section is the name given to the probability of an

interaction occurring. Reported cross sections can be split up to describe different contributions to final states, and they can be collated into what are called “production cross sections” which describe the probabilities of particular intermediate states “occurring” (even though intermediate states never exist in reality). In the example above, the Z boson would be a possible intermediate state.

The main point of the example above is that physicists can learn about the Z boson by observing only electrons and muons. They never see the Z boson itself. This can be generalized to any interesting particle that interacts with other particles. An increased probability of certain initial states resulting in certain final states can teach the observer about the role of the interesting particle without ever directly seeing it.

1.1 Measurement of the Higgs Cross Section

The purpose of the following document is to present the methods and results of measuring the strength of the coupling between the Higgs boson and bottom quarks. In this context, the Higgs boson makes up one of the previously mentioned intermediate states that cannot be shown as present in a given event. The coupling strength is directly related to the probability contribution the Higgs field has on processes involving bottom quarks. However, the cross section measurement also relies on a number of other physics processes.

To measure this coupling, the Higgs boson must first be “produced” before measuring its coupling strength to bottom quarks. This analysis takes advantage of a process known as associated production, where a vector boson, one of the intermediate particles of the weak nuclear force, couples to and radiates a Higgs boson. The vector boson is in turn produced by the collision of high energy protons.

The measurement is not complete once the intermediate state is generated. The Higgs boson can decay into a number of different particles, with bottom quarks being only one type. The bottom quarks must be identified. The vector bosons have multiple decay modes as well. In this analysis, we only use the leptonic decays

because these give us the cleanest signature, where enough of the contribution to the final state probability is from associated production for it to be measured.

There are also other physics processes that create the same final states, as well as processes that create final states that look similar enough to be practically indistinguishable. These processes must also be well understood before a Higgs boson cross section measurement can be undertaken.

1.2 Motivation for the Measurement

The measurement of a cross section of a known particle is “normal science,” and that is the space in which this analysis operates [1]. Much of the community of physics researchers have been operating under the paradigm of the Standard Model for the better part of a century. The Standard Model has known gaps, such as missing explanations for neutrino mass, as well as the origin of Dark Matter and Dark Energy. However, none of these research fields have yielded any results that will trigger a paradigm shift. In fact, the most exciting discoveries of new particles, such as the weak W [2] and Z [3] bosons and the Higgs boson [4, 5] only confirmed predictions by the Standard Model.

In the meanwhile, precision measurements are performed on processes that we expect to already understand very well. Repeating measurements while the state of the art is improving is interesting, no matter the outcome. Over time, the uncertainty in the measurement outcome shrinks, leading to more precise knowledge of parameters of the Standard Model. If the precise parameters cause excessive tension in that they cannot exist assuming the Standard Model is true, the discrepancies would need to be explained by a different or amended model.

The Higgs decaying to $b\bar{b}$, or a bottom quark and bottom anti-quark, was observed in 2018 [6, 7]. The measurement outlined in this document goes further in that it measures the contribution of $H \rightarrow b\bar{b}$ in associated production to final states of different energy. **Phil** collision energy or production energy? be clear This is called a differential cross section, and places greater constraints on the parameters of the

Standard Model. These constraints lead to more precise measurements of the parameters, and have the potential to discover discrepancies that have hitherto been missed.

Phil Maybe you will go into more detail, but I feel like you can add a paragraph or two to motivate why differential xs are so important and what we learn from them. Recall we did not know exactly how the Higgs boson behaved when we discovered it, there was no reasonable mass prediction at all.

1.3 Using the CMS Detector at the LHC

This measurement is only possible due to massive efforts by the scientific community. The Higgs boson is not typically generated in conditions on Earth. The Large Hadron Collider (LHC) at the European Organization for Nuclear Research (CERN) was constructed due to the efforts of thousands of scientists and engineers, as well as the funding from countries distributed all around the globe. The LHC performs the proton-proton collisions needed.

To observe the final states of collisions at the LHC, multiple detectors have also been constructed, due to the efforts of hundreds or thousands of individuals. This analysis is done using data from the Compact Muon Solenoid (CMS). Other experiments are ATLAS, ALICE, LHCb, TOTEM, LHCf, MoDEL, and FASER. The CMS detector is a general purpose detector, which was used in the discovery of the Higgs boson. Its detection capabilities make it instrumental in a number of state of the art measurements of the Standard Model as well as in searches for physics beyond the Standard Model.

Phil This section feels incomplete, I think you need to motivate the measurement more. Also you should motivate the LHC more. Why is there such a device?

Chapter 2

Theory

The Standard Model describes the interactions of all observable matter. There are many textbooks that cover the Standard Model, as there are many scientists and students who study it. For an in depth presentation of the Standard Model beyond what is presented in this chapter, please refer to Reference [8].

Matter is made up of 12 kinds of fermions. The forces between the fermions are mediated by the gauge fields created by the Standard Model's $SU(3) \times SU(2)_L \times U(1)_Y$ symmetry. Fermions with the appropriate charge are affected by the gauge fields. Fermions are classified as quarks or leptons. There are six types of quarks, separated into three generations of two quarks each. Each quark has a color charge associated with the $SU(3)$ symmetry. The interactions arising from this is called QCD. Each pair of quarks in a family also have an approximate $SU(2)$ symmetry, which allows interactions via the weak force. Quarks also have a hypercharge, which is a relation of electromagnetic charge and weak isospin, meaning the gauge field from the $U(1)$ symmetry affects them as well. Three charged leptons and three neutral leptons, called neutrinos, comprise the other six fermions. The leptons do not carry a color charge, so they are not affected by the $SU(3)$ symmetry, but they do carry weak isospin and hypercharge. Table 2.1 displays the values of these charges for all fermions. Fermions also have anti-particles, which carry the opposite charges of their counterparts.

In the Standard Model, the gauge fields from the $SU(2)_L \times U(1)_Y$ symmetries are

Table 2.1: All of the fermions are listed below, along with their charges and weak isospin values. The three generations are listed from least to most massive, meaning only the first generation of quarks and charged leptons is stable. The masses and decays of neutrinos is beyond the scope of the Standard Model and this analysis.

| | 1st gen. | 2nd gen. | 3rd gen. | Color | Q | $I_W^{(3)}$ | Y |
|------------------|----------|-----------|------------|-------|----------------|----------------|----------------|
| down-type quarks | d | s | b | yes | $-\frac{1}{3}$ | $-\frac{1}{2}$ | $-\frac{1}{3}$ |
| up-type quarks | u | c | t | yes | $+\frac{2}{3}$ | $+\frac{1}{2}$ | $+\frac{5}{3}$ |
| charged leptons | e | μ | τ | no | -1 | $-\frac{1}{2}$ | -1 |
| neutral leptons | ν_e | ν_μ | ν_τ | no | 0 | $+\frac{1}{2}$ | -1 |

mixed into what is known as the electroweak force. The electroweak force is mediated by the neutral photon and Z boson, and the charged W boson. This mixing happens due to the Higgs boson, a scalar which grants all of the charged fermions and the Z and W bosons mass through interactions.

PhilI think you should try to tie this paragraph with the above flow. Firstly, elude to the measuerement as being the focus of your thesis. Secondly, you should connect it to the SM In associated production, a Higgs boson is created through radiation from the Z and W bosons, which are also known as vector bosons. The measurement of $H \rightarrow b\bar{b}$ in associated production is therefore a measurement of interactions between the Higgs boson and vector bosons and the coupling between the Higgs boson and bottom, or b , quarks. The generation and observable decay of the vector bosons are also important to make this measurement. The parameters for those interactions are measured more accurately by other analyses not involving an observation of the Higgs [9]. Since the coupling of the Higgs boson with the vector bosons and with fermions also gives rise to the masses of each in the Standard Model through what is called electroweak symmetry breaking or the Higgs mechanism, most discussions of Higgs couplings include an explanation of the Higgs mechanism.

PhilI am confused by the paragraph below. Personally, I find expalanations about what I am about to read confusing and a waste of time, I think this paragraph doesn't fit with the flow of this intro to theory. The treatment of these topics in this chapter are arranged as follows. First, I will give a brief explanation of Higgs

field's non-zero vacuum energy, a trait that makes the electroweak symmetry breaking possible. After that, the coupling of the Higgs boson to the W and Z vector bosons will be described. These interactions arise from a mixing of the weak force with the electromagnetic force. What results is collectively known as the electroweak force. The coupling of the electroweak force to fermions is then discussed to understand both the generation of the vector boson intermediate states and the resulting final state that can be observed. Finally, the decay of the Higgs boson itself into bottom quarks is explained. This is allowed because the Higgs boson couples directly to massive fermions and gives them mass through the Higgs mechanism.

The Standard model is described by its Lagrangian, which is the difference between a system's kinetic and potential energies. Equations of motion are extracted from a Lagrangian \mathcal{L} for a particle field ϕ_i using the Euler-Lagrange equations.

$$\delta_\mu \left(\frac{\delta \mathcal{L}}{\delta(\delta_\mu \phi_i)} \right) - \frac{\delta \mathcal{L}}{\delta \phi_i} = 0 \quad (2.1)$$

Phil I really like you brought in the E-L equation. However, I think you need to tie the text into it more. It looks like you are going to go into detail into EWSB later. I might restructure this section to either put less details here, and more later or more now, and less later. Perhaps being a bit more abstract to start before you mention all the gauge fields will do that Gauge bosons are produced by symmetries of the Lagrangian. The Standard Model has $SU(3) \times SU(2)_L \times U(1)_Y$ symmetry. The $SU(3)$ symmetry produces gluons which mediate the strong force between quarks [10]. The $SU(2)_L \times U(1)_Y$ symmetry produces two gauge fields. The first interacts with left handed fermions, and the second interacts with all fermions through their hypercharge, Y . These forces are ultimately mixed into what is known as the electroweak force due to electroweak symmetry breaking [11]. Electroweak symmetry breaking is the required solution of the problem that vector gauge bosons of the electroweak force had mass. It was not possible to grant these bosons mass while maintaining the symmetries of the Standard Model without the Higgs boson [12–14]. The granting

of mass happens for two reasons: the Higgs field has a non-zero vacuum expectation value, and the Higgs field couples directly to vector boson and massive fermion fields.

2.1 Electroweak Symmetry Breaking

First, consider the $SU(2) \times U(1)$ symmetry where the Higgs interacts with the electroweak bosons. The L and Y of the $SU(2)_L \times U(1)_Y$ symmetry can be forgotten for the moment, since they describe fermion interactions with the electroweak force. To preserve $SU(2)$ symmetry, the Higgs boson is described as two complex scalar fields in a weak isospin doublet with a quartic potential. The $SU(2)$ symmetry means rotations between the doublet states must be equivalent in the Lagrangian. The Lagrangian for a free Higgs is then

$$\mathcal{L} = (\delta_\mu \phi)^\dagger (\delta^\mu \phi) - (\mu^2 (\phi^\dagger \phi) + \lambda (\phi^\dagger \phi)^2) \quad (2.2)$$

Through the virial theorem [15], the potential has a minimum value when

$$\phi^\dagger \phi = \frac{-\mu^2}{2\lambda} = \frac{v^2}{2} \quad (2.3)$$

This potential of the Higgs field breaks the $SU(2) \times U(1)$ symmetry of the Standard Model Lagrangian. Through this non-zero vacuum expectation value, the Higgs then has a constant influence in other parts of the Standard Model Lagrangian. In this way, it gives mass to electroweak vector bosons, to itself, and to massive fermions.

The first two sets of masses manifest when we force the $SU(2) \times U(1)$ symmetry back onto the Lagrangian in Equation (2.2). The derivatives must be replaced.

$$\delta_\mu \rightarrow D_\mu = \delta_\mu + i \frac{g_W}{2} \boldsymbol{\sigma} \cdot \mathbf{W}_\mu + i g' \frac{Y}{2} B_\mu \quad (2.4)$$

To simplify the expansion of Equation (2.2), a particular gauge, or particular doublet state, is chosen. ϕ is written to satisfy the vacuum expectation value of Equation (2.3) in the gauge that will give us the massless neutral boson known as a photon. **Phil**It

might be good to motivate this choice either through a reference or actually not setting the other double to zero now, and showing that it needs to be zero to ensure the photon is massless

$$\phi(x) = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v + h(x) \end{pmatrix} \quad (2.5)$$

This leads to the following expansion for the kinetic term of the Lagrangian.

$$\begin{aligned} (D_\mu \phi)^\dagger (D^\mu \phi) &= \frac{1}{2} (\delta_\mu h) (\delta^\mu h) + \frac{1}{8} g_W^2 (W_\mu^{(1)} + i W_\mu^{(2)}) (W^{(1)\mu} - i W^{(2)\mu}) (v + h)^2 \\ &\quad + \frac{1}{8} (g_W W_\mu^{(3)} - g' B_\mu) (g_W W^{(3)\mu} - g' B^\mu) (v + h)^2 \end{aligned} \quad (2.6)$$

Terms that are quadratic in terms of the gauge boson fields reveal the mass of the fields. Taking $h(x) \rightarrow 0$, the terms for $W^{(1)}$ and $W^{(2)}$ are the just **Phil** are just

$$\frac{1}{4} g_W^2 v^2 W_\mu^{(1)} W^{(1)\mu} \quad \text{and} \quad \frac{1}{4} g_W^2 v^2 W_\mu^{(2)} W^{(2)\mu},$$

giving the mass.

$$m_W = \frac{1}{2} g_W v \quad (2.7)$$

The quadratic terms for $W^{(3)}$ and B mix to give a non-diagonal mass matrix \mathbf{M} .

$$\frac{v^2}{8} \begin{pmatrix} W_\mu^{(3)} & B_\mu \end{pmatrix} \mathbf{M} \begin{pmatrix} W^{(3)\mu} \\ B^\mu \end{pmatrix} = \frac{v^2}{8} \begin{pmatrix} W_\mu^{(3)} & B_\mu \end{pmatrix} \begin{pmatrix} g_W^2 & -g_W g' \\ -g_W g' & g'^2 \end{pmatrix} \begin{pmatrix} W^{(3)\mu} \\ B^\mu \end{pmatrix} \quad (2.8)$$

The non-diagonal matrix allow $W^{(3)}$ and B to mix. Physical states must be represented by a diagonal Hamiltonian. Diagonalizing the term above gives masses of the physical states.

$$\frac{1}{8} v^2 \begin{pmatrix} A_\mu & Z_\mu \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & g_W^2 + g'^2 \end{pmatrix} \begin{pmatrix} A^\mu \\ Z^\mu \end{pmatrix} = \frac{1}{2} \begin{pmatrix} A_\mu & Z_\mu \end{pmatrix} \begin{pmatrix} m_A^2 & 0 \\ 0 & m_Z^2 \end{pmatrix} \begin{pmatrix} A^\mu \\ Z^\mu \end{pmatrix} \quad (2.9)$$

This gives us the masses of the neutral gauge bosons.

$$m_A = 0 \quad \text{and} \quad m_Z = \frac{1}{2}v\sqrt{g_W^2 + g'^2} \quad (2.10)$$

From the simple act of requiring $SU(2) \times U(1)$ symmetry on the Lagrangian of a scalar doublet with non-zero vacuum expectation value, the masses of all the electroweak gauge bosons have been produced. A similar procedure will produce the masses of fermions due to their coupling to the electroweak force. Phil While I agree it is similar, I think this should be in your thesis, you are doing $H \Rightarrow b \bar{b}$, the mechanisms are different enough that people thought there might be 2 Higgses

2.2 Associated Production

The next thing to consider is the couplings also produced by this process. The couplings will allow us to determine more precisely the parameters above by measuring cross sections.

The physical states of W^+ and W^- bosons can be written as the raising and lowering operators for isospin.

$$W^\pm = \frac{1}{\sqrt{2}} (W^{(1)} \mp iW^{(2)}) \quad (2.11)$$

The second term of Equation (2.6) can be further expanded.

$$\frac{1}{4}g_W^2 W_\mu^- W^{+\mu} (v + h)^2 = \frac{1}{4}g_W^2 v^2 W_\mu^- W^{+\mu} + \frac{1}{2}g_W^2 v W_\mu^- W^{+\mu} h + \frac{1}{4}g_W^2 W_\mu^- W^{+\mu} h^2 \quad (2.12)$$

The second term on the right hand side of Equation (2.12) gives us the coupling strength of a vertex with a Higgs and two W bosons.

$$g_{HWW} = \frac{1}{2}g_W^2 v = g_W m_W \quad (2.13)$$

The coupling to the Z boson can also be found from Equation (2.9) by substituting




Figure 2-1: Above is the Feynman diagram for associated production. The W or Z boson radiates a Higgs boson. Both bosons later decay into particles detected by CMS.

$(v + h)^2$ back in for v^2 and extracting the terms proportional to $hZ_\mu Z^\mu$.

$$g_{HZZ} = \frac{1}{2} (g_W^2 + g'^2) v = \sqrt{g_W^2 + g'^2} m_Z \quad (2.14)$$

When arranged in a way that the W or Z boson radiates the Higgs, as opposed to a Higgs decaying into a pair of W or Z bosons, the process is called associated production or *Higgstrahlung*. The vertex showing associated production is pictured in Figure 2-1. PhilThe above didn't render in my pdf, but maybe thats me

2.2.1 Coupling Between Vector Bosons and Fermions

The W and Z bosons are themselves intermediate states, never existing in a directly observable manner. They must be produced through interactsPhilinteractions with stable fermions. Since the LHC is a hadron collider, considering the vector bosons' couplings with quarks would be most relevant. PhilI would rephrase left

Quarks couple to each other through the strong force, resulting from a $SU(3)$ symmetry. There are three generations of quarks each consisting of a pair of quark types. Their mass eigenstates are denoted as down-type or up-type. A feature of quarks is that their mass eigenstates do not match their weak eigenstates. There is a mixing among the down-type quarks that is parametrized by the Cabibbo-Kobayashi-




Figure 2-2: Above are diagrams for generating W^+ and W^- bosons. The u and d quarks in the diagram can be replaced with any up-type or down-type quark, respectively. The CKM matrix element would in the vertex element would be changed accordingly.

Maskawa (CKM) matrix.

$$\begin{pmatrix} d' \\ s' \\ b' \end{pmatrix} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix} \quad (2.15)$$

The mass eigenstates are denoted as d, s , and b , while d', s' , and b' are the weak eigenstates. This mixing allows quarks to change generations through interaction with W^\pm bosons, which raise or lower the weak isospin. The following is the charge current vertex interaction. **Phil**you are missing the W propagator

$$-i \frac{g_W}{\sqrt{2}} \begin{pmatrix} \bar{u} & \bar{c} & \bar{t} \end{pmatrix} \gamma^\mu \frac{1}{2} (1 - \gamma^5) \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix}$$

The vertices for this interaction is shown in Figure 2-2 arranged in a way to show the processes of generating a W^+ or W^- boson from annihilating quarks. The γ matrices in the interaction are present because the SU(2) component of the Standard Model only interacts with left-handed fermions and right-handed anti-fermions. For this reason, the SU(2) component is more accurately labelled SU(2)_L. From Equation (2.11), the W^\pm bosons are completely made up of the $W^{(1)}$ and $W^{(2)}$ components

of the $SU(2)_L$, so they also only interact with left-handed fermions and right-handed anti-fermions.

Both the photon and the Z boson mix the $SU(2)_L$ and $U(1)_Y$ components of the Standard Model. Production of the Z boson needs to be directly understood for this measurement, but it is more straightforward to determine the strength of the Z boson couplings to left- and right-handed fermions by exploiting the symmetry of photon interactions. That is, the photon interacts the same with left and right handed charged fermions, and not at all with neutral fermions. This is shown directly with experiments with leptons. The charged leptons, electrons, muons, and taus, interact with photons, while the respective neutrinos do not. From the mixing in Equation (2.8), the photon and Z fields can be expressed as the following. PhilThe above paragraph is very confusing, can you rewrite it.

$$A_\mu = B_\mu \cos \theta_W + W_\mu^{(3)} \sin \theta_W \quad (2.16)$$

$$Z_\mu = -B_\mu \sin \theta_W + W_\mu^{(3)} \cos \theta_W \quad (2.17)$$

θ_W is known as the weak mixing angle. The relative strengths of the B and $W^{(3)}$ couplings are determined directly through lepton electromagnetic characteristics, keeping in mind that $W^{(3)}$ only interacts with left handed particles. The following are the electromagnetic interaction strengths of left- and right-handed electrons and neutrinos.

$$e_L : \quad Qe = \frac{1}{2}g'Y_{e_L} \cos \theta_W - \frac{1}{2}g_W \sin \theta_W \quad (2.18)$$

$$\nu_L : \quad 0 = \frac{1}{2}g'Y_{\nu_L} \cos \theta_W - \frac{1}{2}g_W \sin \theta_W \quad (2.19)$$

$$e_R : \quad Qe = \frac{1}{2}g'Y_{e_R} \cos \theta_W \quad (2.20)$$

$$\nu_R : \quad 0 = \frac{1}{2}g'Y_{\nu_R} \cos \theta_W \quad (2.21)$$

Y_{e_L} and Y_{ν_L} must be equal to maintain $SU(2)_L$ symmetry. To satisfy these constraints,




Figure 2-3: Above are diagrams for generating Z bosons. Left- and right-handed fermions are both coupled to, but with different coupling strengths.

the follow definition of Y is needed.

$$Y = 2 \left(Q - I_W^{(3)} \right) \quad (2.22)$$

The following relationship also arises from these experimental constraints.

$$e = g_W \sin \theta_W = g' \cos \theta_W \quad (2.23)$$

Returning to the Z boson, from Equation (2.17), and defining

$$g_Z = \frac{e}{\sin \theta_W \cos \theta_W}, \quad (2.24)$$

we have the following couplings to left- and right-handed fermions.

$$-\frac{1}{2} g' \sin \theta_W (Y_{f_L} \bar{u}_L \gamma^\mu u_L + Y_{f_R} \bar{u}_R \gamma^\mu u_R) + I_W^{(3)} g_W \cos \theta_W (\bar{u}_L \gamma^\mu u_L) = g_Z \left((I^{(3)} - Q \sin^2 \theta_W) \bar{u}_L \gamma^\mu u_L - Q \sin^2 \theta_W \bar{u}_R \gamma^\mu u_R \right) \quad (2.25)$$

Now the coupling of the Z to left- and right-handed quarks can be calculated from Table 2.1, remembering that $I_W^{(3)}$ for right-handed fermions is 0. Diagrams showing the interaction strengths of fermion- Z vertices are shown in Figure 2-3.

Thus vector bosons couple to quarks, the constituents of hadrons, which means they can be produced at the LHC. As mentioned earlier in this section, quarks interact

through an SU(3) symmetry that results in the strong force. The three states that this symmetry supports are known as color states, and they are labelled red, green, and blue, or r , g , and b . There are also anti-states for each color state, labelled \bar{r} , \bar{g} , and \bar{b} . The resulting gauge bosons are known as gluons, and they carry the following color states.

$$r\bar{g}, g\bar{r}, r\bar{b}, b\bar{r}, g\bar{b}, b\bar{g}, \frac{1}{\sqrt{2}}(r\bar{r} - g\bar{g}) \text{ and } \frac{1}{\sqrt{6}}(r\bar{r} + g\bar{g} - 2b\bar{b})$$

Since gluons carry color charge, they interact with other gluons. As the distance between two color-charged particles grows, the energy density of the self-interacting gluon field remains constant. It soon becomes energetically favorable for new a particle/anti-particle pair to pop into existence if it simultaneously reduces the distance that the strong force is interacting. As a result, all observable hadronic states are color singlets. The most common hadronic states are mesons, made of a quark/anti-quark pair with the color singlet state

$$\psi(q\bar{q}) = \frac{1}{\sqrt{3}}(r\bar{r} + g\bar{g} + b\bar{b}), \quad (2.26)$$

and baryons, made of three quarks with the following color singlet state.

$$\psi(qqq) = \frac{1}{\sqrt{6}}(rgb - rbg + gbrgrb + brg - bgr) \quad (2.27)$$

Baryons can also be composed of three anti-quarks, which has a state corresponding to Equation (2.27), but with anti-color. The resulting spray of hadronic particles generated from the vacuum to restore color singlets are called jets. **Phil**I think your definition of jets is a bit liberal, I would say spray from a high energy quark, high energy is the key here.

For this measurement, protons are collided at the LHC. The proton consists of two u quarks, and one d quark. Since the three quarks inside the proton interact strongly, there are also many virtual gluons and quark/anti-quark pairs present at all times. The quantity and energies of all these partons are not able to be calculated




Figure 2-4: The Parton Distribution Function for protons is shown above. Most of the proton's momentum is carried by u and d quarks, but virtual s quarks as well as gluons can also interact with particles passing through the proton.

since QCD is non-perturbative. They can be measured in deep inelastic scattering experiments though. In these, electrons are scattered off of protons, and parton distribution functions (PDFs) can be measured. The PDFs for protons are shown in Figure 2-4.

Combining the known proton energy, PDFs, the CKM matrix, and the theory of the electroweak force, we can predict the cross section of generating W and Z bosons at the LHC. These initial vector bosons will be off-shell, which means they will have a mass much different than the resonance peak. Then they will radiate a Higgs in order to most commonly produce an on-shell vector boson and on-shell Higgs boson. The cross section of generating off-shell particles are suppressed according to the required center-of-mass energy, E , and the resonance mass M . The suppression is in the form of the relativistic Breit-Wigner formula.

$$f(E) = \frac{k}{(E^2 - M^2)^2 + M^2\Gamma^2} \quad (2.28)$$

Phil Maybe you are getting to it, but I think its important to discuss what is offshell and what isn't, why is the Higgs not offshell for example. One way around this is




Figure 2-5: Above are the Feynman diagrams for other production mechanisms of the Higgs boson. Gluon fusion is shown on the left, and vector boson fusion is shown on the right.

to write the whole ME for the production This associated production is one of three production mechanisms of the Higgs Boson. The other two are gluon fusion, where gluons form a top loop, and vector boson fusion, both shown in Figure 2-5. In these other two production mechanisms, only the Higgs is in the final state. These events can only offer additional identification through initial state radiation. In contrast, associated production also results in leptons from the vector boson decay, which allow for tighter selection criteria for event identification.

2.2.2 Decay Channels of Vector Bosons

Due to the couplings described in Section 2.2.1, the vector bosons decay into quarks. **Phil**decay predominantly into However, in the hadronic environment produced at the LHC these are not the best indicators of a vector boson intermediate state. This measurement uses leptonic decays in the final state since they are easier to identify and separate from background processes.

There are three generations of leptons. **Phil**This paragraph feels out of place, I think you said most of this, can you push this up to earlier Each generation consists of a charged lepton, and a neutral lepton, also referred to as a neutrino. The left-handed charged lepton and neutrino of each generation form an electroweak $SU(2)_L$ doublet. In order of increasing mass, the three generations are called electron, muon,




Figure 2-6: Heavier leptons can decay to lighter leptons while emitting two neutrinos. Above is an example of a decay of $\tau \rightarrow \nu_\tau \mu \bar{\nu}_\mu$.

and tau. Heavier charged leptons decay into lighter leptons via the weak force. Two neutrinos result from this decay, as shown in Figure 2-6, making the characteristics of the parent lepton's parent difficult to reconstruct. The tau lepton has a short enough lifetime to consistently decay before reaching the CMS detector. The tau lepton is also massive enough to also decay into quarks, making its measurement even more complicated. PhilI would also mention their are neutrino decays here too Muons have an average lifetime long enough to penetrate the entire detector, and electrons are stable particles. As a result, only final states with muons and electrons are considered in this analysis. The Feynman diagrams for the decay channels of interest are shown in Figure 2-7.

2.3 Decay Channels of the Higgs

What we are ultimately interested in measuring is the contribution of the Higgs intermediate state to the final state of $b\bar{b}$. Since the Higgs is a $SU(2)_L$ doublet of scalar fields, the term $-g_f(\bar{L}\phi R + \bar{R}\phi^\dagger L)$ in the Standard Model Lagrangian is invariant under $SU(2)_L \times SU(1)_Y$, where L is a left-handed fermion doublet, and R is a right-handed singlet. If the Higgs doublet is expanded around the vacuum




Figure 2-7: Above are the three different vector boson decays we are interested in. τ decays do also contribute to the charged lepton final states as seen by a detector, but the energy carried away by neutrinos significantly reduces those decay modes' contribution to the accepted states.

expectation value, as Equation (2.5), the Lagrangian term becomes the following.

$$\mathcal{L}_f = -\frac{g_f}{\sqrt{2}}v(\bar{f}_L f_R + \bar{f}_R f_L) - \frac{g_f}{\sqrt{2}}h(\bar{f}_L f_R + \bar{f}_R f_L) \quad (2.29)$$

In Equation (2.29), f refers to the lower field of the fermion's $SU(2)_L$ doublet. The Lagrangian also includes terms for the upper field since the conjugate of ϕ has the same symmetries as ϕ . **Phil** I think you should put this part earlier when you discuss the Higgs mechanism in vector bosons, also, you should put a decay table or plot of decays so its clear why $b\bar{b}$ is important

The Lagrangian showing fermion-Higgs interactions in Equation (2.29) consists of two terms. Since v is constant, the first term is consistent with a fermion's mass, assuming an appropriate coupling constant.

$$g_f = \sqrt{2}\frac{m_f}{v} \quad (2.30)$$

The second term is the coupling of the fermion to the Higgs field with the same coupling constant. This is the mechanism by which the Higgs give fermions their masses, and also why the Higgs couples more strongly to massive particles. The Feynman rule for the interaction vertex between the Higgs and fermions is proportional to the fermion's mass. Of the quarks, the b quark is the second most massive. The most massive t quark is too massive to be the final decay product of an on-shell Higgs. The required off-shell Higgs would need a center of mass energy, E , at least 350 GeV, and the cross section for this production drops off as the tail of a relativistic Breit-Wigner function in Equation (2.28) with a mass M of 125 GeV. The Higgs can also decay to two vector bosons. In this case, instead of requiring the Higgs to be off-shell, one of the unstable vector bosons can be less massive than its resonance mass. However, this still results cross section suppression in the form of Equation (2.28).

As the decay mode with the highest coupling requiring no off-shell particles, the predicted branching ratio of $H \rightarrow b\bar{b}$ is 57.8%. Therefore, measuring $H \rightarrow b\bar{b}$ is the most direct measurement to confirm this theory of quark masses. The diagram for this decay can be combined with the Feynman diagrams in Figure 2-1, Figure 2-2




Figure 2-8: Above is the full Feynman diagram for $ZH \rightarrow \ell^+ \ell^- b \bar{b}$.

or 2-3, and one of the decays in Figure 2-7 in order to generate the full Feynman diagrams for the processes being measured in this analysis. One such full diagram is shown in Figure 2-8.

Phil This section is a bit jumbled, I think it needs some reorganization of the material, you mostly cover everything needed to explain the process but it is a bit hard to follow. Lastly, you SHOULD add a new sub section here about why it is important to measure $VHbb$, what do we learn about the Higgs? Is there anything interesting about the kinematic dependence? Are there other alternate Higgs theories. Point is you should explain why this measurement is so important.

Chapter 3

The CMS Detector

The Compact Muon Solenoid (CMS) detector, located at the LHC, consists of multiple sub-detectors. The analysis in this work is quite complex, and depends on all parts of the detector. Therefore, a full description of CMS is presented in this chapter.

First, a brief description of the LHC is given in Section 3.1. Then design requirements and considerations are outlined for the CMS detector in Section 3.2. Specific design decisions and descriptions of subdetectors are given in Section 3.3. Section 3.4 describes event reconstruction algorithms, Section 3.5 describes the triggers used to collect data, and Section 3.6 outlines simulation techniques used for CMS. Finally, Section 3.7 describes how data is stored and accessed by members of the collaboration. More can be learned about the design and motivations for the detector in the TDR [16]. Information presented on the physical CMS design parameters are taken directly from that document unless otherwise noted.

3.1 The Large Hadron Collider

The CMS detector only observes events. Before describing the devices that are used to observe and record events, the method of generating interesting events must be described. The CMS detector is located at the Large Hadron Collider (LHC). Described in detail in multiple publications [17], a brief description is given here.

The LHC, with a circumference of 26.7 km, is large enough to be considered

located in multiple towns and countries, but it will suffice to say it is near Geneva, Switzerland at the European Organization for Nuclear Research (CERN), the main campus of which is addressed in Meyrin, Switzerland. This campus itself also spans the border between Switzerland and France. This large circumference is needed since charged particles traveling in a circular path with radius r emit synchrotron radiation at the following rate.

$$P = \frac{q^2 p^4}{6\pi\epsilon_0 m^4 c^5 r^2} \quad (3.1)$$

The amount of power lost by the particles decreases quadratically with the size of the collider. In addition, the energy lost decreases with the mass of the accelerated particles to the fourth power. The LHC was built in the same tunnels that were used for LEP, which was a collider for electrons and positrons that took much of its data at $\sqrt{s} = 91$ GeV in order to study the Z boson resonance. The resulting LHC is designed to collide protons at energies of $\sqrt{s} = 14$ TeV, with the data for this analysis taken at $\sqrt{s} = 13$ TeV. This is more than enough energy to generate the massive off-shell vector bosons that are needed for *Higgstrahlung*, as well as many accompanying jets, via the mechanisms described previously in Chapter 2.

The luminosity of the LHC is given by the following formula.

$$\mathcal{L} = \frac{N_B^2 f_{\text{rev}} k_B}{4\pi B^* \epsilon_{xy}} \times F \quad (3.2)$$

N_B is the number of protons per bunch, f_{rev} is the frequency of beam revolutions, k_B is the number of bunches per beam, B^* and ϵ_{xy} describe the goodness of the beam, and F is a geometric collision factor.

$$F = \frac{1}{\sqrt{1 + \frac{(\sigma_s \tan \phi)^2}{\epsilon_{xy} \beta^*}}} \quad (3.3)$$

σ_s is the length of each bunch, ϕ is the crossing angle, and β^* is the value of the amplitude function at the focal point. In order to generate as much collision data as possible, the LHC operates at a high frequency of collisions, and generates many

simultaneous collisions. For Run 2, there is a proton bunch crossing every 25 ns. The CMS detector must be able to read out and process data on that timescale. Each proton bunch includes over 100 billion protons [18].

3.2 Detector Requirements

One configuration of possible final state particles was shown previously in Figure 2-8. There, two oppositely charged leptons and two b quarks are the end decay products. The b quarks also hadronize **Phil**to form color singlets well before reaching the detector, but the resulting jets can actually be distinguished well from the jets resulting from the fragmenting protons.

Hadrons containing b quarks decay through the weak force since they require a flavor change. As mentioned before, the CKM matrix in Equation 2.15 quantifies the mixing between the different quark flavors. The value of V_{tb} is close to unity, and since the CKM matrix is unitary, V_{cb} and V_{ub} are small. **Phil**wonder if you want to put thi sin the theory section, also good to mention why b-quarks don't decay into tops + stuff This means the matrix element weak decays of the b hadrons is small. This is the only decay channel available to the lightest b hadrons, so their lifetimes are relatively long. The delayed decay results in a jet with a secondary vertex where many of its particles are generated from the vacuum at a distance from the initial collision point.

Alternate signatures of interest can be seen by substituting other vector boson final states from Figure 2-7. In these, there may be one or zero charged leptons, with one or two neutral leptons, respectively. Neutral particles are difficult to detect, with neutral leptons being capable of passing through the entire Earth without being part of a detectable interaction. The CMS detector therefore ignores the neutrinos, but their presence can still be inferred. Even with the variation in momentum along the beam direction, all partons in each proton have approximately zero momentum in the transverse direction. Therefore, the sum of the transverse momenta of all final state particles must also be zero. Many events in CMS have an overall imbalance in

the transverse plane. This imbalance is labelled Missing Transverse Energy, E_T^{miss} , or MET. Large MET in an event is often a sign of high energy neutrinos that the detector cannot detect. **Phil**split this paragraph into two, MET deserves its own paragraph, rework some of the explanation

We need to identify all of these interesting particles, as well as be able to reconstruct missing transverse momentum. In addition, the additional hadronic activity in the event, called pileup, must be mitigated. The energy of the decay products have energies on the scale of the masses of the parent particles. The detector must be capable of measuring jets and leptons with energies on the order of 10s or 100s of GeV. Better energy resolution for each of these decay products allows better separation of our signal process from background processes that generate very similar final states.

3.3 Detector Design

The CMS detector as a whole has cylindrical symmetry around the proton beams. It is 21 meters long and 15 meters in diameter. There are gaps at either end to allow the beams to enter and leave, but otherwise the design tries to cover the full solid angle around the collision point. The azimuthal angle of a particle relative to the beam axis is described by pseudorapidity, η .

$$\eta = -\ln \left[\tan \left(\frac{\theta}{2} \right) \right] \quad (3.4)$$

The barrel portion of CMS detects particles up to $|\eta| < 1.5$, while the forward caps of the detector can reach $|\eta| < 5.0$. The muon and pixel trackers reach up to $|\eta| < 2.5$, with additional space covered by calorimetry. **Phil**Why just pixel? where are the strips? I spent 2 years of my life plugging in power cables to the strips and now they are forgotten :(

Different technologies are better for measuring the energy or other kinematics variables of different particles. As a result, the CMS detector is made up of different sub-detector systems, arranged in cylindrical layers. Each layer consists of a “barrel”

portion and two end caps on either side.

The innermost sub-detector is designed to extrapolate the tracks of charged particles back to their point of origin. This is called the Silicon Tracker. One key design feature of the pixel detector is that it is non-destructive. Particles it detects pass through to the rest of the detector for additional measurement and identification. The next sub-detector encountered by most particles is the Electromagnetic Calorimeter, which is designed to measure the energies of photons and electrons. The next sub-detector, the Hadronic Calorimeter, measures the energies of both charged and neutral hadrons. The two calorimeters are destructive because **Phil**they absorb the particles that interact with them in order to measure their full energy. Outside of these three sub-detectors is a superconducting solenoid, which generates a magnetic field for the entire detector. On the very outside of the detector are gas chambers designed to detect muons interspersed with the iron return yoke for the solenoid. A slice of the CMS detector showing the relative positions of each layer is shown in Figure 3-1.

The magnet is described first since the magnetic field it produces is a key part of most of the rest of the detector. After that, the sub-detectors are summarized in the order of closest to farthest from the beamline, since this is the order that particles would interact with the layers. Each sub-detector section also describes the measured performance during Run 2. Note that this measurement is an iterative process that depends on the event reconstruction described in Section 3.4, which in turn depends on the performance of the entire detector. The performance numbers are presented with each sub-detector design though so that it is immediately clear how effective each design has been.

3.3.1 Solenoid Magnet

Part of the CMS acronym acknowledges the role of the solenoid magnet. **Phil**This is a weird sentence for a non-expert The presence of a magnetic field is paramount for accurate measurements of charged particles passing through the silicon detector and the muon chambers.




Figure 3-1: A slice of the CMS detector is shown above [19]. The four detector layers are labelled and show the penetration depths of various particles stable enough to travel a measurable distance.

The magnetic field generated is designed to cause the path of a muon with 1 TeV of energy to bend enough to have a momentum resolution of 10%. Inside the solenoid, the magnetic field operates at 3.8 T, with the solenoid design being capable of achieving 4 T. The return field is large enough to cause muon tracks to curve throughout the muon chambers outside the magnet.

A super-conducting solenoid enables the creation of a magnetic field with the required strength. A current of 19.5 kA is sent through 2168 turns over 12.9 m. The magnetic field stores 2.7 GJ of energy. In order to hold this, the structural components holding the magnet and the detector in place are strong enough to withstand 64 atm of hoop pressure.

3.3.2 Silicon Tracker

The layer closest to the beamline is designed to obtain a precise track pointing to the origin of particles passing into the detector. It is made up of layers of many small

pixels to do this. As distance from the interaction point increases, the pixel size also increases since the absolute spatial resolution does not need to be as fine.

The innermost three layers, with the closest layer being a distance of $r = 4$ cm from the interaction point, are made of hybrid pixel detectors. Each pixel has dimensions of $100 \times 150 \mu\text{m}$ in order to achieve fine resolution of where the detected particles originated. The TDR also claims an occupancy of 10^{-4} per pixel per LHC bunch crossing, which improves the pixel's longevity and reduces problems from detector downtime. Outside of the pixel detector layers, silicon strip detectors are used. Finally strips, I would rework above These are placed in the region that is $20 < r < 55$ cm from the beamline. Strip dimensions give a cell size of approximately $10 \text{ cm} \times 80 \mu\text{m}$. 2–3% of cells are activated during a typical bunch crossing. The outermost layers are made of larger strips with cell sizes of $25 \text{ cm} \times 180 \mu\text{m}$. About 1% of these pixels are triggered each bunch crossing.

The active material of the silicon pixel detector is semi-conducting silicon. When charged particles pass through, electron-hole pairs are generated and drift apart due to a bias voltage. The voltage change when these pairs reach their respective electrodes indicates a charged particle passed through. Because of this, the silicon pixel detector cannot detect neutral particles, but it gives a precise point of origin for charged particles. The points of origin allow for the determination of locations of primary and secondary collision vertices, which plays an important role in the identification of pileup.

In the beginning of 2017, the pixel detector was upgraded to handle the higher radiation environment of Run 2 [20]. A layer was added to both the barrel and endcap sections of the pixel detector. Firmware was also upgraded to keep the pixel detector operating at a frequency higher than the Run 2 collision frequency. With this upgrade, the detector operated with a 97% hit efficiency for all layers at the highest instantaneous luminosity. Layers beyond the first performed with greater than 99% hit efficiency [21].

3.3.3 Electromagnetic Calorimeter

The next layer of the detector is called the Electromagnetic Calorimeter or ECAL. This layer is designed to fully capture and accurately measure the energy of photons and electrons. The ECAL is made of crystals of the scintillating material Lead Tungstate (PbWO_4). Each crystal is placed in the detector so that its smallest face is facing the collision point. These small faces have dimensions of 22×22 mm. The length of each crystal is 230 mm, and the far face is slightly larger at 26×26 mm. PbWO_4 has a radiation length of $\chi_0 = 8.9$ mm and a Moliere radius of 21 mm. This means each crystal is 25.8 radiation lengths, containing the full shower within the ECAL, and each shower is also localized to within one crystal from the initial ionization. There are gaps in active detecting volume of the ECAL, which are needed to accommodate various electronics and structural components. The gaps are located in symmetric locations on either side of the ECAL from $|\eta| = 1.4442$ to $|\eta| = 1.5660$.

The scintillating properties of PbWO_4 are also desirable for observing LHC collisions. The photodiodes at the far end of the crystals ultimately detect 4.5 photons for every MeV of energy deposited in the ECAL. This is a low number for other experiments, but the only photons and electrons of interest in this measurement deposit at least tens of GeV of energy. This gives the ECAL energy resolutions in the range of 5 – 10%. More importantly, the scintillation is very fast. About 80% of the light from an interaction is emitted within the 25 ns between bunch crossings, making it easy to associate the readouts with the appropriate bunch crossing.

When exposed to the high radiation environment of the LHC, the ECAL crystals are damaged by radiation. Damage to the crystal structure causes it to become more opaque to the scintillated light. Much of this damage happens within the first 30 minutes of operation. Some recovery occurs as the crystal structure falls back into the ground state, but over time, the performance of the crystals degrades. That degradation happens at different rates in different areas of the detector, but, aside from the initial darkening, is slow enough to be able to correct for it during the run. Lasers are used to calibrate the ECAL online during the gaps between

beams [22]. Resolution is measured by looking at $Z \rightarrow ee$ events. For Run 2, the barrel region of the ECAL performed with 1.6% resolution, and the other regions had a 5% resolution [23].

3.3.4 Hadronic Calorimeter

The ECAL absorbs electromagnetic particles and measures their energies which are dispersed in electromagnetic showers. **Phil**I think you need a qualifer in the first sentence. Unlike the ecal, which..., the Hcal Hadrons deposit energy in hadronic showers, which require a different mechanism to contain and measure. The Hadronic Calorimeter (HCAL) does this. Like the ECAL it contains particles and measures their energy destructively. However, it does this for hadrons, such as protons, neutrons, and stable mesons. Since hadrons are much more massive than electrons, the ionizing collisions in a typical scintillator does not slow them down enough to contain them. Instead, they must interact via nuclear collisions to be attenuated. CMS uses brass for its HCAL due to its relatively short interaction length, the fact that it is non-magnetic, and its affordability.

The barrel of the HCAL is jacketed in stainless steel for structural support. This layer is 61 mm thick on the layer immediately next to the ECAL and 75 mm thick on the outer edge. The inside of the HCAL consists of brass absorber plates interspersed with plastic scintillator tiles. The layers closer to the beamline alternate 50.5 mm brass plates with 3.7 mm scintillator plates. Farther away, the brass plates are instead 56.5 mm thick. Wavelength shifting fibers are run through the scintillator tiles to allow photons to travel to the outside of the HCAL where they are detected by photodiodes.

Like the ECAL, the HCAL performance also degrades as it is exposed to radiation. The calibration for HCAL is performed using an embedded radioactive source, lasers and LEDs, and an *in situ* calibration using assumed symmetry in ϕ . With these methods, a response within 3.4% was maintained in the HCAL barrel and within 2.6% in the HCAL endcap up to $|\eta| < 2$ [24].

3.3.5 Muon Chambers

Muons are the most penetrative particles that CMS detects. Through the calorimeters, muons act as minimum-ionizing particles [25]. They are heavier than electrons, so they are not stopped in the ECAL. They do not interact via the strong nuclear force, so the high density of the HCAL also does not cause significant interactions. Instead of stopping and measuring muons in calorimeters, CMS tracks their trajectory with both the silicon tracker on the inside of the detector and the muon chambers that make up the outer layer of the detector.

This is the only sub-detector system outside of the solenoid, but the returning magnetic field is still present outside of the return yoke [26,27], allowing the momentum of the muons to be extracted from the curvature of their trajectory. Layers of muon chambers act much like the pixel detectors, but at a larger and more distant scale. The muon chambers in the barrel region of $|\eta| < 1.2$ consist of drift tube chambers. In the endcaps, cathode strip chambers are used. The difference is to account for higher neutron backgrounds in the endcap, as well as a greater magnetic field. In both regions, resistive plate chambers are spaced between the layers of the other muon chambers.

Each muon chamber has a detection efficiency greater than 95%. The overall efficiency of the muon trigger, which relies heavily on the muon system and is described in more detail in Section 3.5, increases as a function of muon p_T and plateaus around 90%. The timing of the muon system leads to 1% of muons to be assigned as originating from the wrong bunch crossing [28].

3.4 Event Reconstruction

Each sub-detector reconstructs the particles that passes through it. The independent reconstructions are then linked across the different detector components to identify particles. This overview is taken largely from reference [29]. **Phil**This better be in your own words.

3.4.1 Charged Particle Tracks

Both the Silicon Tracker and the Muon Chambers are designed for charged particles to leave tracks. In both sub-detectors, the basic steps for track reconstruction are the same. First, a track must be seeded. Usually, this is done by finding hits in consecutive layers that are consistent with a particle coming from the beamline. Particles lose energy as they pass through matter, and they can also be redirected through multiple scattering, so the extrapolation is non-trivial. A Kalman Filter is used to find hits in the other layers of the appropriate sub-detector that are consistent with the initial seed. Once more hits have been found, a fit is performed for the precise trajectory of the track.

Tracks are kept or discarded based on the number of layers that are missing hits and on the momentum of a charged track in the magnetic field of the solenoid. Making these parameters looser results in better recovery of tracks, but the high activity within the detector results in a combinatorial background. This background increases exponentially when the momentum cut is reduced, for example. To help reduce this background, tracks with missing hits use an iterative fit. Different seeds are found for each track to make sure that the resulting collections of hits remain the same.

Additional complications arise for each the pixel and muon detectors. The Silicon Tracker is the only tracking detector that deals with electrons. Because of their small mass, electrons are likely to radiate energy while travelling through the magnetic field of CMS. This leads to complications in the calorimeters described in Section 3.4.2. It also means that the radius of curvature of an electron track can decrease appreciably within the pixel detector. This can lead to the Kalman Filter approach missing tracks entirely, depending on the number of hits required. Tracks with a large χ^2 and a certain number of hits are fit again using a Gaussian-Sum Filter (GSF). The GSF allows for fitting tracks that have significant energy loss, recovering electron and positron tracks.

A change of trajectory may also happen in the muon chambers, but this is due

to multiple scattering in the return yoke. No specialized tracking algorithm is used to account for it. The muon tracking performs best when the tracks in the muon chambers are successfully linked to a track in the pixel detector. The most common backgrounds in the muon chambers is caused when hadrons manage to punch through the HCAL. This is often mitigated by considering the amount of energy deposited along the particle track in the other sub-detector systems.

3.4.2 Calorimeters

Clusters are identified in calorimeters, also using a seeding algorithm. First clusters with a large energy deposit are identified, and then nearby crystals are checked against noise thresholds. The energy deposition is assumed to have a Gaussian profile, and a fit is performed to disentangle overlapping energy depositions.

The dimensions of the ECAL crystals are comparable to the Moliere radius, keeping clusters localized. Although a complication arises due to curvature of the electromagnetic shower caused by the magnetic field, leading to the need for using GSF to find tracks. Superclusters that are linked to electrons have a larger allowed range over ϕ to account for this.

The calibration of HCAL is complicated by the fact that particles reaching it have to first pass through the ECAL. Initial calibration was done with a 50 GeV pion test beam, but the actual response is non-linear in energy as well as different for charged and neutral particles. Reasons for this difficulty include particles losing energy in the region between the ECAL and the HCAL, in addition to the energy lost in the ECAL. Therefore, there are calibration coefficients that are used depending on if the energy deposits are all in the HCAL, or in preceding sub-detectors as well.

3.4.3 Linking and Particle Identification

An important step in making sense of the various sub-detector readouts is linking tracks to calorimeter clusters. The general procedure is to extrapolate tracks from the inner tracker out to each calorimeter. A shower that originates within one radiation

or interaction length in the calorimeter along that track is linked with the track.

The bremsstrahlung from GSF electrons is linked to the track by looking along track tangents. A dedicated conversion finder is used to identify pair production within the pixel detector caused by either bremsstrahlung or prompt photons in order to not mistakenly link a charged particle track with what should otherwise be measured as a photon. This step, in addition to some ECAL clusters that do not have a track make it possible to identify isolated photons. On the other hand, it is still difficult to determine whether an electron track is well isolated or not. The large number of variables that go into identifying an electron leads to the training of a Boosted Decision Tree (BDT) to identify an electron. Separate BDTs are needed for the barrel and endcap regions of the detector.

For accurate HCAL readings, the linking algorithm also ECAL clusters and HCAL clusters along a path. These may not always be along a charged particle track. Multiple calorimeter links of this nature may be found, but only a single link is kept based on a distance assigned to each link. HCAL hits without a track are identified as neutral hadrons. HCAL hits with a linked track are likely charged hadrons. Though the ECAL clusters must be linked in order to determine the energy coefficient to calibrate the HCAL, ECAL hits without tracks are still identified as photons because photons carry some of the energy of jets.

Of particular interest to this analysis is also the secondary vertex step of linking. Charged particle tracks that do not go back to the interaction vertex are linked together if they share a common secondary vertex. These tracks must have a mass greater than 0.2 GeV to be kept. There must also be a track from the secondary vertex to the primary vertex, which would belong to a long-lived hadron. As mentioned in Section 3.2, this is the signature of a b jet. It is possible, however, that the secondary vertex is generated by a nuclear scattering, pair production, or other long-lived particles like K_S or Λ within the pixel detector, so additional analysis is needed for each secondary vertex.

The final link is made between tracks in the muon chambers and tracks in the inner tracker. Muons are identified as tracker muons if they only leave tracks in the inner

tracker. This can often happen with low energy muons. They are called standalone muons when only the track in the muon chamber is identified. When tracks are successfully linked in both sub-detectors, the resulting reconstructed particle is called a global muon. When a global muon is not well-isolated from other energy deposits, it must have tighter requirements on how it behaves in the muon chambers. This is to prevent energy from a jet from being attributed to a muon or vice versa. This is important for b jets because the decay that happens at a displaced vertex is a decay through the weak nuclear force, which can result in leptons being present inside of a jet.

3.5 Trigger

Bunch crossings happen every 25 ns, with each bunch crossing producing on average 20 collisions. The amount of data that the detector generates for each bunch crossing is too large to store all of it at this rate. Luckily, most collisions result in uninteresting data. PhilYou should use an euphemism for uninteresting A trigger system is used to identify interesting events and reduce the frequency of event writing to 1 kHz. This is done using two stages. The Level-1 (L1) trigger passes events with a frequency of 100 kHz, and the High Level Trigger (HLT) picks from the remaining events with a frequency of 1 kHz [30].

The L1 trigger is implemented in hardware. It was upgraded for Run 2 of the LHC to run on FPGAs on an Advanced Mezzanine Card. There are two main components of the L1 trigger. One considers calorimeter deposits, and the other examines the muon chambers. The overall L1 trigger fires when there are high energy, resolved calorimeter hits or if a possible muon is reconstructed. Due to the flexibility of FPGAs, the exact conditions of the firing are configurable [31].

After the L1 is fired, the data is sent to the High-Level Trigger (HLT), which is a computing farm that makes a final decision on whether or not to save the data using a rough event reconstruction. The use of 30,000 cores in the HLT allows for buffering data so the HLT has plenty of time to make this decision [32].

For this analysis, only a few of the possible HLT paths are of interest. The exact trigger names are given in Chapter 4, but for the most part, they only depend on three different identifiable objects. Figures 2-7 and 2-8 show the different final states of interest. b jets are difficult to identify quickly because we must rely on the Silicon Tracker’s reconstruction of the secondary vertex, but the decay mode of the vector boson can be used for the trigger. More boosted vector bosons leave a signature with a higher trigger efficiency. They also will cause the b jets to have a higher p_T , leading to easier identification and measurement. In that case, only events with one of the following are worth saving and examining for $VH \rightarrow b\bar{b}$.

- ECAL deposits consistent with a high p_T electron
- muon chamber hits consistent with a high p_T muon
- an overall energy imbalance consistent with MET from neutrinos

For the specific decay channel in Figure 2-8, the HLT also includes paths where two electrons or two muons are identified.

3.6 Simulation

After the detector is well understood, predictions on how it responds in the LHC environment can be made. The number of ways the detector could possibly respond are nearly infinite. Therefore, simulation is performed using Monte Carlo methods, and the resulting analysis is statistical in nature. The data format for simulation results is similar to the data format for data collected from the detector. Unobservable information about intermediate steps in the simulation is also stored, but otherwise the data is the predicted output of a collection of events.

The simulation itself consists of several steps each outlined in a separate section of this chapter. First the background processes that will appear in our analysis must be known. Identifying all of these processes is necessary to quantify and characterize the signal events that are also mixed in to our selection. Then each of these processes

must be simulated to determine the final state particles that the detector will observe. Each process will look slightly different in our signal selection. **Phil** will look => look. Events outside of the selection must also be simulated so that they can be studied for accuracy in separate phase spaces that do not include the Higgs events. After the final state particles are predicted, the detector response to those particles passing through must be simulated. This allows researchers to compare the physical readouts they can observe to predicted detector signals. Finally, using the phase spaces outside of the desired signal process, minor corrections to the simulation can be made. Simulated energies from the detector model might not be the exact same as what the physical detector produces, for example, and they must be made to match to make the signal process cleanly appear in the analysis.

The physical processes that occur at the LHC all contain QCD-driven phenomena. As a result, the part of the simulation that predicts the particles present in the detector has two distinct parts. QCD is perturbative at small distances, and other forces are perturbative at all distances. The collisions themselves are in this regime, so the initial- and final-state particles over a distance of femtometers can be simulated using typical calculations using perturbative rules described by Feynman diagrams. The decay of unstable particles can also be simulated this way. Once particles interacting through QCD exceed this distance, well before reaching the detector, hadronization, or parton showers, must be simulated differently. The following two sub-sections describing these techniques. The exact generators and configurations used to simulate each process for this analysis are detailed in Appendix B. After final state particles are generated, their propagation through the detector is simulated in a third step.

3.6.1 Short-Scale Simulation

Events are generated by selecting results and assigning weights in a way proportional to the phase space and the matrix element squared of the event. The phase space

integral has the following form [33].

$$\int d\Pi_n = \left(\prod_f \int \frac{d^3 p_f}{(2\pi)^3} \frac{1}{2E_f} \right) (2\pi)^4 \delta^{(4)}(P - \sum p_f) \quad (3.5)$$

P is the total initial 4-momentum and f runs over all final state particles. This phase space integral is Lorentz invariant. Once an event has been selected, the available phase space can then be used to assign directions to the final particles.

The proportional matrix elements are described in Chapter 2, but most of the diagrams described were Leading Order (LO). Generators used in this analysis can also simulate Next to Leading Order (NLO) processes thanks to the FKS method of subtracting particles to avoid double counting them during the showering calculation. **Phil**FKS needs a citation However, the option to use NLO simulation is not always used. The results of NLO calculations more accurately predict physical processes, but they also take more computational resources, resulting in larger measurement uncertainties due to statistical limitations.

The generators used for short-scale simulation in this analysis are POWHEG [34] and MadGraph5 [35].

3.6.2 Parton Showers

The final state particles from the short-scale simulation include a number of free quarks and gluons. As mentioned in Section 2.2.1, it is energetically favorable for color-charged particles to create additional quark/anti-quark pairs to screen the color charge. This is known as hadronization or parton showering and is simulated separately from the calculation of tree-level processes. Hadronization happens well before particles reach the CMS detector, so the results are needed to predict the detector response. Accurate simulation of this process is important for all collisions at the LHC, which produces much QCD background. It is also important to accurately simulate the constituents of individual jets because this analysis includes detailed inspection of each jet in order to identify b -jets and to estimate the amount of energy carried away by neutrinos.

To be able to analyze the simulation in the same way data is processed, Monte Carlo simulation is used to predict precise final states of the jets. CMS uses the Lund model [36] as implemented in `PYTHIA8` [37].

3.6.3 Detector Simulation

After determining all of the final state particles that will reach the detector, the interaction between these particles and the detector components must be simulated. Multiple simulations of pp collisions are combined to simulate pileup, and then the particle propagation through various materials is done with `GEANT4` [38]. The full CMS detector geometry is maintained within `CMSSW` [39] using a framework written in the Unified Modelling Language [40]. To be able to process the simulated data in the exact same fashion as the measured data, the readout of the electronics is also simulated.

3.7 Accessing Data

The final important piece of the CMS detector is its offline computing resources. The data that is gathered by the detector must be processed and stored. This is done by using computing resources spread around the world. They are grouped into Tier-1 and Tier-2 sites, based largely on the geographic space that they are meant to provide computing services for. Tier-1 sites typically provide 30,000 CPU cores, while the more numerous Tier-2 sites provide another 60,000 [41]. Together they also provide around 100 Petabytes of space [42]. Keeping these services running smoothly is important for the CMS collaboration to function, and two projects that aid in this are outlined in Appendix A. Since the data is event-based, they are stored in n-tuple format. **Phil** put a reference or describe what an n-tuple is. N-tuple is a bit jargony. These n-tuples are created and read by the CMS Software (`CMSSW`) [43].

Phil This section is overall good, its better towards the end, don't think I have any major comments about what is missing. I am glad you put your work in the appendix, otherwise, I would have made a comment about putting your work in this

part of your thesis.

Chapter 4

Event Selection

This chapter gives the specific selection requirements on each physics object that allows us to count particle candidates and to reject or otherwise classify events, based on the description of physics processes described thus far. First, objects are defined in terms of variables and particle candidates provided by the detector reconstruction algorithms. Then selection requirements based on these objects used to reject events entirely from the analysis are given. **Phil**rewrite this sentence, it sounds like German translated to english directly, its not english After that, selection requirements used to classify events into different decay channels of the vector boson are specified. **Phil**ditto There are also selection requirements that allow events to be treated separately when the Higgs decay products can be resolved as separate jets and where they are merged into a single massive jet.

4.1 Object Definitions

Detector responses are linked to possible physical particles. Most of the particle ID techniques described so far can give false positives for individual particle candidates or provide composite physics objects that are in reality composed of background particles. What follows are tighter selections used in order to reduce these backgrounds. Once objects are more strictly defined, they can be used for more reliable event classification.

Each type of object generally has a method of loose pre-selection and additional tighter requirements for a selection. The distinction is particularly useful for categorizing events. Each category is designed to be enriched with a particular physics process. Each physics process would result in certain final states with specific multiplicities for some particles. For objects, passing the loose selection often means that the object is defined well enough to veto events for inclusion in categories that would not include the corresponding particle. **Phil** I am struggling to understand this sentence, rewrite Additional selection requirements are added for objects to classify as a particular particle candidate in order to reduce false positives of events that are included in a given category.

4.1.1 Variable Definitions

Many of the object definitions use variables that are derived from reconstructed quantities. They can be understood in terms of the reconstruction described in Section 3.4. Lepton isolation is quantified using the following formula.

$$I = \frac{1}{p_T^\ell} \left(\sum p_T^{\text{charged}} + \max \left[0, \sum p_T^{\text{neutral}} + \sum p_T^\gamma - p_T^{\text{PU}} \right] \right) \quad (4.1)$$

The sums are over charged hadrons originating from the primary vertex and all neutral hadrons and photons within a distance of $\Delta R < 0.4$ from the lepton if it is a muon, or $\Delta R < 0.3$ from an electron, where ΔR is a distance on the (η, ϕ) plane.

$$\Delta R = \sqrt{\Delta\eta^2 + \Delta\phi^2} \quad (4.2)$$

The term p_T^{PU} is defined as the following for muons.

$$p_T^{\text{PU}} = 0.5 \times \sum p_T^{\text{PU,charged}} \quad (4.3)$$

Electrons use a different definition.

$$p_T^{\text{PU}} = \rho \times A_{\text{eff}} \quad (4.4)$$

A_{eff} is the area of the isolation cone, and ρ is the median of the p_T density of neutral particles in that area.

Particles can also be defined as coming from the primary vertex of an event or from pileup. Vertices are defined through deterministic annealing [44], using the closest approach of tracks to the beamline [45]. The primary vertex is the vertex with the greatest sum of E_T of the charged particles originating from it. After identification of the primary vertex, charged particles are classified as originating from the primary vertex or as pileup using their extrapolated track's distance in the transverse plane, d_{xy} and distance along the beamline, d_z .

4.1.2 Isolated Muons

An isolated muon gives one of the cleanest signatures in CMS, with only perhaps the exception of an isolated photon that does not undergo pair production in the pixel tracker. Muons can also show up in jets from weakly decaying hadrons, in which case they are not isolated. Since weakly decaying b jets are central to this analysis, events with non-isolated leptons are not rejected, but the distinction is important. Loosely selected muons must meet the following requirements so that they are more likely from the decay of W and Z bosons.

- The muon must have a relatively high transverse momentum of $p_T > 5 \text{ GeV}$.
- The muon should pass through the inner tracker within $|\eta| < 2.4$.
- The muon originates from the primary vertex, satisfying both $d_{xy} < 0.5 \text{ cm}$ and $d_z < 1.0 \text{ cm}$.
- The muon must pass a loose isolation cut of $I < 0.4$.
- The muon must be a PF muon.

- The muon is either a global muon or a tracker muon.

Tightly identified muons have some additional criteria they must pass.

- They must have a higher transverse momentum at $p_T > 25 \text{ GeV}$. In events with two muons, such as caused by $Z \rightarrow \mu\mu$, only one muon only needs to satisfy this p_T requirement. **Phil**muon only needs=> muon needs Only $p_T > 15 \text{ GeV}$ is required of the second muon. **Phil**Only => A
- The muon must be a global muon, leaving tracks in both the central tracker and the muon chambers.
- There must be more than five hits in the inner tracker with at least one hit in the pixel detector.
- The fit for the global muon track must be good with $\chi^2/ndof < 10$.
- The muon must be well isolated with $I < 0.06$

These definitions are an accepted standard in the CMS collaboration as loose and tight working points, respectively. This allows the analysis to use efficiency measurements created for wider use.

4.1.3 Isolated Electrons

The kinematic variables associated with an electron are extracted from the GSF fit. Loosely selected electrons must meet the following requirements.

- They must have a transverse momentum satisfying $p_T > 7 \text{ GeV}$.
- They should be centered in the detector with $|\eta| < 2.4$.
- The distance from the primary vertex is limited, requiring $d_{xy} < 0.05 \text{ cm}$ and $d_Z < 0.2 \text{ cm}$.
- They pass a loose isolation cut of $I < 0.4$.

To optimize the electron selection, electrons are identified with the aid of an MVA [46]. Fully selected electrons pass the tight working point used by the CMS collaboration. In order to also match the samples of simulated electrons used in the training sample, the selected electrons must also pass the following cuts.

- The electron must have higher energy with $p_T > 15 \text{ GeV}$.
- The deposit of HCAL energy must be less than 9% of the ECAL energy deposit along the electron track.
- The track sum p_T component of the isolation must be less than 18% of the electron p_T .
- There is a gap in the ECAL geometry, so the electron must be outside the gap $1.4442 < |\eta| < 1.5660$. The gap is between the barrel and endcap regions of the detector. Due to detector geometry and different conditions in each region, the remaining selection criteria differs based on the electron's location.
- For electrons with $|\eta| < 1.4442$ in the detector barrel:
 - The shower shape must satisfy $\sigma_{i\eta i\eta} < 0.012$
 - Isolation in the ECAL cluster must be less than 0.4, and isolation in the HCAL must be less than 0.25.
 - The difference between super cluster and track location of the electron must be small with $\Delta\eta < 0.0095$ and $\Delta\phi < 0.065$.
- For electrons with $|\eta| > 1.5660$ in the detector endcap:
 - The shower shape must satisfy $\sigma_{i\eta i\eta} < 0.033$
 - Isolation in the ECAL cluster must be less than 0.45, and isolation in the HCAL must be less than 0.28.

4.1.4 Jets

Strong interactions cause jets of particles when quarks or gluons hadronize. Conservation of energy and momentum means that the sum of jet constituents give the kinematics of the initial parton that produced them. Jets are constructed by clustering all particle-flow candidates with the anti- k_T algorithm [47] using the jet clustering parameter $R = 0.4$. Due to factors like pileup, imperfect detector response, and differences in various jet-clustering configurations the energy of the reconstructed jets are corrected [48].

Loose jet cuts, based on the constituents, are applied to remove jets constructed from detector noise. Jets that get a significant fraction of their energy from pileup are also removed. Pre-selected electrons and muons are also often reconstructed as jets. Any jet within $\Delta R < 0.4$ from a pre-selected lepton is removed. Phil You should forward reference the selections, even though I don't like forward referencing, wait maybe you don't do that in which case you should reference these ids with a citation

To be considered for the study of decay products of the Higgs boson, jets must be within the inner tracker of the detector with $|\eta| < 2.5$. This allows pileup to be removed and for accurate vertexing of the constituents. The jets must satisfy $p_T > 25 \text{ GeV}$ for the zero and one lepton signatures. The two lepton signature from $Z(\ell\ell)H$ is cleaner, and looser jet selection criteria of $p_T > 20 \text{ GeV}$ are applied to the jets.

4.1.5 Identification of b Jets and Energy Regression

Jets containing b hadrons have a distinct signature. This includes secondary vertices displaced from the beamline, as well as non-isolated leptons from weak decays. When looking at jets in the inner tracker, all of these features are considered in a deep neural network (DNN) called Deep Combined Secondary Vertex (DeepCSV) designed to identify b jets [49]. The output of DeepCSV has three working points that are defined based on the amount of false positives that are expected in a collection of jets passing the cut. The specific values are different for each year of operation. The detector and

Table 4.1: The minimal cut value on the neural network output for each DeepCSV working point are defined for each year of Run 2 of the LHC. The working points are defined by their mis-tag rates. **Phil**mis-tag rates of what data, simulation?

| Working Point | Mistag Rate | 2016 | 2017 | 2018 |
|---------------|-------------|--------|--------|--------|
| Loose | 10% | 0.2219 | 0.1522 | 0.1241 |
| Medium | 1% | 0.6324 | 0.4941 | 0.4184 |
| Tight | 0.1% | 0.8958 | 0.8001 | 0.7527 |

collision conditions change, and a separate model is trained each year to account for that. The working points for each year are given in Table 4.1.

The non-isolated leptons within b jets are caused by flavor-changing weak decay of b hadrons. This decay mode also results in neutrinos which carry away a portion of the jet energy. In order to more accurately reconstruct the di-jet mass of a candidate Higgs decay, a Deep Neural Network (DNN) is trained in Tensorflow [50], which is designed to improve the energy measurement and resolution of all b jets for CMS [51]. Can you list the training targets, or specify the loss definitin? The following variables are used as inputs to the regression:

- the jet's p_T , η , mass and transverse mass
- the event's median energy density, commonly denoted as ρ
- information about the hardest lepton clustered into the jet, including momentum perpendicular to the jet, distance ΔR from the center of the jet, and the lepton's flavor
- the p_T , mass, and number of tracks from any secondary vertex linked to the jet, as well as the secondary vertex's distance from the collision point and associated uncertainties
- the fractions energy in the jet due to charged and neutral hadrons and electromagnetic constituents
- the highest p_T of charged hadron constituents

- the energy fraction contained in five concentric rings around the jet center binned by $\Delta R \in [0, 0.05, 0.1, 0.2, 0.3, 0.4]$
- number of PF candidates in a jet
- energy sharing computed by

$$\frac{\sqrt{\sum_i p_{T,i}^2}}{\sum_i p_{T,i}}$$

where i runs over the jet constituents

This list results in 41 input variables for the DNN. All selections depending on the p_T values of b -tagged jets are evaluated after applying the b -jet energy regression.

4.1.6 Fat Jets

When the Higgs boson is highly boosted, its decay products are closer together in the frame of the detector. At some point, the jets from the b quarks would not be clustered separately. These highly boosted events are interesting for the differential cross section measurements, and they are more enriched with the signal associated production process than events with lower p_T intermediate particles. To not loose the events, single jets are analyzed for evidence of containing two b hadrons. In order to handle the transition from resolved jets to boosted single jets, larger jets, labelled fat jets, are used. A second collection of jets is made from the same set of PF candidates that are clustered to create the jets described previously. This collection also uses the anti- k_T algorithm, but with a larger jet clustering parameter of $R = 0.8$ to capture the majority of boosted events. The requirement on the fat jet which ensures that Higgs decay products are contained in the jet when they are at a maximum opening angle is $p_T > 250 \text{ GeV}$.

Being significantly larger in area, the fat jets also contain much additional radiation from the underlying event. As a result, the mass of the constituents collected into the jet is also significantly larger than the original mass of the primary parent particle whose daughters make up most of the jet constituents. A number of grooming algorithms, which are designed to remove particles from pileup from the jets, were

considered within CMS [52]. The soft drop algorithm [53] was chosen as the standard in the experiment and is used in this analysis. The resulting groomed mass of the jet is close to its original parent particle. The soft drop algorithm has the additional benefit of forcing pileup jets to low mass, which is why a second requirement on fat jets considered for the analysis is $m_{\text{SD}} > 50 \text{ GeV}$. **Phil** I actually don't know but I would presume you guys are using PUPPI jets, since it is the default, can you cite PUPPI please

4.1.7 Missing Transverse Momentum

Missing transverse energy, which is actually the missing transverse momentum, also labelled E_T^{miss} or MET, is a vector that takes advantage of the fact that momentum transverse to the beamline is conserved. Particle flow MET is calculated by taking the negative vector sum of the transverse momentum of all particle flow candidates in the event. The resulting vector is then adjusted by taking into account the difference between the uncorrected and corrected jet energies [54]. The resulting magnitude and direction is a proxy for the transverse momentum of any neutrinos in the event. Since particle flow MET is the type of MET that is used most often in this analysis, it is referred to throughout as just MET. Large MET values can be generated by instrumental and beam effects as well. Therefore, there are additional event filters applied to events with large MET that removes events where these known instrumental and beam effects have been identified.

An additional technique to select interesting events with a large MET **Phil**with large MET is to compare the particle flow MET to other calculations of MET. This analysis also uses track MET, or track MET. **Phil** I am seeing double, please fix Track MET is calculated using just the reverse vector sum of charged particles that are detected by the tracker. In collisions without leptons, charged and neutral particles energy distributions are similar in the detector, so differences in track and particle flow MET directions can indicate instrumental failures in purely hadronic environments.

4.1.8 Soft Hadronic Activity

In signal VH events, hadronic activity outside of the $b\bar{b}$ decay of the Higgs is expected to be low. This hadronic activity is defined by considering the additional charged PF tracks coming from the primary vertex. An exclusion region is defined in an ellipse in (η, ϕ) space containing the two selected b jets with a major axis length of $\Delta R(b\bar{b}) + 1$ and a minor axis length of 1. All charged tracks outside of this ellipse that also do not correspond with the selected leptons and that satisfy $p_T > 300 \text{ MeV}$ and $d_Z < 0.2 \text{ cm}$ are clustered using the anti- k_T algorithm [47] with $R = 0.4$. The resulting collection of soft jets is used to define four variables:

- H_T^{soft} – The scalar sum of soft jets' p_T for jets with $p_T > 1 \text{ GeV}$
- N_2^{soft} – The number of soft jets with $p_T > 2 \text{ GeV}$
- N_5^{soft} – The number of soft jets with $p_T > 5 \text{ GeV}$
- N_{10}^{soft} – The number of soft jets with $p_T > 10 \text{ GeV}$

These variables are used in the training of the BDT that discriminates signal and background events. **Phil**Not your fault, but I don't think we can really simulate these variables well, its a bit of a shock that they use this

4.1.9 Kinematic Fit

In the two-lepton region, the Z boson's decay products are observed directly, and the Z boson momentum is precisely reconstructed. A kinematic fit is performed to constrain the momenta of the two leptons by requiring their combined mass to match the Z boson mass. After this constraint is applied, the transverse momentum of the di-jet system, along with identified initial state radiation jets, is balanced with the transverse momentum of the di-lepton system. The fit adjusts the momentum of each identified component of the event in a way that minimizes the chi-squared value meeting measured and theoretical physics constraints.

For electrons and muons, the corrections defined in centralized efforts by the CMS collaboration include uncertainties. The energy uncertainty used for the b -jets and




Figure 4-1: The Higgs di-jet mass in the 2-lepton signal samples is shown above. Peaks from the raw jet, the regressed jet energy, and the kinematic fit are compared.

recoil jets is the same as that used by the $HH \rightarrow b\bar{b}b\bar{b}$ analysis from CMS [55]. The mass of the Z boson is given a Gaussian uncertainty of 5 GeV, and it is assumed that the MET in the event is 0. **Phil**with what resolution? The kinematic fit minimizes the chi-squared value of these constraints, resulting in new energies for all particles in the fit. The full implementation is in the `PhysicsTools/KinFitter` package of `CMSSW_10_2_0_pre3` [56]. The fitted b -jet energies are then used for all values in the remainder of the two-lepton channel analysis. The improvement of the di-jet mass in the signal sample as a result of the kinematic fit are shown in Figure 4-1.

4.2 Simplified Template Cross Section Bins

The measurement performed in this analysis is a binned differential cross section of Higgs production using the Simplified Template Cross Section (STXS) approach [57]. The allows the result to be compared more thoroughly to Standard Model predictions while reducing the model dependence of the measurements. For this measurement, the vector boson produced as well as the p_T of the vector boson separates data points into different bins. The clean signal of the $Z \rightarrow \ell\ell$ decay channel allows for more bins to be measured for ZH processes. In addition, the middle p_T bin for Z boson

production is split by multiplicity of additional jets. There are eight bins overall:

- $WH, 150 \text{ GeV} < p_T(V) \leq 250 \text{ GeV}$
- $WH, 250 \text{ GeV} < p_T(V) \leq 400 \text{ GeV}$
- $WH, 400 \text{ GeV} < p_T(V)$
- $ZH, 75 \text{ GeV} < p_T(V) \leq 150 \text{ GeV}$
- $ZH, 150 \text{ GeV} < p_T(V) \leq 250 \text{ GeV}, n_{\text{jet}} = 0$
- $ZH, 150 \text{ GeV} < p_T(V) \leq 250 \text{ GeV}, n_{\text{jet}} \geq 1$
- $ZH, 250 \text{ GeV} < p_T(V) \leq 400 \text{ GeV}$
- $ZH, 400 \text{ GeV} < p_T(V)$

All of the selections in the following sections are also divided into the appropriate set of STXS bins for the generation of datacards and fits. Since the fat jets are most helpful in events where intermediate particles are highly boosted, they are only considered in selections for the bins where $p_T(V) > 250 \text{ GeV}$.

4.3 Treatment of Background Yields

In order to effectively measure Higgs production, we need to be able to accurately estimate other events that end up in our selection. To start, the most significant contaminant physics processes with a final state identical to the $VHb\bar{b}$ process are identified. Then, selections that are enriched with these processes, called control regions, are created, which allows more direct comparison between simulation of these processes and collected data. When fitting for the signal strength, the background process yields are simultaneously fit from these control regions.




Figure 4-2: Above is the Feynman diagram matching the two lepton final state coming from Drell-Yan and jets.

4.3.1 Identification of Background Processes

The final state for the two lepton decay in Figure 2-8 contains two oppositely-charged, same flavor leptons and two b -tagged jets. It can also be achieved by a Drell-Yan process radiating b jets or a $t\bar{t}$ event where both W bosons from the top decays decay leptonically. Feynman diagrams in Figure 4-2 and Figure 4-3 show how the two respective processes can result in the same final state as the signal process. The Drell-Yan process can also radiate jets initiated by lighter flavor quarks that are mistakenly identified as b -jets, and those make up a significant portion of the backgrounds as well. Less significant, but still important backgrounds include processes like di-boson production, QCD jets, and single top processes.

The backgrounds for the one- and zero-lepton signal decay channels are caused by similar processes. For the one-lepton decays of WH , the Drell-Yan background in Figure 4-2 is replaced with a flavor changing current of $W + \text{jets}$. The $t\bar{t}$ background would instead be caused by either a hadronic decay of one of the W bosons in Figure 4-3 or by one of the pictured leptons travelling out of the detector without being observed. For the zero-lepton channel, the Drell Yan process is instead replaced with $Z \rightarrow \nu\bar{\nu}$. The $t\bar{t}$ process still needs high MET in order to appear to contain a hard neutrino presence, so it is most often caused when a single W decays leptonically with the lepton falling outside of the detector acceptance. For both of these channels,



Figure 4-3: Above is the Feynman diagram matching the two lepton final state coming from fully leptonic $t\bar{t}$ decay. In events with little energy carried away by the neutrinos, this can appear to be the same as the two-lepton signal process.

di-boson, QCD, and single top backgrounds can also contribute, but not significantly enough to require a tight constraint on their yields.

4.3.2 Goals for Control Regions

To accurately estimate the contribution of each of these backgrounds, control regions are used. These are selections that are in similar phase spaces as the signal selection, but are instead enriched with background events. By comparing the prediction from Monte Carlo to data, scaling corrections to the simulation are made for the relevant phase space and extrapolated to the signal region.

For each channel, three control regions are defined. There are two control regions for the appropriate vector boson radiating jets. The region is enriched with light flavor jets, labelled $udsg$ for up, down, and strange quarks, and gluons, by requiring a di-jet system that fails the b -tagging working points. The heavy flavor $V + \text{jets}$ control regions are differentiated from the signal selection primarily by requiring the di-jet mass to be different than Higgs boson mass. The $t\bar{t}$ process tends to radiate more jets than the signal process, but the lack of a di-lepton resonance is taken advantage of in the two-lepton regions. Specific cuts for each signal region and the control regions are given in the next section.

4.4 Resolved Analysis Selection

First, the selection for events with two b -tagged jets, also known as resolved events, will be given. The next section will explain the selections relying on a single fat jet. With objects defined, the selections differ mostly in counting the number of charged leptons present in the event. However, other adjustments are made per channel to optimize the presence of signal events. Therefore, the first channels described will have the most thorough selection description, with later channel sections noting many similarities and differences. For each channel, multiple control region selections are also used in order to more accurately estimate the contribution of each physics process to the events in each phase space, and these will be described after each channel's signal region. The most significant contributions to background contamination of the signal region are from $Z/W + \text{jets}$ (depending on the number of leptons in the selection) and $t\bar{t}$ processes. The vector boson plus jets regions are separated into heavy-flavor and light-flavor jets to control the relative contributions independently since they are not well-known. For each channel, there are therefore three control regions in addition to the signal region.

A summary of cuts for each region in each channel is given in Table 4.2. A few channel- or region-specific cuts are left out, but are described in the appropriate sections. The common variables that are cut on are the following:

- $p_T(V)$ – The transverse momentum of the reconstructed vector boson
- $p_{T,j}$ – The minimum transverse momentum of both jets, which may be an asymmetric cut
- $\max b$ – The working point that the jet with the higher b -tag value must satisfy
- $\min b$ – The working point that the jet with the lower b -tag value must satisfy
- $p_{T,jj}$ – The summed transverse momentum of the di-jet system
- m_{jj} – The mass of the di-jet system
- N_{aj} – The number of additional jets outside of the selected di-jet system

Table 4.2: Below is a summary of common cuts for all regions in the resolved channels. See the text for each channel for an explanation of variables. All energy equivalent values are in GeV. A “!” in front of a working point label means the jet is required to fail the relevant b -tagging criteria. **Phil**minor point, cuts is jargon. Bigger point, I really think you need a diagram for these control regions its really hard to flow otherwise

| 0-lepton channel | | | | | | | | |
|------------------|----------|-----------|---------|---------|------------|-----------------|-----------------|---------------------|
| Region | $p_T(V)$ | $p_{T,j}$ | max b | min b | $p_{T,jj}$ | m_{jj} | N_{aj} | $\Delta\phi(jj, V)$ |
| Signal | 170 | 60, 35 | med. | loose | 120 | $>90, <150$ | ≤ 1 | > 2.0 |
| $Z + b$ | 170 | 60, 35 | med. | loose | 120 | <90 or >150 | ≤ 1 | > 2.0 |
| $Z + uds g$ | 170 | 60, 35 | !med. | loose | 120 | $>50, <500$ | ≤ 1 | > 2.0 |
| $t\bar{t}$ | 170 | 60, 35 | med. | loose | 120 | $>50, <500$ | ≥ 2 | – |
| 1-lepton channel | | | | | | | | |
| Region | $p_T(V)$ | $p_{T,j}$ | max b | min b | $p_{T,jj}$ | m_{jj} | N_{aj} | $\Delta\phi(jj, V)$ |
| Signal | 150 | 25 | med. | loose | 100 | $>90, <150$ | ≤ 1 | > 2.5 |
| $W + b$ | 150 | 25 | med. | loose | 100 | <90 or >150 | ≤ 1 | > 2.5 |
| $W + uds g$ | 150 | 25 | !med. | loose | 100 | $>50, <250$ | – | > 2.5 |
| $t\bar{t}$ | 150 | 25 | med. | loose | 100 | $>50, <250$ | ≥ 2 | – |
| 2-lepton channel | | | | | | | | |
| Region | $p_T(V)$ | $p_{T,j}$ | max b | min b | $p_{T,jj}$ | m_{jj} | N_{aj} | $\Delta\phi(jj, V)$ |
| Signal | 50 | 20 | med. | loose | 50 | $>90, <150$ | – | > 2.5 |
| $Z + b$ | 50 | 20 | med. | loose | 50 | <90 or >150 | – | > 2.5 |
| $Z + uds g$ | 50 | 20 | !loose | !loose | 50 | $>50, <250$ | – | > 2.5 |
| $t\bar{t}$ | 50 | 20 | tight | loose | 50 | $>50, <250$ | – | – |

Table 4.3: Below are the trigger paths used for the 0 lepton selections for all three years of Run 2 of the LHC.

| Year | Trigger path(s) |
|------|---|
| 2016 | HLT_PFMET110_PFMHT110_IDTight HLT_PFMET120_PFMHT120_IDTight HLT_PFMET170_NoiseCleaned HLT_PFMET170_BeamHaloCleaned HLT_PFMET170_HBHECleaned |
| 2017 | HLT_PFMET120_PFMHT120_IDTight HLT_PFMET120_PFMHT120_IDTight_PFHT60 |
| 2018 | HLT_PFMET120_PFMHT120_IDTight |

- $\Delta\phi(jj, V)$ – The distance in angle ϕ between the di-jet system and the recoiling reconstructed vector boson

4.4.1 0 Leptons

In the 0-lepton channel, the transverse momentum carried away by the neutrinos in the Z boson decay results in a large amount of MET. CMS has triggers that identify events with large values of MET. Slightly different triggers are used for each year of Run 2 of the LHC, and they are listed in Table 4.3. The MET for the event must be larger than 170 GeV. The trigger efficiency does not quite plateau at that point, as can be seen in Figure 4-4. However, the extracted efficiency scale factor for MC matches for both $t\bar{t}$ and $W + \text{jets}$ simulation at and above this value. During the 2018 run, a number of HCAL endcap modules were taken offline due to power supply problems. These modules were all located in the region $-1.57 < \phi < -0.87$, so an excess number of high MET events with ϕ_{MET} in that region were recorded in later 2018 runs when jets would have been registered in the deactivated detector elements. The resulting peak can be seen in Figure 4-5. To handle this, all events with $-1.86 < \phi_{\text{MET}} < -0.7$ that occurred during and after run 319077, when the faulty detector elements were shut off.

In addition to the neutrino decay of the Z boson, the $b\bar{b}$ decay of the Higgs also needs to be selected and backgrounds need to be removed. Many of the following cuts are similar for all of the channels. For the Higgs decay, two jets are selected. One jet




Figure 4-4: The efficiency for the MET triggers are shown for all three years in data.




Figure 4-5: Above compares the MET ϕ distributions before and after shutting off problematic HCAL modules during for run 319077. The excess of events in the region $-1.86 < \phi_{\text{MET}} < -0.7$ is caused by mis-measuring the momenta of forward jets in that region.

is the one with the highest b -tag score out of jets with $p_T > 60$ GeV. The other jet is the highest b -tag score out of jets remaining with $p_T > 35$ GeV. This distinction is important, because these two jets may not have the highest b -tag score of the event if the two highest b -tag scored jets both have $p_T < 60$ GeV, for example. However, we are not interested in such relatively low p_T systems since the di-jet system should be reasonably balanced against the MET requirement. To additionally enforce this balance, the di-jet system must satisfy $p_T > 120$ GeV. Also the di-jet system is selected to be back-to-back with the MET by requiring $\Delta\phi(\text{MET}, jj) > 2.0$. The di-jet mass is required to be less than 500 GeV to remove high energy combinatoric backgrounds. To increase purity of $Z\nu\nu$ processes, events are not considered if there are any isolated leptons with $|\eta| < 2.5$ and $p_T > 15$ GeV. Finally, to reduce QCD background contributions, all jets in the event with $p_T > 30$ GeV must be a minimum distance from the event MET satisfying $\Delta\phi(\text{MET}, j) > 0.5$.

In the signal region, it is further required that the selected b -jets to be of high quality, the di-jet system has a mass close to the Higgs, and low additional hadronic activity. At least one of the b -jets must pass the medium working point, and the other b -jet must pass the loose working point. The value of the di-jet mass must satisfy $90 \text{ GeV} < m_{jj} < 150 \text{ GeV}$. At most one jet with $p_T > 30$ GeV is allowed in addition to the selected b -jets. Since the event has no high energy leptons, the particle flow MET and track MET must have good agreement with $\Delta\phi(\text{MET}, \text{trackMET}) < 0.5$.

As mentioned earlier, three processes make up the majority of the backgrounds in the signal selection, and control regions must be defined for them. Two of them when a $Z \rightarrow \nu\bar{\nu}$ occurs recoiling off of jets. In one process, the recoiling jets are b -jets, and in the other, the jets are light jets. Since the fraction of actual $Z \rightarrow \nu\bar{\nu}$ events containing b -jets is not well known, they are scaled separately. The third set of background events that needs to be separately measured in this phase space are the $t\bar{t}$ semi-leptonic decays where the lepton falls outside of the detector acceptance. This type of background also results in a final state with high MET and b -jets.

Of the three processes, the $Z + \text{heavy flavor jets}$ selection is the most similar to the signal selection. The only difference is the di-jet mass. Events with a mass

Table 4.4: Below are the trigger paths used for the 1 lepton selections for all three years of Run 2 of the LHC. Different triggers are used for the electron and muon channel, and are labelled separately.

| Year | Lepton | Trigger path(s) |
|------|--------|----------------------------------|
| 2016 | e | HLT_Ele27_WPTight_Gsf |
| 2016 | μ | HLT_IsoMu24 HLT_IsoTkMu24 |
| 2017 | e | HLT_Ele32_WPTight_Gsf_L1DoubleEG |
| 2017 | μ | HLT_IsoMu27 |
| 2018 | e | HLT_Ele32_WPTight_Gsf |
| 2018 | μ | HLT_IsoMu24 |

between $50 \text{ GeV} < m_{jj} < 500 \text{ GeV}$ but not between $90 \text{ GeV} < m_{jj} < 150 \text{ GeV}$, which is the signal region window, are selected to quantify the $Z + \text{heavy flavor}$. The $Z + \text{light flavor}$ selection includes the entire mass range, without a veto for the Higgs mass. It is instead enriched with light jets by requiring that the selected b -jet with a higher b -tag score fails the medium working point. The selection for $t\bar{t}$ events is different from the signal selection mostly because more hadronic activity is expected. It uses the same full di-jet mass window as the $Z + \text{light flavor}$ region, but requires at least two additional jets with $p_T > 30 \text{ GeV}$ instead of zero or one. Also, the $\Delta\phi(\text{MET}, \text{trackMET})$ requirement is dropped.

4.4.2 1 Lepton

In the 1-lepton channel, a single fully-selected isolated lepton as defined in Section 4.1.2 or Section 4.1.3 is required. CMS has triggers for isolated leptons, and the ones used in the analysis are listed in Table 4.4. That lepton must point in a similar direction of the MET, satisfying $\Delta\phi(\ell, \text{MET}) < 2.0$. The p_T of the reconstructed W boson, consisting of the vector sum of MET and the lepton, must satisfy $p_T > 150 \text{ GeV}$. If there are any additional leptons, the event is not used. The presence of an isolated lepton provides a much cleaner signal than in the zero lepton channel. Therefore, the kinematic cuts on the selected b -jets can be looser. The b -jets only need to satisfy $p_T > 25 \text{ GeV}$, and the di-jet system only needs $p_T > 100 \text{ GeV}$. Any events with $m_{jj} \geq 250 \text{ GeV}$ are not considered for any regions. The b -tagging

Table 4.5: Below are the trigger paths used for the 2 lepton selections for all three years of Run 2 of the LHC. Different triggers are used for the electron and muon channel, and are labelled separately.

| Year | Leptons | Trigger path(s) |
|------|--------------|--|
| 2016 | e^+e^- | HLT_Ele23_Ele12_CaloIdL_TrackIdL_IsoVL_DZ |
| 2016 | $\mu^+\mu^-$ | HLT_Mu17_TrkIsoVVL_Mu8_TrkIsoVVL HLT_Mu17_TrkIsoVVL_TkMu8_TrkIsoVVL HLT_Mu17_TrkIsoVVL_Mu8_TrkIsoVVL_DZ HLT_Mu17_TrkIsoVVL_TkMu8_TrkIsoVVL_DZ |
| 2017 | e^+e^- | HLT_Ele23_Ele12_CaloIdL_TrackIdL_IsoVL_DZ |
| 2017 | $\mu^+\mu^-$ | HLT_Mu17_TrkIsoVVL_Mu8_TrkIsoVVL_DZ_Mass3p8 HLT_Mu17_TrkIsoVVL_TkMu8_TrkIsoVVL_DZ_Mass8 |
| 2018 | e^+e^- | HLT_Ele23_Ele12_CaloIdL_TrackIdL_IsoVL_DZ |
| 2018 | $\mu^+\mu^-$ | HLT_Mu17_TrkIsoVVL_Mu8_TrkIsoVVL_DZ_Mass3p8 |

requirement is the same as the one in the 0-lepton channel. There is a slightly tighter requirement on the di-jet direction of $\Delta\phi(\text{MET}, jj) > 2.5$.

In the signal region, the additional selection criteria are again similar to the 0-lepton channel. The di-jet mass window is the same, as is the requirement of at most one additional jet. The only other difference from the 0-lepton selection aside from the adjusted criteria listed above is that there is no dependence on the track MET direction. The change to create the W + heavy flavor jets control region is the exact same as the Z + heavy flavor region in the 0-lepton channel. The mass window for the Higgs is vetoed. The cut for the W + light flavor control region is also the same in terms of b -tagging, but the additional jet requirement is also removed. The $t\bar{t}$ control region is also the same in that the only changes from the signal region are a relaxed mass window, the requirement of at least two additional jets, and no requirement on the di-jet direction relative to MET.

4.4.3 2 Leptons

For the 2-lepton channel, two oppositely charged leptons with the same flavor are required. The triggers used that are designed to match this requirement are given in Table 4.5. They must have a mass satisfying $75 < m_{\ell\ell} < 105 \text{ GeV}$. This process is clean enough to relax the kinematic cuts on the selected b -jets even further than

the relaxed selection criteria of the 1-lepton channel. The selected b -jets only need to have a $p_T > 20$ GeV, and the di-jet system only needs $p_T > 50$ GeV. There are no requirements on the number of additional jets, outside of categorization for STXS bins. The di-jet system still needs to satisfy the tighter cut of $\Delta\phi(jj, V) > 2.5$.

The signal region uses the same b -tag and mass value requirement as the other two channels. For the $Z + \text{heavy flavor}$ control region, the di-lepton mass is narrowed to $85 < m_{\ell\ell} < 97$ GeV in order to remove $t\bar{t}$ events, and the usual Higgs mass veto is applied. The MET is also required to be low with $\text{MET} < 60$ GeV for the $Z + \text{heavy}$ region, but no other. For the $Z + \text{light flavor}$ region, purity is achieved by requiring that both selected jets fail the loose working point for b -tagging. The $t\bar{t}$ region is then selected by requiring one selected jet to pass the tight working point. The di-lepton mass value also must be either $10 < m_{\ell\ell} < 75$ GeV or $m_{\ell\ell} > 120$ GeV.

4.5 Boosted Analysis Selection

When the Higgs has an adequately high p_T , the jet clustering algorithms will often find both daughter particles as being part of a single jet. The boosted analysis only targets the STXS bins with $p_T(V) > 250$ GeV. The selection differs primarily in the fact that a single fat jet which passes a double b -tag cut [49] is used to reconstruct the potential Higgs instead of two b -tagged jets. The double b -tagger used in this analysis is DeepAK8, a DNN as opposed to a BDT tagger. The output is decorrelated with mass, allowing for the jet mass to be used to generate control regions, as is done in the resolved analysis. [Phil](#) can you reference the decorrelation strategy

Additional b jets outside of the fat jet are also counted to define selections. These come from the regular jet collections of Section 4.1.4. In order to count as an additional b -jet, the jet must pass the DeepCSV medium working point, have a $p_T > 25$ GeV, and be outside of the selected fat jet with $\Delta R(j, fj) > 0.8$.

4.5.1 0 Leptons

The boosted selection regions are designed to study the same processes as the resolved selection regions. The 0-lepton channel is enriched with signal events where the Z boson decays to neutrinos, so the 0-lepton channel has high MET. A requirement of $\text{MET} > 250 \text{ GeV}$ is applied, which is consistent with the p_T requirement of the fat jet. As for the resolved analysis, the presence of any leptons leads to the event not being considered for the 0-lepton channel. To remove QCD background for all regions, the same cut from the resolved analysis of $\Delta\phi(\text{MET}, j) > 0.5$ for all jets with $p_T > 30 \text{ GeV}$ is used.

In the signal region, jets must have a score of 0.8 or higher in the `bbVsLight` output node of the `DeepAK8` tagger. They must also have a soft drop mass in the range of $90 < m_{\text{SD}} < 150 \text{ GeV}$. No additional jets outside of the fat jet are allowed in the event. The control regions are the same as for the resolved analysis: $Z + \text{heavy flavor}$, $Z + \text{light flavor}$, and $t\bar{t}$. For the $Z + \text{heavy flavor}$ control region, the mass cut is changed to instead veto the Higgs mass window. For the $Z + \text{light flavor}$, there is no mass requirement outside of the $m_{\text{SD}} > 50 \text{ GeV}$ required for all fat jets. Orthogonality is enforced by requiring the `bbVsLight` score to be less than 0.8. For $t\bar{t}$, the lack of a mass requirement is also present, but there must be at least one b jet outside of the fat jet.

4.5.2 1 Lepton

For the single lepton channel, exactly one selected lepton must be present. It must also point in the same direction as the MET with $\Delta\phi(\text{MET}, \ell) < 2.0$. Otherwise, the selection criteria for the different 1-lepton regions are the exact same as for the boosted 0-lepton regions.

4.5.3 2 Leptons

For the two lepton channel, two oppositely charged, same flavor leptons must be present, as described in Section 4.4.3. These leptons must also have a mass near the

Z boson mass for the signal region and the two $Z + \text{jets}$ regions of $87 < m_{\ell\ell} < 95 \text{ GeV}$. The selection for the signal and control regions are otherwise similar to the selections for the 0- and 1-lepton channels in the boosted analysis. The only difference is that instead of requiring a b jet outside of the fat jet for the $t\bar{t}$ control region, a mass veto of the di-lepton mass is applied, just as was done for the resolved analysis.

4.6 Overlap in Resolved and Boosted Selections

An important note is that each signal and control region described in this chapter must be orthogonal to all other regions. To prevent any statistical bias in the analysis, both simulated and measured events must not be double counted. It is easy to make orthogonal selections within resolved or boosted categories by counting the number of leptons, comparing their flavor, and reversing other selections to go from signal to control regions. However, when making selections for the resolved and boosted channels, enforcing orthogonality is not as straightforward. The same PF candidates are reused to define two different kinds of jet collections, making it harder to reverse a single selection cut to define a different category. The selections described so far result in some events simultaneously passing both boosted and resolved selections. To maximize the expected sensitivity of this analysis, most events that are in both resolved and boosted categories are assigned to the resolved category and omitted from the boosted. The only exception to this rule is when the event is assigned to one of the resolved control region and a boosted signal region. In that case, the event is assigned to its boosted category. **Phil**It shocks me that resolved is better than boosted, can you add a few sentence explanation why?

PhilOne note is that the word cut is jargon and should not be in your thesis. Overall I thought the selections part is written well, very concise and precise, it is easy to follow. However, I think you should add a diagram this would be good for your talk too.

Chapter 5

Corrections and Uncertainties

Despite efforts to simulate LHC collisions accurately, a number of differences in the distributions predicted by MC and present in data arise. There are many reasons for this. The theory calculations used for simulation are not accurate since only leading order or next-to-leading order perturbations are used by the generators. After the generator step, the detector must be simulated, which is difficult due to the many composite parts, both active detector material and inactive electronics, that must be modeled in `GEANT4`. This is made more difficult since the detector degrades in its high radiation environment. The relation of generating particles and detector geometry also depends on the beam conditions, which are also not predicted exactly. The most accurate measurements of the beam profile are destructive to the beam, so measurements cannot be made during the run. Predictions from simulations are also made harder because there are so many events recorded at the LHC, so tails will be generated in obscure areas of phase space. These tails are difficult to model correctly, yet often contain the most interesting events. Corrections are made to the simulation by re-weighting based on detector conditions and efficiencies of particle identification, and by scaling the predicted energies based on particle type.

5.1 Efficiency Scale Factors

The identification of particles based on working points happen at different rates between simulation and collected data. These efficiencies are measured separately in data and Monte Carlo. The difference depends on a scale factor, which is measured as a function of particle energy and location within the detector. Inclusive selections are used for the measurement, which is then transferred to the events in the analysis. Since this analysis relies on counting leptons for its categorization, selection efficiency for both muons and electrons must be measured. A scale factor for the MET trigger efficiency is also derived for the 0-lepton categories.

5.1.1 Muons

To remove fake muons from events, muons are selected in three ways. Muon identification cuts are applied, isolation cuts are applied, and certain triggers are required. These three things each behave differently in MC and data. For each of these, a separate efficiency is derived in MC and data, and then a total scale factor is derived via the following formula.

$$\epsilon^\mu = \epsilon_{\text{ID}}^\mu \times \epsilon_{\text{ISO|ID}}^\mu \times \epsilon_{\text{Trig|ISO}}^\mu \quad (5.1)$$

The scale factor for muon ID, which is tight for positively counting muons, and loose for counting muons to veto, is ϵ_{ID}^μ . Passing the isolation cut, given the ID is scaled by $\epsilon_{\text{ISO|ID}}^\mu$, and the scale factor for the trigger, after passing the isolation cut is $\epsilon_{\text{Trig|ISO}}^\mu$.

These are each separately measured via the tag and probe method. A single tag muon is selected in events with a single muon trigger. For events with a second, oppositely-charged muon that reconstructs the Z boson resonance with the first, the second muon, called a probe, is checked for identification efficiency. Since the probe muon does not rely on a categorization for the selection, only on the successful reconstruction of a Z boson, the collection of probe muons make up an unbiased sample for the efficiency measurement. A Breit-Wigner convoluted with a Gaussian and a falling




Figure 5-1: Example tag and probe fits are shown above. The left plot shows the dilepton mass fit for all tagged events. The tagging procedure yields little background, which allows for a precise measurement. The middle plot shows the events where the probe passed muon identification, and the right plot show the events when the probe failed.

combinatoric background is fit to the peak to estimate the contribution of the Z boson resonance in both the passing and failing probes. A scale factor is then applied to MC to match the data efficiency. Figure 5-1 shows one of these measurements of efficiency in Data. Each of the efficiency measurements and scale factors are binned in muon p_T and η . The p_T bin boundaries are $[20, 25, 30, 40, 50, 60, \infty)$ in GeV. Bins in $|\eta|$ are delimited at $[0, 0.9, 1.2, 2.1, 2.4]$.

The uncertainties from the scale factors are also combined in the final analysis. The identification efficiencies in data are slightly lower than in Monte Carlo. Loosely identified muons require an average scale factor of 0.998 ± 0.002 applied to simulated events, and tightly identified muons require an average scale factor of 0.98 ± 0.005 [58]. In general, the scale factors are half a percent or one percent lower, respectively, at values of $|\eta| > 2.0$, and flat across p_T bins.

5.1.2 Electrons

The electron scale factors are measured in a similar manner as the muon scale factors, using the tag and probe method. The relative difficulty in reconstructing electrons, which are contained in the ECAL, compared to muons leaving tracks through the muon chambers means that a reconstruction scale factor, ϵ_{RECO}^e , is also factored into the full scale factor.

$$\epsilon^e = \epsilon_{\text{RECO}}^e \times \epsilon_{\text{ISO+ID|RECO}}^e \times \epsilon_{\text{Trigger|ISO+ID+RECO}}^e \quad (5.2)$$

The rest of the efficiency scale factors follow the same definition patterns given for the muons. The binning for the electron scale factors is also the same as the binning for the muon scale factors. The scale factor is around 1.0 ± 0.1 for most of the bins. **Phil** Is it really 0.1 per bin, this seems high?, I would have expected an order of magnitude lower. Integrated over all bins, I recall we had it under 1 percent.

5.1.3 MET Trigger Scale Factors

The MET trigger efficiencies for each year are measured in events with a single electron and large MET. This allows tagging events with the single electron triggers, and selecting mostly events from W +jets and $t\bar{t}$ events. The events are additionally required to be similar to the events of interest in the analysis. There must be two jets with $p_T > 20$ GeV and $|\eta| < 2.5$, and the events must pass the MET filters. The event must also not be biased by the L1 MET triggers activated by the calorimeters, which is achieved by requiring that the $\Delta\phi$ between the electron and MET is less than 2.5. **Phil** I am confused by this cut, do mean MET triggers from the electron mismeasurement, or at least influenced by the electron? The nominal values for the scale factors are derived by measuring the MET trigger efficiency in data and scaling the trigger efficiency in the $t\bar{t}$ Monte Carlo sample to match. Uncertainties are derived by repeating the measurement using a W +jets sample. The scale factors for the nominal measurement and the W +jets sample are shown in Figure 5-2.

5.2 Energy Corrections of b -Jets

This analysis does depend in particular on b -jet energy predictions, which do not have recommended corrections from the CMS collaboration, so a bespoke measurement is done for this analysis. Even when distributions of individual variables agree between MC and data, correlations are often different. These correlations are also important in the evaluation of a DNN. The DNN used to estimate the energy of b -jets therefore has differing performance in MC and data. In particular, it is better at estimating the true energy of a b -jet in MC. The energies evaluated in MC must be smeared in order




Figure 5-2: **Phillabel** y-axis, say MET Trigger scale factor too, I was confused. Also mention how the uncertainty is computed given the two lines. The MET scale factors for two varying simulations are given. The $t\bar{t}$ sample is used for the nominal scale factor, and the W +jets is used to estimate systematic uncertainties. The top left plot shows the measurements for 2016, the top right plot shows 2017, and the bottom plot shows the MET scale factors for 2018. Each scale factor is a function of the minimum MET or missing hadronic energy. Remember the MET cut for the 0-lepton channel is at 170 GeV, so the different behavior in 2018 below 150 GeV does not affect the analysis. The kink in the 2016 W +jets line is caused by the merging of two fits, but since this is a variation to assess uncertainties, it is not necessary to fix.

to accurately simulate the resolution of jets in data after they have been modified by the DNN regression.

One way to measure jet energy resolution is to consider an event where a jet is recoiling off of a Z boson that decays into leptons. In principle, the Z boson's transverse momentum is balanced with the jet's transverse momentum. Measurements of lepton energies in the CMS detector is relatively precise, so the ratio of the reconstructed jet's p_T to the Z boson's p_T allows measurement of the jet energy resolution. Ideally, this measurement would be done with an collision resulting in one Z boson decay, and one jet. However, this is an infrequent occurrence. Instead, events with two jets are selected, with one jet having relatively low p_T . A fit is performed to estimate resolution characteristics where the second jet would have $p_T = 0$ GeV.

These events are selected using the following requirements:

- Exactly two muons or two electrons must pass the selection criteria for the di-leptons channels described in Section 4.4.3.
- The two selected leptons must be oppositely charged.
- The di-lepton kinematics must satisfy
 $p_{T,\ell\ell} > 100$ GeV and $71 \text{ GeV} < m_{\ell\ell} < 111$ GeV.
- Exactly two jets must pass the pre-selection described in Section 4.4.3.
- The leading jet must also satisfy $\Delta\phi(j, \ell\ell) > 2.8$
- The ratio between the sub-leading jet p_T and the di-lepton p_T must be less than 0.3.
- The leading jet must pass the tight working point for the b -tagger, as defined for each year in Table 4.1.

The selected events are divided into four bins of $\alpha = p_{T,j2}/p_{T,\ell\ell}$ with bounds (0, 0.155, 0.185, 0.23, 0.3). The distribution of α is shown in Figure ??, along with the bin boundaries. These bins were selected to give approximately the same number




Figure 5-3: The distribution for α is shown above, along with the binning boundaries.

of data events in each bin. The jet response ($p_{T,j1}/p_{T,\ell\ell+j2}$) is plotted in each bin, with uncertainties from renormalization and refactorization scale weights and parton shower weights. These histograms of jet response are shown in Fig. 5-4. From each plot, the mean (μ) and the standard deviation (σ) are extracted. The relative resolution (σ/μ) is fit as a function of α .

$$f(\alpha) = (m \times \alpha) \oplus b \times (1 + c_k \times \alpha) \quad (5.3)$$

Phil Is it not the magnitude of the right formula, since there is a direct sum

The c_k term is fixed by a linear fit to the MC's intrinsic jet resolution ($p_{T,reco}/p_{T,gen}$) over α as $c_k = m_0/q_0$. The fit results are shown in Fig. 5-5. The resolution is corrected by scaling the difference between $p_{T,reco}$ and $p_{T,gen}$ by b_{data}/b_{MC} . This causes the post-smearing fits to agree at $\alpha = 0$. Uncertainties are extracted from the fit uncertainties of b for data and MC. The resulting smearing values are in Table 5.1. The




Figure 5-4: The histograms of response for each event are shown above. The top row shows 2016, the middle shows 2017, and the bottom row shows 2018 histograms.




Figure 5-5: The fits to Data, MC, and intrinsic resolutions are shown. From left to right are the fits for 2016, 2017, and 2018.

Table 5.1: The extracted corrections needed for each year of data as a percent of the jet's p_T .

| Year | Scaling | Resolution Difference |
|------|-------------------|-----------------------|
| 2016 | 0.998 ± 0.019 | 0.017 ± 0.060 |
| 2017 | 1.020 ± 0.023 | 0.088 ± 0.071 |
| 2018 | 0.985 ± 0.019 | 0.080 ± 0.073 |

difference in smearing values for 2016 can be explained by a few characteristics of the simulation and data. First, the 2016 data has been more thoroughly studied and can generally be expected to have more robust calibration. The detector behavior also degrades over time as it accumulates radiation damage. Though this is accounted for in calibrating, it does make more precise simulations of detector response less accurate. Finally, the simulation for 2016 uses a different PYTHIA tune.

5.3 Theoretical Corrections

There are known inaccuracies in the simulations used in this analysis that can be understood at the theoretical level. However, less accurate calculations are run because they are simpler and faster to compute the results for. A trade-off must be made between the accuracy of the simulation and being able to generate enough simulated events to fill the phase space relevant for all analyses CMS collaborators are undertaking.

5.3.1 LO to NLO Reweighting

For each V +jets process, $Z \rightarrow \nu\nu$, $W \rightarrow \ell\nu$, and $Z \rightarrow \ell\ell$, there are three distinct processes that are simulated separately.

- b -quarks are generated in the matrix element
- b -quarks are not in the matrix element, but are produced in the parton shower
- No b -quarks are present in the event

For the first two, VBJets and VJetsBGenFilter datasets are used respectively, as listed in Appendix B, but they are only generated for $p_T(V) > 100 \text{ GeV}$. This small phase space and low relative cross section for generating b quarks means that it is feasible to simulate these processes to Next to Leading Order (NLO) diagrams. However, expected events with no b -quarks and events with $p_T(V) < 100 \text{ GeV}$ are more numerous, so more simulated events must be generated. To speed up calculations, Leading Order (LO) samples are used for these processes.

One common way to make LO samples more accurate is to generate NLO samples and reweight the LO sample to match generator-level kinematics. This can be done with an inclusive selection so that the statistical limitations of the available NLO samples are not significant. While the samples in this analysis were generated using just MadGraph5 [35], the following inclusive samples using aMC@NLO [59] are used to reweight the LO simulation as a function of $p_T(V)$. PhilI understand NLO/LO reweighting for inclusive production, and I think what you are doing is fine, but for $z+b$ production, I am confused what is done. Are you reweighting? If so are you computing a k -factor with an NLO b -sample, what is your choice of b -generation, do you use 5-flavor pdfs, this should be specified.

- Z1JetsToNuNu_M-50_LHEZpT_50-150_TuneCP5_13TeV-amcnloFXFX-pythia8
- Z1JetsToNuNu_M-50_LHEZpT_150-250_TuneCP5_13TeV-amcnloFXFX-pythia8
- Z1JetsToNuNu_M-50_LHEZpT_250-400_TuneCP5_13TeV-amcnloFXFX-pythia8

- Z2JetsToNuNu_M-50_LHEZpT_50-150_TuneCP5_13TeV-amcnloFFFX-pythia8
- Z2JetsToNuNu_M-50_LHEZpT_150-250_TuneCP5_13TeV-amcnloFFFX-pythia8
- Z2JetsToNuNu_M-50_LHEZpT_250-400_TuneCP5_13TeV-amcnloFFFX-pythia8
- WJetsToLNu_0J_TuneCP5_13TeV-amcatnloFFFX-pythia8
- WJetsToLNu_1J_TuneCP5_13TeV-amcatnloFFFX-pythia8
- WJetsToLNu_2J_TuneCP5_13TeV-amcatnloFFFX-pythia8
- DYJetsToLL_M-50_TuneCP5_13TeV-amcatnloFFFX-pythia8

The change to the $p_T(V)$ spectra and the resulting improved agreement between data and MC are shown in Figure 5-6.

5.3.2 Electroweak and QCD Corrections to Background

Though the LO to NLO reweighting does improve the agreement between MC and data, there is also an additional small correction. Higher order electroweak and QCD corrections predict a slightly softer $p_T(V)$ spectrum [60]. The reweighting function is shown in Figure 5-7.

5.3.3 Corrections to Signal Process

The samples produced for signal are NLO. For an accurate measurement of the VH cross section, the simulation is scaled to Next to Next to Leading Order (NNLO) QCD effects, where two gluons are added to the Feynman diagrams. An example NNLO VH process is shown in Figure 5-8. Electroweak corrections can still be factored out, and total VH production cross sections are given as the following [61].

$$\sigma^{WH} = \sigma_{\text{NNLOQCD}}^{WH,DY} (1 + \delta_{\text{EWK}}) + \sigma_{t\text{-loop}} + \sigma_{\gamma} \quad (5.4)$$

$$\sigma^{ZH} = \sigma_{\text{NNLOQCD}}^{ZH,DY} (1 + \delta_{\text{EWK}}) + \sigma_{t\text{-loop}} + \sigma_{\gamma} + \sigma^{ggZH} \quad (5.5)$$




Figure 5-6: The shapes for $p_T(V)$ in the V+light control regions are shown for 2017. The left plots are before LO to NLO reweighting, and the right plots are after the correction factor is applied. The top row of plots show the 0-lepton control region, the middle row shows the 1-lepton selection, and the bottom row shows the 2-lepton control region.




Figure 5-7: Electroweak correction as a function of boson $p_T(V)$ for the V+jets samples.




Figure 5-8: Above is an example diagram used to calculate NNLO QCD corrections to the VH production cross section.




Figure 5-9: **Phil** where is the caption??

Phil Please define all of the terms in the formula, I don't know what σ_{t-loop} is even. The full correction shape as a function of $p_T(V)$ is shown in Figure 5-9.

5.4 Application of Uncertainties

In addition to the corrections and their associated uncertainties listed so far, there are additional experimental and theoretical uncertainties. Some of these uncertainties only affect the normalization of samples. Other uncertainties affect the shape of variables that are used in the DNN discriminator described in the following chapter. The final set of uncertainties is a set of migration uncertainties to account for variation near the STXS bin boundaries.

5.4.1 Normalization Uncertainties

The following uncertainties only affect the normalization of the simulated samples.

- The luminosity measurement has an uncertainty of 2.5% for 2016 and 2018, and 2.3% for 2017 [62–64]. Since the method of luminosity measurement was the same for all three years, the uncertainties are partially correlated across the three years.

- The theoretical uncertainty of the branching ratio of the Higgs Boson to bottom quarks is 0.5% [61].
- QCD scale uncertainties for the signal production cross section are implemented as acceptance uncertainties for each of the STXS bins. **Phil** why do you say for each bin, is this to imply some correlation model?
- The uncertainties to the proton's PDF and α_s for the signal processes are 1.6% for ZH production, and 1.9% for WH production.
- At the high $p_T(V)$ regions of this analysis, the theoretical uncertainties grow. For the NLO electroweak corrections, the uncertainties are 2%. For the NNLO QCD correction, the uncertainty is 5%.
- For smaller background processes without a dedicated control region, primarily single-top and di-boson processes, a 15% normalization uncertainty is applied. This number comes from measured cross sections for single-top [65] and di-boson [66].
- The lepton identification efficiency uncertainties are applied as a flat uncertainty to each channel, as appropriate.
- The MET trigger efficiencies are also applied as a normalization uncertainty to the 0-lepton regions. **Phil** you show shape uncertainties before, please explain why you don't apply shape

5.4.2 Shape Uncertainties

In addition to normalization, there is some uncertainty in the shapes of the various distributions that are fit. This is typically caused by uncertainties in variables that are used in the event selection, but can also be a result of normalization uncertainties that only affect a subset of events in a selection. The shape variations are generated by varying different parameters up and down and re-applying selections and weights

to determine the final shape of each distribution. Uncertainties of simulated processes that cause shape differences are the following.

- The energy resolution, and therefore the exact final energy, of simulated b jets are varied by increasing or decreasing the distance between the generator-level p_T and the reconstructed jet p_T .
- The energy scale is varied by varying all jet energies up and down by one standard deviation. **Phil**for this can you put a reference
- The b -tagging efficiency uncertainties are calculated for multiple p_T and η bins. The uncertainties for these different bins are uncorrelated, so they each affect a different subset of events when applied. **Phil**do you have a plot of the uncertainties for b-tagging it would be good to see. Also, can you explain why its uncorrelated, at minimum put a reference
- The b -tagging for fat jets is applied for two different working points, medium and tight, and is binned in p_T . These working points and p_T bins are decorrelated from each other. **Phil** please explain why its uncorrelated. In fact, I think this b-tagging deserves a sub sectin in this chapter where you explainhow these uncertiainties are derived.
- The uncertainties from the limited number of Monte Carlo events is handled using an approximation of the Barlow-Beeston method [67].

Phil This section is pretty light on details, I really think you should elaborate on a few things, I have pointed them out along the way. In particular, I think you should put a page or two on the b-tagging uncertainties. Additionally, you don't talk about the uncertainties in the boosted category (double-b) and how they are different.

5.4.3 Migration Uncertainties

An additional uncertainty is a applied that accounts for simulated events crossing the STXS bin boundaries due to uncertainties in the vector boson p_T . The effect of each

nuisance parameter is anti-correlated between the bins on either side of the boundary. Most bins have two uncertainties allowing events to cross into lower or higher $p_T(V)$ bins, with the exception of the highest STXS bins, which have no upper bounds. Due to limited availability of studies on this effect, these systematic uncertainties are given large values before the global fit is performed, ranging from 20% to 50% for each individual boundary. These are worth noting since they have the largest affect of all systematic uncertainties on the pre-fit simulation uncertainties, though they are significantly constrained by the fit process.

Chapter 6

Analysis Results

6.1 Run 2 Data Collection

The CMS detector collected proton-proton collision data at $\sqrt{s} = 13$ TeV over three years during Run 2 of the LHC. In 2016, CMS collected 35.9 fb^{-1} of data [62]. In 2017, data corresponding to 41.5 fb^{-1} were collected [63]. In 2018, CMS collected 60 fb^{-1} of collision data [64]. This data, totaling about 137 fb^{-1} was used for the analysis.

6.2 Control Region Plots

A number of variables are compared in all of the control regions to ensure that the data and simulation are in reasonable agreement. A selection of these plots are shown in Figures 6-1 through 6-6. These are the distributions before fitting the control and signal regions to data, so agreements improve after the fit. The number of variables, selections, and separation of the data collection years means that this is just a fraction of the total number of plots produced. However, this subset shown demonstrate several discrepancies that needed to be accounted for in our systematics, such as the $p_T(V)$ trend and normalization of certain background samples.




Figure 6-1: The vector boson p_T is plotted for multiple control regions and bins in an inclusive selection of both the $Z \rightarrow \mu\mu$ and $Z \rightarrow ee$ two lepton channels with 2016 data. The top row shows the $t\bar{t}$ control region, the middle row shows the Z + heavy jets, and the bottom row shows the Z + light jets control region. The left column shows the low $p_T(V)$ bin of 75 to 150 GeV, and the right column shows the medium $p_T(V)$ bin of 150 to 250 GeV.




Figure 6-2: The p_T of the jet with a higher DeepCSV value is plotted for multiple control regions and bins in an inclusive selection of both the $W \rightarrow \mu\nu$ and $W \rightarrow e\nu$ one lepton channels with 2016 data. The top row shows the $t\bar{t}$ control region, the middle row shows the W + heavy jets, and the bottom row shows the W + light jets control region. The left column shows the medium $p_T(V)$ bin of 150 to 250 GeV, and the right column shows the high $p_T(V)$ bin of 250 GeV and above.




Figure 6-3: The DeepCSV value of the jet with a higher DeepCSV value is plotted for multiple control regions and bins in an inclusive selection of both the $Z \rightarrow \mu\nu$ and $Z \rightarrow e\nu$ $\text{PhilZ}/\rightarrow\mu\nu$ one lepton channels with 2017 data. The top row shows the $t\bar{t}$ control region, the middle row shows the Z + heavy jets, and the bottom row shows the Z + light jets control region. The left column shows the medium $p_T(V)$ bin of 150 to 250 GeV, and the right column shows the high $p_T(V)$ bin of 250 GeV and above.




Figure 6-4: The p_T of the jet with a lower DeepCSV score is plotted for multiple control regions and bins in the $Z \rightarrow ee$ two lepton channel with 2017 data. The top row shows the $t\bar{t}$ control region, the middle row shows the Z + heavy jets, and the bottom row shows the Z + light jets control region. The left column shows the low $p_T(V)$ bin of 75 to 150 GeV, and the right column shows the medium $p_T(V)$ bin of 150 to 250 GeV. Phil Could you comment in the text on the NLO correction; from what I read its only on the p_T and not on jet distributions like the sub leading jet, this will impact that high boson p_T and low sub leading jet p_T (bottom right figure).




Figure 6-5: The DeepCSV value of the jet with a higher DeepCSV value is plotted for multiple control regions in an inclusive selection that contains all STXS bins with $p_T(V)$ greater than 150 GeV in the one lepton channels with 2018 data. The top row shows the $t\bar{t}$ control region, the middle row shows the W + heavy jets, and the bottom row shows the W + light jets control region. The left column shows the $W \rightarrow e\nu$ channel, and the right column shows the $W \rightarrow \mu\nu$ channel.




Figure 6-6: The vector boson p_T is plotted for multiple control regions and bins in the $Z \rightarrow \mu\mu$ two lepton channel with 2018 data. The top row shows the $t\bar{t}$ control region, the middle row shows the Z + heavy jets, and the bottom row shows the Z + light jets control region. The left column shows the low $p_T(V)$ bin of 75 to 150 GeV, and the right column shows an inclusive selection of the medium and high $p_T(V)$ bins of greater than 150 GeV. I am having a lot of trouble reconciling the chi2 values with what is shown in the plots. Furthermore why is the normalization so different from teh 2017, I hope you comment on this in the text.

6.3 Multivariate Discriminator

In each STXS bin, a multivariate discriminator is plotted which separates the signal events from background events. A Deep Neural Network (DNN) is trained for the resolved selection, and a Boosted Decision Tree (BDT) is trained for the boosted selection.

6.3.1 Resolved DNN

The DNN classifier for distinguishing background and signal events is prepared using Keras with a Tensorflow back-end using an Adam optimizer. It has five hidden layers. The number of nodes in each layer, from input to output, is 512, 256, 128, 64, 64, and 64. The final layer is a softmax layer with the target of predicting the probability of each event belonging to a particular class.

Each channel of 0-, 1-, and 2-leptons is trained separately, and has slightly different input variables. The list of input variables is given in Table 6.1. All variables that are affected by the kinematic fit in the 2-lepton region use the values calculated by the fit. PhilWhat is the loss function?

6.3.2 Boosted BDT

The BDT used to classify signal and background events in the boosted region was trained using ROOT. The model uses 100 trees with 20 cuts and a minimum node size of 0.05. The QCD multi-jet backgrounds were not used in the training since the sample's large weights of individual events affected the training.

The list of input variables for the BDT is the following:

- Soft-drop mass of the reconstructed fat jet
- Transverse momentum of the fat jet
- Transverse momentum of the reconstructed vector boson
- Number of soft-track jets with $p_T > 5 \text{ GeV}$

Table 6.1: The list of input variables used for each DNN training is shown.

| Variable | Explanation | 0-lepton | 1-lepton | 2-lepton |
|--------------------------------------|---|----------|----------|----------|
| m_{jj} | Di-jet mass | ✓ | ✓ | ✓ |
| $p_{T,jj}$ | Di-jet transverse momentum | ✓ | ✓ | ✓ |
| MET | Missing transverse energy | ✓ | ✓ | ✓ |
| $m_{T,V}$ | Vector boson transverse mass | | ✓ | |
| $p_T(V)$ | Vector boson p_T | | ✓ | ✓ |
| $p_{T,jj}/p_T(V)$ | Redundant ratio | | ✓ | ✓ |
| $\Delta\phi(V, jj)$ | Azimuthal angle between vector boson and di-jet | ✓ | ✓ | ✓ |
| $b\text{-tag}_{\text{max}}$ WP | 1, 2, or 3 if higher b -tag discriminate meets the tight, medium, or loose working point respectively | ✓ | ✓ | ✓ |
| $b\text{-tag}_{\text{min}}$ WP | 1, 2, or 3 if lower b -tag discriminate meets the tight, medium, or loose working point respectively | ✓ | ✓ | ✓ |
| $\Delta\eta(jj)$ | η difference between jets | ✓ | ✓ | ✓ |
| $\Delta\phi(jj)$ | Azimuthal angle between jets | ✓ | ✓ | |
| $p_{T,\text{lead}}$ | Leading jet p_T | ✓ | ✓ | ✓ |
| $p_{T,\text{trail}}$ | Trailing jet p_T | ✓ | ✓ | ✓ |
| SA5 | Number of soft jets, $p_T > 5$ GeV | ✓ | ✓ | ✓ |
| N_{aj} | Number of additional jets | ✓ | ✓ | |
| $b\text{-tag}_{\text{add}}$ | Maximum b -tag of additional jets | ✓ | | |
| $p_{T,\text{add}}$ | Maximum p_T of additional jets | ✓ | | |
| $\Delta\phi(\text{add}, \text{MET})$ | Azimuthal angle between additional jet and MET | ✓ | | |
| $\Delta\phi(\ell, \text{MET})$ | Azimuthal angle between lepton and MET | | ✓ | |
| m_t | Reconstruction top mass | | ✓ | |
| m_V | Vector boson mass | | | ✓ |
| $\Delta R(V, jj)$ | Separation between vector boson and di-jet | | | ✓ |
| ΔR_{jj} | Separation between jets | | | ✓ |

- Double b -tagger output node for boosted jets

All of these variables were used in the 0-, 1-, and 2-lepton regions, even though the regions were trained separately.

6.4 Combination Fit

A simultaneous fit is run over all channels, control regions, and the signal selection region in order to determine the most likely values for all parameters with systematic uncertainties, called nuisance parameters, as well as the most likely scale factors for all the MC backgrounds and signal. The fit is done by using the `combine` tool [68] as part of `CMSSW_10_2_13` [56]. Part of the method for discovery involves finding the most likely values for all nuisance parameters and signal strength simultaneously. This maximum likelihood feature is used for the measurement of signal strength in the STXS bins. The difference in signal strength uncertainty when allowing the nuisance parameters to float from their most likely values or not show the overall effect of the systematic uncertainties on the final measurement.

In addition, `combine` includes a variety of diagnostic tools to ensure the assumed model is adequate. The first step before unblinding results is to throw toys using the best-fit nuisance parameter values and comparing a test statistic of these toys to that of the observed data. **Phil**You need a paragraph talking about what unblinding means, there are many ways to define this. The Higgs approach is specific to the Higgs group, B2G for example has a very different unblinding procedure. The distributions of a the values from the set of toys allows the evaluation of the p -value of the test statistic observed in data. **Phil**of the values The `combine` tool also evaluates the impacts of each nuisance parameter on the likelihood by varying them individually. If nuisance parameters with large impacts are pulled far from their initial value, that means that an important aspect of the measurement was not well understood.

The final results will therefore show comparison of data and best-fit toys, called Goodness of Fit, as well as the final scan results. A post-fit plots will also reveal the way that the fit moves the background and signal distributions within uncertainties

in order for the background to describe the data. **Phil** A post-fit=> Post-fit Impacts plots are shown in Appendix D due to the high multiplicity of nuisance parameters. Before running these checks and measurements for the VH measurement, they were first tested on VZ . **Phil** I think this section could use an intro section explaining the strategy towards getting the results.

6.4.1 VZ Cross Check Analysis

The VZ cross check analysis mostly duplicated the VH . **Phil** This needs more than one sentence, also your sentence is not grammatically correct The primary differences are that a di-boson sample with generator-level b hadrons produced is used as the signal sample, and the signal mass window is moved from $90 \text{ GeV} < m_{jj} < 150 \text{ GeV}$ to $60 \text{ GeV} < m_{jj} < 120 \text{ GeV}$. However, other differences should be mentioned. Most of the potential effects of these differences are difficult to quantify, with the exception of the following. Additional measured electroweak corrections are applied to the di-boson samples for each STXS bin. The values of these corrections vary from as low as 4% for the low $p_T(V)$ bins and as high as 23% for the high $p_T(V)$, but all will increase the measured signal strength compared to VH . Other differences include the dedicated VZ DNN training and the fact that the Z boson will decay into light flavored jets at a higher proportional rate than the Higgs boson. The DNN training for VH has to account for the similar VZ background, while the VZ is not influenced as much by the rare VH process. Also, the light flavor VZ process will have more influence over the control region fits than the VH process has.

Data from all three years of Run 2 are used again so that any differences that may affect the VH analysis can also be investigated using VZ . The Goodness of Fit plots are shown in Figure 6-7. **Phil** You need a reference for GOF, use Bob Cousins' paper on this The high p -value means that the simulated processes were able to match the data very well while staying within the fit constraints of the systematic uncertainties. **Phil** Do you have the GOF without systematics, or reducing the systematics, it would be good to show in your thesis as a justification of this explanation. Such a low GOF is worrisome for me otherwise.




Figure 6-7: The Goodness of Fit test results are shown for VZ. The fit is performed using STXS bins, and the test statistic is generated by toys for all control regions and signal regions.

Due to the splitting of STXS bins, the multiple channels for each bin, and the combination of three different years, 486 distributions are fit in the VZ cross check analysis. **Phil**The caculation of 486 should go in your talk, I am struggling to get to that number. $486/3\text{years}/3\text{finalstates}/6\text{categories?} = 9.9$ seems to high for the number of categories. For the sake of brevity, the effect of the fit will only be shown on eight of these distributions for the VZ cross-check analysis. These are split into two figures, each showing distributions closely related to each other. On the left of each figure are simulation values and uncertainties before the nuisance parameters are fit to match the simulation to data. On the right are the simulated values after fitting the nuisance parameters. In the lower parts of each plot, ratios between observed data and expected yields improve as a result of the fit. The uncertainties of the simulations also decrease as a result of the fit. Some nuisance parameters end up with smaller individual uncertainties from the fit, «««< HEAD including the initially large migration uncertainties. Figure 6-8 shows this decrease in bin migration




Figure 6-8: The most significant impacts in on WH $250 < p_T(V) < 400$ GeV. Note the decrease in migration uncertainties, such as CMS_vhbb_Vpt250_TT_1lep_13TeV2016.

uncertainties, especially in $t\bar{t}$ in line 19. The uncertainty also decreases since each named nuisance parameter is at first assumed to be uncorrelated to the other nuisance parameters. The fit reveals many of the nuisance parameters to be correlated, decreasing the overall possible variation. ===== but the main reason for the dramatic improvement in the uncertainty is that each named nuisance parameter is at first assumed to be uncorrelated to the other nuisance parameters. The fit reveals many of the nuisance parameters to be correlated, decreasing the overall possible variation. Phil This explanation does not make sense to me. If you have two nuisances that are uncorrelated, they remain uncorrelated throughout the fit. If they are both describing the same thing they will each get constrained so that the sum of their uncertainties in quadrature is equal to the data. Thinking through, I think I knwo what you are saying, but you should rephrase this. >>>> phil/phil

Figure 6-9 shows all of the distributions fit for the low p_T bin in the $Z \rightarrow \mu\mu$

channel in 2017. Several additional features aside from the pre-fit and post-fit differences are worth noting. There is only one bin in the $t\bar{t}$ control region, so it is only used to constrain the normalization of the $t\bar{t}$ background. In the $Z + \text{heavy jets}$ control region, the DNN is trained with the same inputs as the signal region DNN. This constrains the shape uncertainties that would affect both classifiers. In all of the regions, we can see improved agreement between the simulation and data after the fit, and the combination fit also decreases the uncertainty on each bin. The relative contributions of the main background processes can also be seen at a glance at the signal region plots. These distributions can be compared to those in Figure 6-10, which shows distributions from the boosted selection of the $W \rightarrow e\nu$ channel in 2018. In this case, the $W + \text{heavy jets}$ control region is not shown because the related process contributes less in the signal region. Instead, the $t\bar{t}$ background is dominant in this boosted topology. **Phil** is dominant. The control regions use the double b -tag score of the fat jet instead of $p_T(V)$ for the distribution. The signal region is also split into two higher $p_T(V)$ bins, while the control regions stay combined to constrain both STXS bins. As in the low p_T bin, though, both agreement between simulation and data and estimated uncertainties are improved as a result of the fit.

From these post-fit distributions, the strength and uncertainty of the signal samples are extracted. A value of 1.0 corresponds to the expectation based on simulation of the Standard Model. By correlating all of the signal samples across STXS bins, an inclusive signal strength is obtained. The likelihood scan of this strength is shown in Figure 6-11. The maximum likelihood with respect to the Standard Model is $1.182^{+0.118}_{-0.112}(\text{stat})^{+0.098}_{-0.097}(\text{sys})$. The uncertainty of this measurement means the result has a p -value of 22%, assuming Standard Model couplings, or within 1.2 standard deviations of the Standard Model. Figure 6-12 shows the measured kinematic distributions of the various VZ processes. Assuming the Standard Model, the measurement has a p -value of 23%. Overall, the VZ cross-check is in agreement with the Standard Model.

Phil You should show the WZ and ZZ combined results, also do you have an explanation why WZ in particular is so off? It looks like 2.5 sigma




Figure 6-9: Above are $Z \rightarrow \mu\mu$ pre-fit (left) and post-fit distributions (right) for the low p_T STXS bin in 2017 in the VZ cross-check analysis. The top row shows the Z + light jets control region, the second row shows the Z + heavy jets control region, and the third row shows the $t\bar{t}$ control region. The bottom row shows the signal region. **Phil** Can you make the signal region plot aggregated over the years, ideally weighted by their significance contribution, also can you make it log so that we see the VZ contribution lastly can you make the VZ signal the redline not the VH




Figure 6-11: The likelihood scan of the inclusive signal strength of the VZ cross check analysis is shown above. The dashed line is generated by freezing all nuisance parameters to their most likely value, so the likelihood variation for that curve is from statistical uncertainties only. Using quadratic subtraction between the two curves allows the systematic uncertainties to be evaluated.




Figure 6-12: The measured most likely values of all STXS bins in the VZ cross check analysis. **Phil** Can you show the ggZH and qqZH split as well. Also, the WZ vs ZZ 2D contour, it seems that the scale factors are quite different

6.4.2 VH Combination Fit Results

Phil big comment about this section is you should make some summary distributions combining the signal across years ideally (weighted by significance in fit). Also all of your DNN plots need to be log, I can't see the signal otherwise. The first step to show that the processes in the detector are well-modeled by simulation is the Goodness of Fit test. The Goodness of Fit for the VH selection is shown in Figure 6-13. The observed value for the saturated test statistic is in the bulk of the distribution created by throwing post-fit toys, with an associated p -value of 70%, suggesting both good modeling and reasonable values for post-fit uncertainties.

A selection of post-fit distributions are also shown. Like, the VZ analysis, there are too many distributions to show all of them in this work, so what follows are only a few illustrative examples. Figure 6-14 shows the zero-lepton channel in the high $p_T(V)$ regime. Unlike the two lepton selections shown, the signal distributions display contamination from $W + \text{jets}$ background processes. There is no zero-lepton




Figure 6-13: The Goodness of Fit test results are shown for VH . The fit is performed using STXS bins, and the test statistic is generated by toys for all control regions and signal regions.

control region for $W + \text{jets}$. Instead, these backgrounds are controlled only by one-lepton control regions, so the five channels are correlated in the fit. Figure 6-15 shows the $W\mu\nu$ distributions for the medium $p_T(V)$ STXS bin. Like in the VZ cross-check analysis, the single lepton signal selection is largely made up of $t\bar{t}$ background. Figure 6-16 shows the medium $p_T(V)$ bin for $Z \rightarrow ee$. In the ZH STXS bins, the signal region is split between events with no extra jets, and events with extra jets, which can be seen in the distributions.

The signal strength extracted from the fit can be separated a number of ways. The inclusive signal strength is shown in Figure 6-17 to be $0.568^{+0.154}_{-0.147}(\text{stat})^{+0.134}_{-0.133}(\text{sys})$. The uncertainties put this measurement within 2.1 standard deviations of the Standard Model. Figure 6-18 shows that this low value is caused by a low cross section measurement of ZH , which alone is 2.9 standard deviations below the Standard Model. Figure 6-19 shows the full STXS measurement results. Both ZH and WH show increasing trends in differential cross section as $p_T(V)$ increases. However, substituting




Figure 6-14: Above are $Z \rightarrow \nu\nu$ pre-fit (left) and post-fit distributions (right) for the high p_T STXS bins in 2016 in the VH measurement using the resolved selection. The top row shows the Z + light jets control region, and the second row shows the Z + heavy jets control region. The third row shows the signal region for the $250 \text{ GeV} < p_T(V) < 400 \text{ GeV}$ bin, and the bottom row shows the signal region for the bin of $p_T(V) > 400 \text{ GeV}$.




Figure 6-15: Above are $W \rightarrow \mu\nu$ pre-fit (left) and post-fit distributions (right) for the medium p_T STXS bins in 2017 in the VH measurement. The top row shows the $W + \text{light jets}$ control region, and the second row shows the $W + \text{heavy jets}$ control region, and the third row shows the $t\bar{t}$ control region. The bottom row shows the signal region.




Figure 6-16: Above are $Z \rightarrow ee$ pre-fit (left) and post-fit distributions (right) for the medium p_T STXS bins in 2018 in the VH measurement. The top row shows the Z + light jets control region, and the second row shows the Z + heavy jets control region. The third row shows the signal region with no additional jets, and the bottom row shows the signal region with additional jets. **Phil** where is the highest DNN score. If it fell off the plt adjust the range.




Figure 6-17: The likelihood scan of the inclusive signal strength of the VH analysis.

the inclusive signal strength in for the Standard Model, the STXS scans have an agreement with a p -value of 9.3%. Standard Model kinematics cannot be entirely ruled out by this measure.

Philgiven how weird the results are have you tried injecting other Higgs mass points?

The total coupling between the Higgs boson and vector bosons, as well as the coupling between the Higgs boson and bottom quarks is measured using the kappa framework [69]. Results of this analysis alone are shown in Figure 6-20. Standard Model values of $\kappa_V = \kappa_b = 1$ fall within the 68% confidence level. The results of this κ scan can be additionally constrained by combining this analysis with others that focus on using either vector bosons for Higgs production or the $b\bar{b}$ decay path.

«««< HEAD ===== **Phil**This chapter really does feel incomplete. You need to comment about the elephant in the room (large and systematic disagreement). Othwerise, it reads really weird. There is nothing evidently incorrect with this approach, so you should stand by your results, and be very clear that things are notn




Figure 6-18: The measured most likely values of WH and ZH in the VH analysis.




Figure 6-19: The measured most likely values of all STXS bins in the VH analysis.




Figure 6-20: The possible values of κ_V and κ_b are shown above. Values of 1.0 reproduce Standard Model predictions.

consistent with the SM. $\ggg>$ phil/phil

Chapter 7

Conclusions

Good agreement between the STXS measurement and Standard Model predictions is observed. **Phil**Please don't say good agreement. There is no such thing, what is good? Quantify the number. This is a major pet-peeve of mine, I will not sign the thesis until you remove this sentence, and put a number. However, study of the Higgs boson, and even study of $H \rightarrow b\bar{b}$ in associated production, is not finished. This analysis is only one of many to characterize the Higgs boson, and this cross section measurement itself will be improved as more data is gathered and measurement techniques are refined.

The first published searches of the Higgs in associated production by CMS were given in 2011 [70], As mentioned in the beginning of this document, the search concluded successfully in 2018 After the discovery, confirmed also by ATLAS, a joint effort produced the STXS framework for additional study of the Higgs boson [71]. **Phil**Can you compare your result with ATLAS, is the uncertainty better, how about the agreement. What about the trends in p_T . This is the stage the presented analysis comes in. ATLAS has already published results of an STXS measurement using Run 2 LHC data [72], and this work will allow a direct comparison between the two experiments. Standard Model confirmation of this independent experiment also immediately serves to reduce any uncertainties from previous analyses. **Phil**Can you rewrite this sentence, I find it very confusing Though the binning and results of this analysis may appear rudimentary in the context of what is required to dispute the Standard Model, the development of the techniques is crucial for studies to come.

While it is impossible to predict the future precisely, international effort is working on two projects to continue to study the Higgs. **Phil**an international? confused The LHC is receiving an upgrade, in what is called the High Luminosity LHC (HL-LHC). All cutting-edge technologies will be used to create a machine that will gather an order of magnitude more data suitable for Higgs studies and other searches beyond the Standard Model [73]. Beyond that, the Circular Electron Positron Collider (CEPC) is currently being designed for use as a Higgs factory for precision studies [74]. **Phil**why pick this one, should you not mention FCC? These experiments will produce very fine-grained descriptions of the Higgs differential cross section, and have the potential to uncover discrepancies from the Standard Model. They may allow us to understand just a little bit more about our universe, along with any other exciting results from physics experiments looking elsewhere.

PhilCan you rewrite the last paragraph. The words are a little jumbled, but the message is nice. It would be good to end your thesis on a high note!

Appendix A

Detector Projects

Each collaborator must contribute to the operation of the CMS detector before his or her name is added to the author list. The operation of the detector is distinct from analyzing the data generated by the detector, so all collaborators must adopt some role outside of being a physicist.

This appendix details projects I completed in order to contribute to the operation of the CMS detector. The first project presented is the Dynamo Consistency project. It is a plugin for the dynamic data management system Dynamo [75] that compares the inventory of files Dynamo expects at a site with the files that are actually at a site. The other project described is known as Workflow Web Tools. This is a dynamic web server that displays errors reported by the CMS computing infrastructure to operators, and allows those operators to perform corrective actions through the web page. Workflow Web Tools also tracks operator actions for future use in training various machine learning models. Both projects are published as software packages written in Python [76,77] and are available through the Python Package Index (PyPI) as `dynamo-consistency` and `workflowwebtools`, respectively.

A.1 Dynamo Consistency

Dynamo Consistency is a plugin for Dynamo Dynamic Data Management System that checks consistency between Dynamo’s inventory and files actually located at managed

sites. Even though Dynamo controls and tracks the history of file transfers between computing sites, a separate check is needed to ensure files are not lost or accumulated due to user or system errors. For example, sites that can no longer access some files after a power outage can cause problems for many related activities. File transfers requested from a inconsistent site to another site will fail when files are missing. Sites will be also be chosen incorrectly for production jobs that assume the presence of a local file. Last disk copies may also be missing, causing a significant delay when a user requests data. Another type of inconsistency arises when files thought to be deleted are still on disk. This leads to wasted disk space for files that are not accessed, except by accident. Dynamo Consistency regularly checks consistency by listing each remote site and comparing the listed contents to Dynamo’s inventory database. The results are reported back to Dynamo, which can then take corrective measures.

A single executable `dynamo-consistency` is provided to run the consistency check. This executable can be used directly in Dynamo’s scheduling system. Most of the behaviour is controlled via a single JSON configuration file, with options for site selection, passed via command line arguments. Differing command line arguments allows Dynamo to run separate schedules for differing site architectures.

Because Dynamo runs in a heterogeneous computing environment, different sites need to be listed remotely using different methods. Currently implemented are listings using XRootD Python bindings, the `gfal-ls` CLI [78], and a `xrd fs` subshell. These listers are easily extensible in Python, allowing for new site architectures to be added to Dynamo Consistency as well.

The default executable performs the check as expected, listing files that are not tracked by Dynamo as orphans and listing files that are not found at sites as missing. A few configurable filters can be added to modify these lists. Dynamo Consistency avoids listing orphan files that have a modification time that is recent. Paths to avoid deleting can also be set. Deletion and transfer requests that are queued are also used to filter the final report to avoid redundant actions from Dynamo.

In addition to tracking the consistency between Dynamo’s inventory and physical site storage, Dynamo Consistency can report all remote files older than a certain age

in general directories. These files can also be filtered with path patterns, just as the regular consistency check. The time-based only reporting allows for cleaning of directories that Dynamo does not track. This is a setting recommended for large file systems that are written to with a high frequency.

Summaries of check results, as well as the statuses of running checks, are displayed in a web page. The page consists of a table that includes links to logs and lists of orphan and missing files. Cells are color coded to allow operators to quickly identify problematic sites. Historic summary data for each site is also accessible through this page.

If the available configuration options and listers are not enough, advanced users can also directly use the Python API to run a custom consistency check. For more details on the Dynamo Consistency package, see [79].

A.1.1 Installation

Dynamo Consistency requires the XRootD [80] Python module to be installed separately. In addition, it uses the Dynamo Dynamic Data Management package to get inventory listings and to report results of the consistency check. Any other needed packages are installed with Dynamo Consistency during installation.

The simplest way to install is through pip:

```
pip install dynamo-consistency
```

The source code is maintained on GitHub [81]. Other typical `setuptools` methods are supported by the repository's `setup.py`.

A.1.2 Inventory Listing

Two listings must be done to compare. One is the Inventory Listing, and the other is the Remote Listing. This section describes the inventory listing, and the next describes remote listing.

The inventory is queried before the site is listed remotely due to possible race conditions. It is not uncommon for a site listing to take multiple days. In the

meanwhile, two things can change in the inventory. A file can be deleted from a site or it can be added to a site. An added file is ignored by setting **IgnoreAge** in the configuration to a large enough value. Files that are deleted during the remote listing are filtered out by checking recent deletion requests after the remote listing.

There are currently multiple ways to get the site contents from Dynamo. One is to access the MySQL database use for Dynamo storage directly. This will work as long as the schema does not change. A more reliable way to keep up with major changes in Dyanmo is to use the Dynamo inventory object. This method is less optimized when working with the MySQL storage plugin, but will work for different schema and any different storage types that are added in the future.

The type of inventory lister is selected via command line options, or by setting `dynamo_consistency.opt.V1` to `True` or `False` before importing any modules that rely on the backend. By implementing the three modules `inventory`, `registry`, and `siteinfo`, described in the full documentation [79], any other method of communicating with an inventory can be added.

After selecting the backend, the inventory can be listed transparently using the following method:

```
from dynamo_consistency import inventorylister
listing = inventorylister.listing(sitename)
```

Here, `listing` is a `dynamo_consistency.datatypes.DirectoryInfo` object that is the root node of the full directory tree. Each node of `DirectoryInfo` contains meta data about a directory, such as its modification timestamp and name. It also holds a list of sub-directories, in the form of `DirectoryInfo` objects, and a list of files. The files are represented as dictionaries containing the name, size, and modification time of the file. Each file and `DirectoryInfo` also stores a hash of the meta data. The `DirectoryInfo` hash includes information from the object's files and sub-directories too. This is to speed up the file tree comparison, shown in Figure A-1




Figure A-1: Comparison algorithm

A.1.3 Remote Listing

The remote listing is equally flexible in terms of having multiple implementations used for the listing. The factory function `dynamo_consistency.backend.get_listers()` reads the configuration file to determine the type of lister for each site. There are currently three different classes implemented, and more can be added by extending the `dynamo_consistency.backend.listers.Lister` class and implementing its `ls_directory` method. The three current listers are the following:

- `dynamo_consistency.backend.listers.XRootDLister` - This listing object uses the `XRootD` Python module to connect to and query each site.
- `dynamo_consistency.backend.listers.GFallister` - This listing object uses the `gfal-ls` command line tool to list remote sites.
- `dynamo_consistency.backend.listers.XRootDLister` - This listing object opens a subshell using the `xrdfs` command line tool and queries the remote site.

Once the type of lister is set in the configuration the contents of the remote site can be gathered with a simple interface:

```

from dynamo_consistency import remotelister
listing = remotelister.listing(sitename)

```

This takes much longer than the Inventory Listing, since every directory of the site needs to be queried over the network. A software layer between the listing class and

the final output creates multiple connections and works on two queues with multiple threads. There is the input queue, which is a list of directories that still need to be listed, and an output queue which holds the result of each directory listed so far. The workflow of the listing is shown in Figure A-2.



Figure A-2: Listing algorithm with retries

A.1.4 Executables

Other tools are available as part of **dynamo-consistency** in order to simplify operating the system. Many of these come as separate executables. A list of some of the executables installed with the package is given below.

dynamo-consistency

This program runs the Site Consistency Check for Dynamo Dynamic Data Management System.

Usage: **dynamo-consistency** [options]

Options:

| | |
|----------------------------|--|
| <code>--version</code> | show program's version number and exit |
| <code>-h, --help</code> | show this help message and exit |
| <code>--config=FILE</code> | Sets the location of the configuration file to read. |

Selection Options:

| | |
|-------------------------------------|--|
| <code>--site=PATTERN</code> | Sets the pattern used to select a site to run on next. |
| <code>--lock=NAME</code> | Sets the lock name that should be used for this run. |
| <code>--date-string=YYYYMMDD</code> | Set the datestring to pull for RAL-Reader listers |

Logging Options:

| | |
|-------------------------------|---|
| <code>--update-summary</code> | Forces the update of the summary table, even if loading trees |
| <code>--email</code> | Send an email on uncaught exception. |
| <code>--info</code> | Displays logs down to info level. |
| <code>--debug</code> | Displays logs down to debug level. |

Behavior Options:

These options will change the backend loaded and actions taken

| | |
|-----------------------------|--|
| <code>--no-orphan</code> | Do not delete any orphan files. |
| <code>--cms</code> | Run actions specific to CMS collaboration data. |
| <code>--no-sam</code> | Disables the SAM readiness check. |
| <code>--more-logs</code> | Clean any "AdditionalLogDeletions" directories. |
| <code>--no-inventory</code> | Do not connect the inventory. Used to test unmerged |
| <code>--unmerged</code> | Run actions on "/store/unmerged". |
| <code>--v1</code> | Connect to Dynamo database directly |
| <code>--v1-reporting</code> | Connect to Dynamo database directly for registry only. |
| <code>--cnf=FILE</code> | Point to a non-default location of a "my.cnf" file. |
| <code>--test</code> | Run with a test instance of backend module. |

set-status

This script changes the status of a site on the summary webpage. It can be used to unlock from a dead process, disable sites from running, and change whether or not to act on the site. This script can take a `-config <FILE>` parameter to point to a configuration file, a la `dynamo-consistency`. For the last two arguments, `SITE` will

Table A.1: Valid statuses for sites as tracked by `dynamo-consistency` are described below.

| Action | Description |
|----------------------|--|
| <code>ready</code> | This sets the site status back to idle. This means the site is ready to run. Should be used on a site that's disabled. |
| <code>halt</code> | This stops a currently running or locked site. This site is still eligible to run. |
| <code>disable</code> | Can be applied to a site that is either running or ready. It halts the site and also prevents it from running until set to <code>ready</code> again. |
| <code>act</code> | Marks a site as one to report results to the registry. |
| <code>dry</code> | Opposite of <code>act</code> , this action prevents this site from making entries into the registry in future runs. |

match the name of the site to change. `ACTION` can be one of the entries in Table A.1

Usage: `set-status [options] SITE ACTION`

Options:

```
--version      show program's version number and exit
-h, --help     show this help message and exit
--config=FILE  Sets the location of the configuration file to read.
```

Logging Options:

```
--info        Displays logs down to info level.
--debug       Displays logs down to debug level.
```

`consistency-dump-tree`

Dumps the `dynamo_consistency.datatypes.DirectoryInfo` tree into the cache directory. By default, it dumps the tree that would be read from the inventory.

If the `[NAME]` argument is not given, defaults to `inventory.pkl` or `remote.pkl` when using the `--remote` option.

Usage: `consistency-dump-tree [options] [NAME]`

Options:

| | |
|----------------------------|--|
| <code>--version</code> | show program's version number and exit |
| <code>-h, --help</code> | show this help message and exit |
| <code>--config=FILE</code> | Sets the location of the configuration file to read. |

Selection Options:

| | |
|-------------------------------------|--|
| <code>--site=PATTERN</code> | Sets the pattern used to select a site to run on next. |
| <code>--remote</code> | Dump the remote site listing instead of the inventory |
| <code>--date-string=YYYYMMDD</code> | Set the datestring to pull for RAL-Reader listers |

Logging Options:

| | |
|----------------------|------------------------------------|
| <code>--info</code> | Displays logs down to info level. |
| <code>--debug</code> | Displays logs down to debug level. |

Behavior Options:

These options will change the backend loaded and actions taken

| | |
|-------------------------|---|
| <code>--unmerged</code> | Run actions on <code>"/store/unmerged"</code> . |
| <code>--v1</code> | Connect to Dynamo database directly |
| <code>--test</code> | Run with a test instance of backend module. |

check-phedex

This program is only useful for double-checking CMS sites. This program checks a site's orphan files against PhEDEx. If any of the datasets are supposed to be at the site, this gives a non-zero exit code.

Usage: `check-phedex [options] SITE`

Options:

| | |
|------------------------|--|
| <code>--version</code> | show program's version number and exit |
|------------------------|--|

`-h, --help` show this help message and exit
`--config=FILE` Sets the location of the configuration file to read.

Logging Options:

`--info` Displays logs down to info level.
`--debug` Displays logs down to debug level.

A.1.5 Configuration

The configuration file for `dynamo-consistency` is a JSON or YAML file with the following keys.

- **AccessMethod** - A dictionary of access methods for sites. Sites default to XRootD, but setting a value of **SRM** causes the site to be listed by `gfal-ls` commands.
- **AdditionalLogDeletions** - A dictionary that lists which directories have logs to be cleaned for different sites. These log directories are treated the same as log directories in `/store/unmerged`. This means they use the **UnmergedLogsAge** parameter to determine cleaning policy.
- **DirectoryList** - A list of directories inside of **RootPath** to check consistency.
- **DeleteOrphans** - By default, is true. If set to false, orphan files will all be filtered out so that none are deleted.
- **FreeMem** - The amount of free memory that is required for a check to run. The memory is given in GBs.
- **GFALThreads** - The number of threads used by the GFAL lists
- **GlobalRedirectors** - The redirectors to start all locate calls from, unless looking for a site that is listed in the **Redirectors** configuration.
- **IgnoreAge** - Ignore any files or directories with an age less than this, in days.

- **IgnoreDirectories** - The check ignores any paths that contain any of the strings in this list.
- **InventoryAge** - The age, in days, of how old the information from the inventory can be
- **ListAge** - The age, in days, of how old the list of files directly from the site can be
- **ListDeletable** - Configuration for unmerged cleaning “listdeletable” module. Details on some of the configuration parameters are documented online [82].
- **MaxMissing** - If more files than this number are missing, then there will be no automatic entry into the register.
- **MaxOrphan** - If more than files than this number are orphan files at a site, then there will be no automatic entry into the register.
- **NumThreads** - The number of threads used by the XRootD listers
- **PathPrefix** - A dictionary of prefixes to place before **RootPath** in the XRootD call. This allows for different paths for different sites. If the prefix is not set for a site, and it fails to list **RootPath**, it falls back to a default `/cms` before giving up.
- **RedirectorAge** - The age, in days, of how old the information on doors from redirectors can be. If this value is set to zero, the redirector information is never refreshed.
- **Redirectors** - A dictionary with keys of sites with hard-coded redirector locations. If a site is not listed in this way, the redirector is found by matching domains from `CMSToolBox.siteinfo.get_domain()` to redirectors found in a generic `xrd fs locate` call.
- **Retries** - Number of retries after timeouts to attempt

- **RootPath** - The directory where all of the listed subdirectories will be under. For CMS sites, this will be `"/store"`
- **SaveCache** - If set and evaluates to True, copies old cached directory trees instead of overwriting
- **Timeout** - This gives the amount of time, in seconds, that you want the listing to try to run on a single directory before it times out.
- **Unmerged** - A list of sites to handle cleaning of `/store/unmerged` on. If the list is empty, all the sites are managed centrally
- **UnmergedLogsAge** - The minimum age of the unmerged logs to be deleted, in days
- **UseLoadBalancer** - A list of sites where the main redirector of the site is used
- **UseTransferQueue** - If true, put missing files into transfer queue table when using `--v1` for reporting. Defaults to true value.
- **VarLocation** - The location for the varying directory. Inside this directory will be:
 - Logs
 - Redirector lists
 - Cached trees
 - Lock files
- **WebDir** - The directory where text files and the sqlite3 database are stored

Configuration parameters can also be quickly overwritten for a given run by setting an environment variable of the same name.

A.1.6 Comparison Script

The full set of operations in a typical run of the `dynamo-consistency` is enumerated below.

1. Points `config.py` to the local `consistency_config.json` file
2. Notes the time, and if it's daylight savings time for entry into the summary database
3. Reads the list of previous missing files, since it requires a file to be missing on multiple runs before registering it to be copied
4. It gathers the inventory tree by calling
`dynamo_consistency.getinventorycontents.get_db_listing()`.
5. Creates a list of datasets to not report missing files in. This list consists of deletion requests fetched from PhEDEx by
`dynamo_consistency.checkphedex.set_of_deletions()`
6. It creates a list of datasets to not report orphans in. This list consists of the following.
 - Datasets that have any files on the site, as listed by the dynamo MySQL database
 - Deletion requests fetched from PhEDEx (same list as datasets to skip in missing)
 - Any datasets that have the status flag set to 'IGNORED' in the dynamo database
 - Merging datasets that are protected by Unified
7. It gathers the site tree by calling
`dynamo_consistency.getsitecontents.get_site_tree()`. The list of orphans is used during the running to filter out empty directories that are reported to the registry during the run.

8. Does the comparison between the two trees made, using the configuration options concerning file age.
9. If the number of missing files is less than **MaxMissing**, the number of orphans is less than **MaxOrphan**, and the site is under the webpage’s “Debugged sites” tab, connects to a dynamo registry to report the following errors:
 - Every orphan file and every empty directory that is not too new nor should contain missing files is entered in the deletion queue.
 - For each missing file, every possible source site as listed by the dynamo database, (not counting the site where missing), is entered in the transfer queue. Creates a text file full of files that only exist elsewhere on tape.
10. Creates a text file that contains the missing blocks and groups.
11. .txt file lists and details of orphan and missing files are moved to the web space
12. If the site is listed in the configuration under the **Unmerged** list, the unmerged cleaner is run over the site:
 - `dynamo_consistency.getsitecontents.get_site_tree()` is run again, this time only over `/store/unmerged`
 - Empty directories that are not too new nor protected by Unified are entered into the deletion queue
 - The list of files is passed through the unmerged cleaner
 - The list of files to delete from unmerged cleaner are entered in the deletion queue
13. The summary database is updated to show the last update on the website

A.2 Workflow Web Tools

Datasets used by the CMS collaboration are produced centrally by the Workflow Team. During production of these datasets, errors can arise from things such as

missing files, too much memory usage, or loss of functionality at a particular site. For each failed dataset, Workflow Team operators must gather information from error reports and decide how to recover in order to complete the workflow.

The Workflow Web Tools has the dual purpose of displaying these errors as well as allowing operators to submit actions directly to the web page, instead of requiring them to open up separate shells to script actions. The existence of this layer, which is aware of both the error reports and the submitted actions, also allows for investigation into automatically attempting to recover simple errors.

A.2.1 Error Information

A matrix of errors is built for each task of each workflow. Each row of the matrix corresponds to a different error code that was reported back as a job failed to complete. Each column corresponds to which computing site that the job was running at. The cell simply contains the number of jobs that threw the particular error code while running at a particular site. At the time when the error report is gathered, Workflow Web Tools also gathers the site readiness reports of each computing site. This can be correlated with the error reports to see if jobs failed due to site problems, rather than problems with the workflow submission itself. An example of the display for this error information is given in Figure A-3.

A.2.2 Possible Actions

Workflows are typically completed after throwing errors by either rerunning the failed jobs, called ACDC, or by discarding the entire workflow, and restarting it. Both options often include adjustment of parameters such as memory requested and the number of events run per job. These actions are all set with a number of radio buttons, meaning that there is a finite set of actions that an operator can take. This makes it easier for an automated system to learn and take over some operations.

Each action submitted by an operator is stored in a database and exposed via an API for Unified, the backend manager of workflows, to read the actions from the

Appendix B

Simulation Details

The datasets use for each year are given in Table B.1, Table B.2, and Table B.3.

| Sample Name | Xsec (pb) | KFactor | Total Events | Weight |
|--|------------|---------|--------------|----------|
| DYBJetsToLL_M-50_Zpt-100to200_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 3.086 | 1.23 | 8.38e+06 | 1.63e-02 |
| DYBJetsToLL_M-50_Zpt-200toInf_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 0.3167 | 1.23 | 1.28e+06 | 1.10e-02 |
| DYJetsToLL_BGenFilter_Zpt-100to200_M-50_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 3.414 | 1.23 | 2.67e+05 | 5.65e-01 |
| DYJetsToLL_BGenFilter_Zpt-200toInf_M-50_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 0.5084 | 1.23 | 1.29e+05 | 1.75e-01 |
| DYJetsToLL_M-50_HT-100to200_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 147.3 | 1.23 | 9.37e+06 | 6.96e-01 |
| DYJetsToLL_M-50_HT-1200to2500_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 0.1512 | 1.23 | 5.96e+05 | 1.12e-02 |
| DYJetsToLL_M-50_HT-200to400_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 41.04 | 1.23 | 8.65e+06 | 2.10e-01 |
| DYJetsToLL_M-50_HT-2500toInf_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 0.003659 | 1.23 | 3.99e+05 | 4.06e-04 |
| DYJetsToLL_M-50_HT-400to600_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 5.676 | 1.23 | 7.66e+06 | 3.28e-02 |
| DYJetsToLL_M-50_HT-600to800_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 1.359 | 1.23 | 8.29e+06 | 7.26e-03 |
| DYJetsToLL_M-50_HT-800to1200_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 0.623 | 1.23 | 2.67e+06 | 1.03e-02 |
| DYJetsToLL_M-50_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 4954.0 | 1.0 | 1.46e+08 | 1.22e+00 |
| QCD_HT100to1500_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 1208.0 | 1.0 | 1.52e+07 | 2.86e+00 |
| QCD_HT100to200_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 27940000.0 | 1.0 | 8.23e+07 | 1.22e+04 |
| QCD_HT1500to2000_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 120.4 | 1.0 | 1.18e+07 | 3.66e-01 |
| QCD_HT2000toInf_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 25.27 | 1.0 | 5.97e+06 | 1.52e-01 |
| QCD_HT200to300_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 1712000.0 | 1.0 | 5.76e+07 | 1.07e+03 |
| QCD_HT300to500_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 345900.0 | 1.0 | 5.46e+07 | 2.28e+02 |
| QCD_HT500to700_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 31630.0 | 1.0 | 6.26e+07 | 1.82e+01 |
| QCD_HT700to1000_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 6828.0 | 1.0 | 3.72e+07 | 6.61e+00 |
| ST_t-channel_antitop_4f_inclusiveDecays_13TeV-powhegV2-madspin-pythia8_TuneCUETP8M1/ | 80.95 | 1.0 | 3.88e+07 | 7.51e-02 |
| TT_TuneCUETP8M2T4_13TeV-powheg-pythia8/ | 831.76 | 1.0 | 7.69e+07 | 3.89e-01 |
| ST_s-channel_4f_inclusiveDecays_13TeV-amcatnlo-pythia8/ | 3.36 | 1.0 | 2.95e+07 | 4.10e-03 |
| ST_t-channel_top_4f_inclusiveDecays_13TeV-powhegV2-madspin-pythia8_TuneCUETP8M1/ | 136.02 | 1.0 | 6.71e+07 | 7.30e-02 |
| ST_tW_top_5f_inclusiveDecays_13TeV-powheg-pythia8_TuneCUETP8M1/ | 35.6 | 1.0 | 6.95e+06 | 1.84e-01 |
| ST_tW_antitop_5f_inclusiveDecays_13TeV-powheg-pythia8_TuneCUETP8M1/ | 35.6 | 1.0 | 6.93e+06 | 1.85e-01 |
| WBJetsToLNu_Wpt-100to200_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 5.969 | 1.22 | 3.98e+06 | 6.59e-02 |
| WBJetsToLNu_Wpt-200toInf_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 0.8744 | 1.22 | 5.43e+05 | 7.07e-02 |
| WJetsToLNu_HT-100To200_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 1349.0 | 1.22 | 7.60e+07 | 7.80e-01 |
| WJetsToLNu_HT-1200To2500_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 1.329 | 1.22 | 3.17e+06 | 1.84e-02 |
| WJetsToLNu_HT-200To400_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 359.3 | 1.22 | 3.90e+07 | 4.05e-01 |
| WJetsToLNu_HT-2500ToInf_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 0.03216 | 1.22 | 2.64e+06 | 5.35e-04 |
| WJetsToLNu_HT-400To600_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 48.86 | 1.22 | 6.07e+06 | 3.54e-01 |
| WJetsToLNu_HT-600To800_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 12.03 | 1.22 | 1.87e+07 | 2.83e-02 |
| WJetsToLNu_HT-70To100_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 1352.0 | 1.22 | 3.26e+07 | 1.82e+00 |
| WJetsToLNu_HT-800To1200_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 5.482 | 1.22 | 6.29e+06 | 3.83e-02 |
| WJetsToLNu_BGenFilter_Wpt-100to200_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 26.89 | 1.22 | 4.19e+05 | 2.82e+00 |
| WJetsToLNu_BGenFilter_Wpt-200toInf_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 3.786 | 1.22 | 2.15e+05 | 7.72e-01 |
| WJetsToLNu_TuneCUETP8M1_13TeV-madgraphMLM-pythia8/ | 61526.7 | 1.0 | 8.69e+07 | 2.55e+01 |
| WW_TuneCUETP8M1_13TeV-pythia8/ | 117.6 | 1.0 | 7.98e+06 | 5.30e-01 |
| WWTo1L1Nu2Q_13TeV_amcatnloFXFX_madspin_pythia8/ | 51.87 | 1.0 | 4.41e+08 | 4.23e-03 |
| WZ_TuneCUETP8M1_13TeV-pythia8/ | 48.1 | 1.0 | 4.00e+06 | 4.33e-01 |
| WZTo1L1Nu2Q_13TeV_amcatnloFXFX_madspin_pythia8/ | 10.875 | 1.0 | 1.94e+08 | 2.01e-03 |
| WminusH_HToBB_WToLNu_M125_13TeV_powheg_pythia8/ | 0.10899 | 1.0 | 4.21e+05 | 9.32e-03 |
| WplusH_HToBB_WToLNu_M125_13TeV_powheg_pythia8/ | 0.17202 | 1.0 | 6.90e+05 | 8.97e-03 |
| ZBJetsToNuNu_Zpt-100to200_TuneCUETP8M1_13TeV-madgraphMLM-pythia8 | 5.93 | 1.23 | 6.79e+06 | 3.85e-02 |
| ZBJetsToNuNu_Zpt-200toInf_TuneCUETP8M1_13TeV-madgraphMLM-pythia8 | 0.6075 | 1.23 | 4.44e+05 | 6.06e-02 |
| ZH_HToBB_ZToLL_M125_13TeV_powheg_pythia8/ | 0.04718 | 1.0 | 2.24e+05 | 7.59e-03 |
| ZH_HToBB_ZToNuNu_M125_13TeV_powheg_pythia8/ | 0.09322 | 1.0 | 4.35e+05 | 7.72e-03 |
| ZJetsToNuNu_BGenFilter_Zpt-100to200_TuneCUETP8M1_13TeV-madgraphMLM-pythia8 | 2.162 | 1.23 | 1.50e+05 | 6.37e-01 |
| ZJetsToNuNu_BGenFilter_Zpt-200toInf_TuneCUETP8M1_13TeV-madgraphMLM-pythia8 | 0.3151 | 1.23 | 9.78e+04 | 1.43e-01 |
| ZJetsToNuNu_HT-100To200_13TeV-madgraph | 280.59 | 1.23 | 2.42e+07 | 5.12e-01 |
| ZJetsToNuNu_HT-1200To2500_13TeV-madgraph | 0.28629 | 1.23 | 5.13e+05 | 2.47e-02 |
| ZJetsToNuNu_HT-200To400_13TeV-madgraph | 77.79 | 1.23 | 2.48e+07 | 1.39e-01 |
| ZJetsToNuNu_HT-2500ToInf_13TeV-madgraph | 0.006912 | 1.23 | 4.05e+05 | 7.56e-04 |
| ZJetsToNuNu_HT-400To600_13TeV-madgraph | 10.755 | 1.23 | 8.84e+06 | 5.38e-02 |
| ZJetsToNuNu_HT-600To800_13TeV-madgraph | 2.5644 | 1.23 | 5.77e+06 | 1.97e-02 |
| ZJetsToNuNu_HT-800To1200_13TeV-madgraph | 1.1802 | 1.23 | 2.17e+06 | 2.41e-02 |
| ZZ_TuneCUETP8M1_13TeV-pythia8/ | 17.2 | 1.0 | 1.69e+06 | 3.66e-01 |
| ZZTo2L2Q_13TeV_amcatnloFXFX_madspin_pythia8/ | 2.387 | 1.0 | 7.34e+07 | 1.17e-03 |
| ZZTo2Q2Nu_13TeV_amcatnloFXFX_madspin_pythia8/ | 4.726 | 1.0 | 1.99e+08 | 8.55e-04 |
| ggZH_HToBB_ZToLL_M125_13TeV_powheg_pythia8/ | 0.0072 | 1.0 | 3.00e+06 | 8.65e-05 |
| ggZH_HToBB_ZToNuNu_M125_13TeV_powheg_pythia8/ | 0.01437 | 1.0 | 3.00e+06 | 1.72e-04 |

Table B.1: The MC datasets used for 2016 are listed above.

| Sample Name | Xsec (pb) | KFactor | Total Events | Weight |
|--|-----------|---------|--------------|----------|
| DYBJetsToLL_M-50_Zpt-100to200_TuneCP5_13TeV-madgraphMLM-pythia8_newgridpack/ | 3.224 | 1.23 | 2.72e+06 | 6.09e-02 |
| DYBJetsToLL_M-50_Zpt-200toInf_TuneCP5_13TeV-madgraphMLM-pythia8_newgridpack/ | 0.3298 | 1.23 | 2.40e+05 | 7.05e-02 |
| DYJetsToLL_BGenFilter_Zpt-100to200_M-50_TuneCP5_13TeV-madgraphMLM-pythia8_newgridpack/ | 2.671 | 1.23 | 1.91e+06 | 7.16e-02 |
| DYJetsToLL_BGenFilter_Zpt-200toInf_M-50_TuneCP5_13TeV-madgraphMLM-pythia8_newgridpack/ | 0.3934 | 1.23 | 3.21e+05 | 6.28e-02 |
| DYJetsToLL_M-50_HT-100to200_TuneCP5_13TeV-madgraphMLM-pythia8/ | 161.1 | 1.23 | 1.51e+07 | 5.45e-01 |
| DYJetsToLL_M-50_HT-1200to2500_TuneCP5_13TeV-madgraphMLM-pythia8/ | 0.1933 | 1.23 | 6.25e+05 | 1.59e-02 |
| DYJetsToLL_M-50_HT-200to400_TuneCP5_13TeV-madgraphMLM-pythia8/ | 48.66 | 1.23 | 9.74e+06 | 2.56e-01 |
| DYJetsToLL_M-50_HT-2500toInf_TuneCP5_13TeV-madgraphMLM-pythia8/ | 0.003468 | 1.23 | 4.19e+05 | 4.24e-04 |
| DYJetsToLL_M-50_HT-400to600_TuneCP5_13TeV-madgraphMLM-pythia8/ | 6.968 | 1.23 | 7.05e+06 | 5.07e-02 |
| DYJetsToLL_M-50_HT-600to800_TuneCP5_13TeV-madgraphMLM-pythia8/ | 1.743 | 1.23 | 6.22e+06 | 1.44e-02 |
| DYJetsToLL_M-50_HT-800to1200_TuneCP5_13TeV-madgraphMLM-pythia8/ | 0.8052 | 1.23 | 2.10e+06 | 1.96e-02 |
| DYJetsToLL_M-4to50_HT-100to200_TuneCP5_13TeV-madgraphMLM-pythia8/ | 204.0 | 1.23 | 9.88e+06 | 1.06e+00 |
| DYJetsToLL_M-4to50_HT-200to400_TuneCP5_13TeV-madgraphMLM-pythia8/ | 54.39 | 1.23 | 4.02e+06 | 6.94e-01 |
| DYJetsToLL_M-4to50_HT-400to600_TuneCP5_13TeV-madgraphMLM-pythia8/ | 5.697 | 1.23 | 2.98e+06 | 9.81e-02 |
| DYJetsToLL_M-4to50_HT-600toInf_TuneCP5_13TeV-madgraphMLM-pythia8/ | 1.85 | 1.23 | 2.93e+06 | 3.24e-02 |
| DYJetsToLL_M-50_TuneCP5_13TeV-madgraphMLM-pythia8/ | 5343.0 | 1.23 | 9.70e+07 | 2.83e+00 |
| QCD_HT1000to1500_TuneCP5_13TeV-madgraph-pythia8/ | 1088.0 | 1.0 | 1.66e+07 | 2.73e+00 |
| QCD_HT1500to2000_TuneCP5_13TeV-madgraph-pythia8/ | 99.11 | 1.0 | 1.16e+07 | 3.55e-01 |
| QCD_HT2000toInf_TuneCP5_13TeV-madgraph-pythia8/ | 20.23 | 1.0 | 5.94e+06 | 1.42e-01 |
| QCD_HT200to300_TuneCP5_13TeV-madgraph-pythia8/ | 1547000.0 | 1.0 | 5.94e+07 | 1.09e+03 |
| QCD_HT300to500_TuneCP5_13TeV-madgraph-pythia8/ | 322600.0 | 1.0 | 6.02e+07 | 2.23e+02 |
| QCD_HT500to700_TuneCP5_13TeV-madgraph-pythia8/ | 29980.0 | 1.0 | 5.62e+07 | 2.22e+01 |
| QCD_HT700to1000_TuneCP5_13TeV-madgraph-pythia8/ | 6334.0 | 1.0 | 7.53e+07 | 3.51e+00 |
| ST_s-channel_4f_leptonDecays_TuneCP5_PSweights_13TeV-amcatnlo-pythia8/ | 3.74 | 1.0 | 9.91e+06 | 1.57e-02 |
| ST_t-channel_antitop_4f_InclusiveDecays_TuneCP5_PSweights_13TeV-powheg-pythia8/ | 80.95 | 1.0 | 6.26e+07 | 5.39e-02 |
| ST_t-channel_top_4f_InclusiveDecays_TuneCP5_PSweights_13TeV-powheg-pythia8/ | 136.02 | 1.0 | 1.75e+08 | 3.23e-02 |
| ST_tW_antitop_5f_inclusiveDecays_TuneCP5_PSweights_13TeV-powheg-pythia8/ | 35.85 | 1.0 | 7.75e+06 | 1.93e-01 |
| ST_tW_top_5f_inclusiveDecays_TuneCP5_PSweights_13TeV-powheg-pythia8/ | 35.85 | 1.0 | 7.95e+06 | 1.88e-01 |
| TTToHadronic_TuneCP5_PSweights_13TeV-powheg-pythia8/ | 377.96 | 1.0 | 1.30e+08 | 1.21e-01 |
| TTTo2L2Nu_TuneCP5_PSweights_13TeV-powheg-pythia8/ | 88.29 | 1.0 | 3.78e+07 | 9.73e-02 |
| TTToSemiLeptonic_TuneCP5_PSweights_13TeV-powheg-pythia8/ | 365.34 | 1.0 | 8.41e+07 | 1.81e-01 |
| WBJetsToLNu_Wpt-100to200_TuneCP5_13TeV-madgraphMLM-pythia8_newgridpack/ | 5.542 | 1.21 | 4.15e+06 | 6.74e-02 |
| WBJetsToLNu_Wpt-200toInf_TuneCP5_13TeV-madgraphMLM-pythia8_newgridpack/ | 0.801 | 1.21 | 7.72e+05 | 5.24e-02 |
| WJetsToLNu_HT-100To200_TuneCP5_13TeV-madgraphMLM-pythia8/ | 1395.0 | 1.21 | 3.45e+07 | 2.04e+00 |
| WJetsToLNu_HT-1200To2500_TuneCP5_13TeV-madgraphMLM-pythia8/ | 1.074 | 1.21 | 1.24e+07 | 4.36e-03 |
| WJetsToLNu_HT-200To400_TuneCP5_13TeV-madgraphMLM-pythia8/ | 407.9 | 1.21 | 2.13e+07 | 9.69e-01 |
| WJetsToLNu_HT-2500ToInf_TuneCP5_13TeV-madgraphMLM-pythia8/ | 0.03216 | 1.21 | 1.28e+07 | 1.26e-04 |
| WJetsToLNu_HT-400To600_TuneCP5_13TeV-madgraphMLM-pythia8/ | 57.48 | 1.21 | 5.63e+06 | 5.15e-01 |
| WJetsToLNu_HT-600To800_TuneCP5_13TeV-madgraphMLM-pythia8/ | 12.87 | 1.21 | 1.88e+07 | 3.45e-02 |
| WJetsToLNu_HT-800To1200_TuneCP5_13TeV-madgraphMLM-pythia8/ | 5.366 | 1.21 | 8.15e+06 | 3.32e-02 |
| WJetsToLNu_BGenFilter_Wpt-100toInf_TuneCP5_13TeV-madgraphMLM-pythia8_newgridpack/ | 20.56 | 1.21 | 1.34e+07 | 7.76e-02 |
| WJetsToLNu_BGenFilter_Wpt-200toInf_TuneCP5_13TeV-madgraphMLM-pythia8_newgridpack/ | 2.936 | 1.21 | 2.48e+06 | 5.96e-02 |
| WJetsToLNu_TuneCP5_13TeV-madgraphMLM-pythia8/ | 52940.0 | 1.0 | 7.77e+07 | 2.84e+01 |
| WW_TuneCP5_13TeV-pythia8/ | 115.3 | 1.0 | 7.71e+06 | 6.24e-01 |
| WWTo1L1Nu2Q_13TeV_amcatnloFXFX_madspin_pythia8/ | 50.858 | 1.0 | 5.05e+06 | 4.20e-01 |
| WWToLNuQQ_NNPDF31_TuneCP5_PSweights_13TeV-powheg-pythia8/ | 50.858 | 1.0 | 8.79e+06 | 2.41e-01 |
| WZ_TuneCP5_13TeV-pythia8/ | 48.1 | 1.0 | 3.93e+06 | 5.11e-01 |
| WminusH_HTtoBB_WToLNu_M125_13TeV_powheg_pythia8/ | 0.10899 | 1.0 | 2.38e+06 | 1.91e-03 |
| WplusH_HTtoBB_WToLNu_M125_13TeV_powheg_pythia8/ | 0.17202 | 1.0 | 4.84e+06 | 1.48e-03 |
| ZBJetsToNuNu_Zpt-100to200_TuneCP5_13TeV-madgraphMLM-pythia8_newgridpack/ | 6.209 | 1.23 | 4.98e+06 | 6.40e-02 |
| ZBJetsToNuNu_Zpt-200toInf_TuneCP5_13TeV-madgraphMLM-pythia8_newgridpack/ | 0.6286 | 1.23 | 5.14e+05 | 6.27e-02 |
| ZH_HTtoBB_ZToNuNu_M125_13TeV_powheg_pythia8/ | 0.09322 | 1.0 | 2.58e+06 | 1.51e-03 |
| ZJetsToNuNu_BGenFilter_Zpt-100to200_TuneCP5_13TeV-madgraphMLM-pythia8_newgridpack/ | 1.689 | 1.23 | 1.37e+06 | 6.31e-02 |
| ZJetsToNuNu_BGenFilter_Zpt-200toInf_TuneCP5_13TeV-madgraphMLM-pythia8_newgridpack/ | 0.2476 | 1.23 | 1.89e+05 | 6.73e-02 |
| ZJetsToNuNu_HT-100To200_13TeV-madgraph/ | 302.8 | 1.23 | 2.27e+07 | 6.83e-01 |
| ZJetsToNuNu_HT-200To400_13TeV-madgraph/ | 92.59 | 1.23 | 2.17e+07 | 2.19e-01 |
| ZJetsToNuNu_HT-2500ToInf_13TeV-madgraph/ | 0.005146 | 1.23 | 6.73e+03 | 3.92e-02 |
| ZJetsToNuNu_HT-400To600_13TeV-madgraph/ | 13.18 | 1.23 | 1.83e+07 | 3.70e-02 |
| ZJetsToNuNu_HT-600To800_13TeV-madgraph/ | 3.257 | 1.23 | 1.13e+07 | 1.47e-02 |
| ZJetsToNuNu_HT-800To1200_13TeV-madgraph/ | 1.496 | 1.23 | 2.06e+06 | 3.73e-02 |
| ZZ_TuneCP5_13TeV-pythia8/ | 14.6 | 1.0 | 1.95e+06 | 3.12e-01 |
| ZZTo2L2Q_13TeV_amcatnloFXFX_madspin_pythia8/ | 10.88 | 1.0 | 2.45e+07 | 1.85e-02 |
| ZZTo4L_13TeV_powheg_pythia8/ | 2.038 | 1.0 | 2.50e+07 | 3.40e-03 |
| ggZH_HTtoBB_ZToLL_M125_13TeV_powheg_pythia8/ | 0.0072 | 1.0 | 2.85e+06 | 1.05e-04 |
| ggZH_HTtoBB_ZToNuNu_M125_13TeV_powheg_pythia8/ | 0.01437 | 1.0 | 2.98e+06 | 2.01e-04 |

Table B.2: The MC datasets used for 2017 are listed above.

| Sample Name | Xsec (pb) | KFactor | Total Events | Weight |
|---|-----------|---------|--------------|----------|
| DYBJetsToLL_M-50_Zpt-100to200_TuneCP5_13TeV-madgraphMLM-pythia8/ | 3.206 | 1.23 | 1.99e+07 | 1.19e-02 |
| DYBJetsToLL_M-50_Zpt-200toInf_TuneCP5_13TeV-madgraphMLM-pythia8/ | 0.3304 | 1.23 | 1.75e+06 | 1.40e-02 |
| DYJetsToLL_BGenFilter_Zpt-100to200_M-50_TuneCP5_13TeV-madgraphMLM-pythia8/ | 2.662 | 1.23 | 6.54e+05 | 3.01e-01 |
| DYJetsToLL_BGenFilter_Zpt-200toInf_M-50_TuneCP5_13TeV-madgraphMLM-pythia8/ | 0.3949 | 1.23 | 1.03e+05 | 2.83e-01 |
| DYJetsToLL_M-50_HT-100to200_TuneCP5_PSweights_13TeV-madgraphMLM-pythia8/ | 160.8 | 1.23 | 1.15e+07 | 1.03e+00 |
| DYJetsToLL_M-50_HT-1200to2500_TuneCP5_PSweights_13TeV-madgraphMLM-pythia8/ | 0.1931 | 1.23 | 5.32e+05 | 2.68e-02 |
| DYJetsToLL_M-50_HT-200to400_TuneCP5_PSweights_13TeV-madgraphMLM-pythia8/ | 48.63 | 1.23 | 1.07e+07 | 3.35e-01 |
| DYJetsToLL_M-50_HT-2500toInf_TuneCP5_PSweights_13TeV-madgraphMLM-pythia8/ | 0.003513 | 1.23 | 4.16e+05 | 6.24e-04 |
| DYJetsToLL_M-50_HT-400to600_TuneCP5_PSweights_13TeV-madgraphMLM-pythia8/ | 6.982 | 1.23 | 3.44e+07 | 1.50e-02 |
| DYJetsToLL_M-50_HT-600to800_TuneCP5_PSweights_13TeV-madgraphMLM-pythia8/ | 1.756 | 1.23 | 8.83e+06 | 1.47e-02 |
| DYJetsToLL_M-50_HT-800to1200_TuneCP5_PSweights_13TeV-madgraphMLM-pythia8/ | 0.8094 | 1.23 | 3.12e+06 | 1.91e-02 |
| DYJetsToLL_M-50_TuneCP5_13TeV-madgraphMLM-pythia8/ | 5343.0 | 1.23 | 9.78e+07 | 4.03e+00 |
| QCD_HT1000to1500_TuneCP5_13TeV-madgraphMLM-pythia8/ | 1088.0 | 1.0 | 1.54e+07 | 4.24e+00 |
| QCD_HT1500to2000_TuneCP5_13TeV-madgraphMLM-pythia8/ | 99.11 | 1.0 | 1.01e+07 | 5.87e-01 |
| QCD_HT2000toInf_TuneCP5_13TeV-madgraphMLM-pythia8/ | 20.23 | 1.0 | 5.41e+06 | 2.24e-01 |
| QCD_HT200to300_TuneCP5_13TeV-madgraphMLM-pythia8/ | 1547000.0 | 1.0 | 5.42e+07 | 1.71e+03 |
| QCD_HT300to500_TuneCP5_13TeV-madgraphMLM-pythia8/ | 322600.0 | 1.0 | 5.46e+07 | 3.55e+02 |
| QCD_HT500to700_TuneCP5_13TeV-madgraphMLM-pythia8/ | 29980.0 | 1.0 | 5.50e+07 | 3.27e+01 |
| QCD_HT700to1000_TuneCP5_13TeV-madgraphMLM-pythia8/ | 6334.0 | 1.0 | 4.47e+07 | 8.50e+00 |
| ST_s-channel_4f_leptonDecays_TuneCP5_13TeV-madgraph-pythia8/ | 3.74 | 1.0 | 1.48e+08 | 1.51e-03 |
| ST_t-channel_antitop_4f_InclusiveDecays_TuneCP5_13TeV-powheg-madspin-pythia8/ | 80.95 | 1.0 | 5.13e+09 | 9.48e-04 |
| ST_t-channel_top_4f_InclusiveDecays_TuneCP5_13TeV-powheg-madspin-pythia8/ | 136.02 | 1.0 | 1.66e+10 | 4.92e-04 |
| ST_tW_antitop_5f_inclusiveDecays_TuneCP5_13TeV-powheg-pythia8/ | 35.85 | 1.0 | 2.66e+08 | 8.07e-03 |
| ST_tW_antitop_5f_NoFullyHadronicDecays_TuneCP5_13TeV-powheg-pythia8/ | 19.56 | 1.0 | 2.42e+08 | 4.86e-03 |
| ST_tW_top_5f_inclusiveDecays_TuneCP5_13TeV-powheg-pythia8/ | 35.85 | 1.0 | 3.35e+08 | 6.42e-03 |
| ST_tW_top_5f_NoFullyHadronicDecays_TuneCP5_13TeV-powheg-pythia8/ | 19.56 | 1.0 | 3.04e+08 | 3.86e-03 |
| TTToHadronic_TuneCP5_13TeV-powheg-pythia8/ | 377.96 | 1.0 | 1.03e+11 | 2.20e-04 |
| TTTo2L2Nu_TuneCP5_13TeV-powheg-pythia8/ | 88.29 | 1.0 | 4.50e+09 | 1.18e-03 |
| TTToSemiLeptonic_TuneCP5_13TeV-powheg-pythia8/ | 365.34 | 1.0 | 5.38e+10 | 4.07e-04 |
| WBJetsToLNu_Wpt-100to200_TuneCP5_13TeV-madgraphMLM-pythia8/ | 5.527 | 1.21 | 7.36e+06 | 5.45e-02 |
| WBJetsToLNu_Wpt-200toInf_TuneCP5_13TeV-madgraphMLM-pythia8/ | 0.7996 | 1.21 | 3.02e+05 | 1.92e-01 |
| WJetsToLNu_HT-100To200_TuneCP5_13TeV-madgraphMLM-pythia8/ | 1392.0 | 1.21 | 2.83e+07 | 3.57e+00 |
| WJetsToLNu_HT-1200To2500_TuneCP5_13TeV-madgraphMLM-pythia8/ | 1.084 | 1.21 | 7.40e+06 | 1.06e-02 |
| WJetsToLNu_HT-200To400_TuneCP5_13TeV-madgraphMLM-pythia8/ | 410.3 | 1.21 | 2.22e+07 | 1.34e+00 |
| WJetsToLNu_HT-2500ToInf_TuneCP5_13TeV-madgraphMLM-pythia8/ | 0.008067 | 1.21 | 3.19e+06 | 1.84e-04 |
| WJetsToLNu_HT-400To600_TuneCP5_13TeV-madgraphMLM-pythia8/ | 57.85 | 1.21 | 5.91e+06 | 7.10e-01 |
| WJetsToLNu_HT-600To800_TuneCP5_13TeV-madgraphMLM-pythia8/ | 12.95 | 1.21 | 1.68e+07 | 5.61e-02 |
| WJetsToLNu_HT-800To1200_TuneCP5_13TeV-madgraphMLM-pythia8/ | 5.45 | 1.21 | 8.36e+06 | 4.73e-02 |
| WJetsToLNu_BGenFilter_Wpt-100to200_TuneCP5_13TeV-madgraphMLM-pythia8/ | 20.49 | 1.21 | 2.36e+05 | 6.30e+00 |
| WJetsToLNu_BGenFilter_Wpt-200toInf_TuneCP5_13TeV-madgraphMLM-pythia8/ | 2.935 | 1.21 | 8.30e+05 | 2.57e-01 |
| WJetsToLNu_HT-70To100_TuneCP5_13TeV-madgraphMLM-pythia8/ | 1353.0 | 1.21 | 7.04e+07 | 1.40e+00 |
| WminusH_HTtoBB_WToLNu_M125_13TeV_powheg_pythia8/ | 0.10899 | 1.0 | 6.40e+05 | 1.02e-02 |
| WplusH_HTtoBB_WToLNu_M125_13TeV_powheg_pythia8/ | 0.17202 | 1.0 | 1.01e+06 | 1.02e-02 |
| ZBJetsToNuNu_Zpt-100to200_TuneCP5_13TeV-madgraphMLM-pythia8 | 6.195 | 1.23 | 6.65e+07 | 6.87e-03 |
| ZBJetsToNuNu_Zpt-200toInf_TuneCP5_13TeV-madgraphMLM-pythia8 | 0.6293 | 1.23 | 7.35e+06 | 6.32e-03 |
| ZH_HTtoBB_ZToLL_M125_13TeV_powheg_pythia8/ | 0.04718 | 1.0 | 5.53e+05 | 5.12e-03 |
| ZH_HTtoBB_ZToNuNu_M125_13TeV_powheg_pythia8/ | 0.09322 | 1.0 | 7.15e+05 | 7.82e-03 |
| ZJetsToNuNu_BGenFilter_Zpt-100to200_TuneCP5_13TeV-madgraphMLM-pythia8 | 1.679 | 1.23 | 2.01e+05 | 6.16e-01 |
| ZJetsToNuNu_BGenFilter_Zpt-200toInf_TuneCP5_13TeV-madgraphMLM-pythia8 | 0.2468 | 1.23 | 3.73e+04 | 4.88e-01 |
| ZJetsToNuNu_HT-100To200_13TeV-madgraph | 303.4 | 1.23 | 2.37e+07 | 9.46e-01 |
| ZJetsToNuNu_HT-1200To2500_13TeV-madgraph | 0.3425 | 1.23 | 3.40e+05 | 7.43e-02 |
| ZJetsToNuNu_HT-200To400_13TeV-madgraph | 91.71 | 1.23 | 2.32e+07 | 2.91e-01 |
| ZJetsToNuNu_HT-2500ToInf_13TeV-madgraph | 0.005263 | 1.23 | 3.50e+05 | 1.11e-03 |
| ZJetsToNuNu_HT-400To600_13TeV-madgraph | 13.1 | 1.23 | 9.48e+06 | 1.02e-01 |
| ZJetsToNuNu_HT-600To800_13TeV-madgraph | 3.248 | 1.23 | 5.73e+06 | 4.19e-02 |
| ZJetsToNuNu_HT-800To1200_13TeV-madgraph | 1.496 | 1.23 | 2.06e+06 | 5.37e-02 |
| ggZH_HTtoBB_ZToLL_M125_13TeV_powheg_pythia8/ | 0.01437 | 1.0 | 2.70e+04 | 3.20e-02 |
| ggZH_HTtoBB_ZToNuNu_M125_13TeV_powheg_pythia8/ | 0.01437 | 1.0 | 5.62e+04 | 1.53e-02 |

Table B.3: The MC datasets used for 2018 are listed above.

Appendix C

Data Card

The tables in this appendix give the observed yields and profit predictions for each channel and each year.

Table C.1: The observed and predicted yields are given for the signal regions for 0-lepton in 2016.

| Process | Medium $p_T(V)$ 0J | Medium $p_T(V)$ GE1J | High $p_T(V)$ | High $p_T(V)$, boosted | Highest $p_T(V)$ | Highest $p_T(V)$, boosted |
|-------------------------|-----------------------|----------------------------|------------------|-------------------------------|---------------------|----------------------------------|
| Data | 3444 | 4036 | 1601 | 2035 | 152 | 423 |
| WH 150_250 0J | 7.297e+00 | 1.392e+00 | 4.894e-01 | 0 | 0 | 0 |
| WH 150_250 GE1J | 7.867e-01 | 1.892e+00 | 2.302e-01 | 3.321e-02 | 0 | 0 |
| WH 250_400 | 1.419e+00 | 9.698e-01 | 3.503e+00 | 7.489e-01 | 4.932e-02 | 1.924e-02 |
| WH 75_150 | 6.560e-01 | 1.731e-01 | 0 | 0 | 0 | 0 |
| WH GT400 | 2.859e-02 | 8.905e-03 | 3.160e-01 | 1.040e-01 | 3.322e-01 | 3.394e-01 |
| ZH 0_75 | 0 | 0 | 2.550e-03 | 0 | 0 | 0 |
| ZH 150_250 0J | 2.299e+01 | 2.596e+00 | 2.481e+00 | 2.669e-02 | 0 | 0 |
| ZH 150_250 GE1J | 2.809e+00 | 5.882e+00 | 1.016e+00 | 1.134e-01 | 0 | 0 |
| ZH 250_400 | 3.374e+00 | 1.271e+00 | 1.613e+01 | 3.322e+00 | 5.961e-01 | 2.165e-01 |
| ZH 75_150 | 1.814e+00 | 4.654e-01 | 6.357e-03 | 0 | 0 | 0 |
| ZH GT400 | 7.561e-03 | 6.880e-03 | 1.040e+00 | 4.143e-01 | 3.322e+00 | 2.615e+00 |
| ggZH 0_75 | 2.600e-04 | 0 | 0 | 0 | 0 | 0 |
| ggZH 150_250 0J | 5.120e+00 | 6.577e-01 | 4.522e-01 | 7.052e-03 | 0 | 0 |
| ggZH 150_250 GE1J | 2.206e+00 | 3.975e+00 | 6.435e-01 | 8.790e-02 | 0 | 0 |
| ggZH 250_400 | 6.495e-01 | 4.919e-01 | 3.457e+00 | 8.091e-01 | 8.446e-02 | 4.281e-02 |
| ggZH 75_150 | 5.832e-01 | 3.063e-01 | 1.357e-03 | 0 | 0 | 0 |
| ggZH GT400 | 7.085e-04 | 4.001e-04 | 9.972e-02 | 6.591e-02 | 3.183e-01 | 2.834e-01 |
| TT | 1.530e+03 | 2.557e+03 | 6.268e+02 | 7.935e+02 | 1.888e+01 | 9.874e+01 |
| VVHF | 8.001e+01 | 4.534e+01 | 5.194e+01 | 1.625e+01 | 6.761e+00 | 6.037e+00 |
| VVLF | 2.920e+01 | 2.805e+01 | 2.264e+01 | 4.700e+01 | 3.771e+00 | 1.411e+01 |
| Wj0b_c | 8.414e+01 | 7.524e+01 | 5.032e+01 | 7.504e+01 | 4.481e+00 | 1.932e+01 |
| Wj0b_udsg | 1.088e+02 | 9.527e+01 | 6.117e+01 | 1.265e+02 | 6.757e+00 | 2.355e+01 |
| Wj1b | 7.093e+01 | 5.827e+01 | 3.482e+01 | 5.805e+01 | 3.440e+00 | 1.271e+01 |
| Wj2b | 1.423e+02 | 1.451e+02 | 9.751e+01 | 1.211e+02 | 1.099e+01 | 3.260e+01 |
| Zj0b_c | 1.314e+02 | 7.419e+01 | 7.680e+01 | 1.128e+02 | 9.821e+00 | 4.681e+01 |
| Zj0b_udsg | 1.027e+02 | 6.430e+01 | 6.435e+01 | 1.522e+02 | 1.174e+01 | 6.052e+01 |
| Zj1b | 1.607e+02 | 9.877e+01 | 7.365e+01 | 1.124e+02 | 8.575e+00 | 3.373e+01 |
| Zj2b | 3.791e+02 | 2.391e+02 | 1.861e+02 | 1.273e+02 | 2.537e+01 | 6.232e+01 |
| s_Top | 1.867e+02 | 1.846e+02 | 6.419e+01 | 7.596e+01 | 3.774e+00 | 1.171e+01 |

Table C.2: The observed and predicted yields are given for the control regions for 0-lepton in 2016.

| Process | Z+udcsg enr., med. $p_T(V)$ | Z+udcsg enr., high $p_T(V)$ | Z+udcsg enr., high $p_T(V)$, boosted |
|-----------|--------------------------------|--------------------------------|---|
| Data | 9160 | 2742 | 78659 |
| TT | 1.367e+03 | 3.053e+02 | 3.191e+03 |
| VVHF | 1.133e+01 | 5.163e+00 | 1.883e+01 |
| VVLF | 1.446e+02 | 6.736e+01 | 1.889e+03 |
| Wj0b_c | 7.463e+02 | 2.018e+02 | 2.943e+03 |
| Wj0b_udsg | 2.264e+03 | 5.809e+02 | 2.496e+04 |
| Wj1b | 9.205e+01 | 2.615e+01 | 3.510e+02 |
| Wj2b | 5.640e+01 | 1.715e+01 | 1.879e+02 |
| Zj0b_c | 7.793e+02 | 2.818e+02 | 4.405e+03 |
| Zj0b_udsg | 1.770e+03 | 7.512e+02 | 3.749e+04 |
| Zj1b | 1.575e+02 | 5.061e+01 | 6.923e+02 |
| Zj2b | 8.495e+01 | 3.031e+01 | 2.533e+02 |
| s_Top | 1.699e+02 | 3.858e+01 | 5.277e+02 |

| Process | Z+bb enr., med. $p_T(V)$ | Z+bb enr., high $p_T(V)$ | Z+bb enr., high $p_T(V)$, boosted |
|-----------|-----------------------------|-----------------------------|---------------------------------------|
| Data | 13062 | 2952 | 2567 |
| TT | 5.855e+03 | 1.058e+03 | 8.015e+02 |
| VVHF | 9.473e+01 | 3.227e+01 | 1.869e+01 |
| VVLF | 7.388e+01 | 2.008e+01 | 8.989e+01 |
| Wj0b_c | 4.188e+02 | 1.023e+02 | 1.498e+02 |
| Wj0b_udsg | 5.343e+02 | 1.285e+02 | 2.215e+02 |
| Wj1b | 3.060e+02 | 7.521e+01 | 9.754e+01 |
| Wj2b | 6.184e+02 | 1.710e+02 | 2.421e+02 |
| Zj0b_c | 4.219e+02 | 1.581e+02 | 2.285e+02 |
| Zj0b_udsg | 3.478e+02 | 1.481e+02 | 3.006e+02 |
| Zj1b | 5.748e+02 | 1.592e+02 | 2.419e+02 |
| Zj2b | 1.294e+03 | 3.618e+02 | 3.005e+02 |
| s_Top | 6.552e+02 | 1.414e+02 | 9.027e+01 |

| Process | $t\bar{t}$ enr., med. $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$, boosted |
|-----------|-----------------------------------|-----------------------------------|---|
| Data | 3395 | 838 | 1229 |
| TT | 2.846e+03 | 6.922e+02 | 1.108e+03 |
| VVHF | 3.394e+00 | 1.482e+00 | 5.775e+00 |
| VVLF | 8.099e+00 | 1.330e+00 | 8.263e+00 |
| Wj0b_c | 3.301e+01 | 1.050e+01 | 1.648e+01 |
| Wj0b_udsg | 3.762e+01 | 1.403e+01 | 1.502e+01 |
| Wj1b | 2.378e+01 | 7.778e+00 | 1.714e+01 |
| Wj2b | 3.117e+01 | 1.160e+01 | 4.850e+01 |
| Zj0b_c | 2.130e+01 | 8.779e+00 | 1.491e+01 |
| Zj0b_udsg | 1.805e+01 | 8.977e+00 | 9.209e+00 |
| Zj1b | 2.265e+01 | 8.555e+00 | 1.992e+01 |
| Zj2b | 4.769e+01 | 2.146e+01 | 7.903e+01 |
| s_Top | 1.281e+02 | 3.589e+01 | 7.903e+01 |

Table C.3: The observed and predicted yields are given for the signal regions for 1-lepton (e) in 2016.

| Process | Medium $p_T(V)$ | High $p_T(V)$ | High $p_T(V)$, boosted | Highest $p_T(V)$ | Highest $p_T(V)$, boosted |
|--------------------|--------------------|---------------|----------------------------|---------------------|----------------------------------|
| Data | 12006 | 1563 | 1921 | 129 | 423 |
| WH 150_250 0J | 2.357e+01 | 1.537e+00 | 1.458e-02 | 0 | 0 |
| WH 150_250 GE1J | 8.049e+00 | 5.998e-01 | 1.080e-01 | 0 | 0 |
| WH 250_400 | 3.431e+00 | 1.148e+01 | 2.492e+00 | 4.427e-01 | 1.747e-01 |
| WH 75_150 | 6.580e+00 | 1.243e-05 | 0 | 0 | 0 |
| WH GT400 | 7.395e-03 | 8.824e-01 | 4.518e-01 | 2.574e+00 | 2.214e+00 |
| ZH 0_75 | 9.896e-04 | 0 | 0 | 0 | 0 |
| ZH 150_250 0J | 6.733e-01 | 2.646e-02 | 0 | 0 | 0 |
| ZH 150_250 GE1J | 2.275e-01 | 1.549e-02 | 0 | 0 | 0 |
| ZH 250_400 | 1.887e-01 | 1.845e-01 | 3.590e-02 | 2.067e-03 | 6.941e-04 |
| ZH 75_150 | 2.842e-01 | 0 | 0 | 0 | 0 |
| ZH GT400 | 1.640e-02 | 3.799e-02 | 2.742e-02 | 2.018e-02 | 2.326e-02 |
| TT | 9.417e+03 | 1.103e+03 | 1.440e+03 | 4.114e+01 | 2.545e+02 |
| VVHF | 6.272e+01 | 1.928e+01 | 6.898e+00 | 3.129e+00 | 2.331e+00 |
| VVLF | 4.461e+01 | 1.783e+01 | 3.013e+01 | 9.757e-01 | 1.840e+01 |
| Wj0b_c | 4.404e+02 | 8.025e+01 | 1.247e+02 | 1.365e+01 | 4.264e+01 |
| Wj0b_udsg | 3.764e+02 | 7.213e+01 | 1.116e+02 | 1.416e+01 | 4.817e+01 |
| Wj1b | 2.422e+02 | 5.471e+01 | 6.768e+01 | 9.447e+00 | 2.286e+01 |
| Wj2b | 3.780e+02 | 1.223e+02 | 1.553e+02 | 2.464e+01 | 7.352e+01 |
| Zj0b_c | 2.214e+01 | 1.496e+00 | 3.790e+00 | 1.217e-01 | 1.344e+00 |
| Zj0b_udsg | 3.634e+01 | 2.586e+00 | 5.240e+00 | 3.025e-01 | 1.067e+00 |
| Zj1b | 4.432e+01 | 3.186e+00 | 3.964e+00 | 1.562e-01 | 6.485e-01 |
| Zj2b | 4.838e+01 | 5.169e+00 | 5.912e+00 | 4.418e-01 | 1.759e+00 |
| s_Top | 1.373e+03 | 2.444e+02 | 2.081e+02 | 1.871e+01 | 7.334e+01 |

Table C.4: The observed and predicted yields are given for the control regions for 1-lepton (e) in 2016.

| Process | W+udcsg enr., med. $p_T(V)$ | W+udcsg enr., high $p_T(V)$ | W+udcsg enr., high $p_T(V)$, boosted |
|-----------|-----------------------------------|-----------------------------------|---|
| Data | 105027 | 24155 | 39175 |
| TT | 1.924e+04 | 4.547e+03 | 5.435e+03 |
| VVHF | 2.438e+01 | 8.982e+00 | 1.229e+01 |
| VVLF | 2.289e+03 | 9.025e+02 | 1.819e+03 |
| Wj0b_c | 1.579e+04 | 3.539e+03 | 4.458e+03 |
| Wj0b_udsg | 5.239e+04 | 1.313e+04 | 2.742e+04 |
| Wj1b | 1.134e+03 | 3.139e+02 | 4.068e+02 |
| Wj2b | 3.572e+02 | 1.527e+02 | 2.367e+02 |
| Zj0b_c | 5.605e+02 | 7.261e+01 | 1.302e+02 |
| Zj0b_udsg | 3.489e+03 | 4.150e+02 | 1.050e+03 |
| Zj1b | 1.292e+02 | 1.385e+01 | 2.376e+01 |
| Zj2b | 2.497e+01 | 4.251e+00 | 8.959e+00 |
| s_Top | 3.104e+03 | 8.494e+02 | 1.057e+03 |
| Process | W+bb enr., med. $p_T(V)$ | W+bb enr., high $p_T(V)$ | W+bb enr., high $p_T(V)$, boosted |
| Data | 20605 | 3470 | 2024 |
| TT | 1.545e+04 | 2.473e+03 | 1.547e+03 |
| VVHF | 5.574e+01 | 1.951e+01 | 6.503e+00 |
| VVLF | 5.773e+01 | 2.360e+01 | 8.361e+01 |
| Wj0b_c | 8.084e+02 | 1.778e+02 | 2.028e+02 |
| Wj0b_udsg | 7.505e+02 | 1.958e+02 | 1.985e+02 |
| Wj1b | 5.339e+02 | 1.292e+02 | 1.141e+02 |
| Wj2b | 8.627e+02 | 2.425e+02 | 3.134e+02 |
| Zj0b_c | 4.896e+01 | 5.681e+00 | 7.085e+00 |
| Zj0b_udsg | 6.818e+01 | 6.542e+00 | 9.776e+00 |
| Zj1b | 7.259e+01 | 6.353e+00 | 6.781e+00 |
| Zj2b | 7.790e+01 | 7.958e+00 | 1.088e+01 |
| s_Top | 2.190e+03 | 5.302e+02 | 2.807e+02 |
| Process | $t\bar{t}$ enr., med. $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$, boosted |
| Data | 38193 | 9459 | 3450 |
| TT | 3.708e+04 | 8.817e+03 | 4.079e+03 |
| VVHF | 6.869e+01 | 4.074e+01 | 4.040e+00 |
| VVLF | 5.154e+01 | 3.035e+01 | 1.061e+01 |
| Wj0b_c | 2.374e+02 | 9.584e+01 | 1.743e+01 |
| Wj0b_udsg | 2.763e+02 | 9.764e+01 | 6.360e+00 |
| Wj1b | 6.486e+02 | 2.500e+02 | 1.625e+01 |
| Wj2b | 8.032e+02 | 3.519e+02 | 5.856e+01 |
| Zj0b_c | 1.436e+01 | 3.906e+00 | 1.312e+00 |
| Zj0b_udsg | 2.280e+01 | 4.638e+00 | 9.908e-01 |
| Zj1b | 7.477e+01 | 1.424e+01 | 9.796e-01 |
| Zj2b | 7.816e+01 | 1.694e+01 | 3.997e+00 |
| s_Top | 2.540e+03 | 1.054e+03 | 3.411e+02 |

Table C.5: The observed and predicted yields are given for the signal regions for 1-lepton (μ) in 2016.

| Process | Medium $p_T(V)$ | High $p_T(V)$ | High $p_T(V)$, boosted | Highest $p_T(V)$ | Highest $p_T(V)$, boosted |
|------------|--------------------|---------------|----------------------------|---------------------|----------------------------------|
| Data | 16249 | 2127 | 2463 | 154 | 581 |
| WH 0_75 | 1.951e-02 | 0 | 0 | 0 | 0 |
| WH 150_250 | 2.983e+01 | 1.981e+00 | 5.517e-03 | 0 | 0 |
| 0J | | | | | |
| WH 150_250 | 1.037e+01 | 8.396e-01 | 1.194e-01 | 0 | 0 |
| GE1J | | | | | |
| WH 250_400 | 4.128e+00 | 1.381e+01 | 2.839e+00 | 5.374e-01 | 2.125e-01 |
| WH 75_150 | 8.437e+00 | 2.732e-04 | 0 | 0 | 0 |
| WH GT400 | 9.994e-03 | 8.083e-01 | 4.759e-01 | 2.982e+00 | 2.604e+00 |
| ZH 150_250 | 9.829e-01 | 4.906e-02 | 8.978e-04 | 0 | 0 |
| 0J | | | | | |
| ZH 150_250 | 3.053e-01 | 1.697e-02 | 0 | 0 | 0 |
| GE1J | | | | | |
| ZH 250_400 | 1.964e-01 | 2.314e-01 | 6.373e-02 | 4.133e-03 | 1.195e-03 |
| ZH 75_150 | 4.484e-01 | 1.194e-03 | 0 | 0 | 0 |
| ZH GT400 | 1.158e-02 | 2.205e-02 | 2.343e-02 | 2.572e-02 | 2.038e-02 |
| TT | 1.227e+04 | 1.362e+03 | 1.762e+03 | 5.458e+01 | 2.972e+02 |
| VVHF | 8.050e+01 | 2.528e+01 | 5.329e+00 | 3.062e+00 | 2.303e+00 |
| VVLF | 6.213e+01 | 1.977e+01 | 3.065e+01 | 7.991e-01 | 2.044e+01 |
| Wj0b_c | 5.133e+02 | 1.044e+02 | 1.527e+02 | 1.645e+01 | 7.126e+01 |
| Wj0b_udsg | 5.218e+02 | 1.039e+02 | 1.568e+02 | 2.600e+01 | 6.334e+01 |
| Wj1b | 3.279e+02 | 6.970e+01 | 8.666e+01 | 1.097e+01 | 2.953e+01 |
| Wj2b | 4.827e+02 | 1.604e+02 | 1.817e+02 | 3.198e+01 | 9.275e+01 |
| Zj0b_c | 2.968e+01 | 3.868e+00 | 5.701e+00 | 4.552e-01 | 1.209e+00 |
| Zj0b_udsg | 4.615e+01 | 3.884e+00 | 1.021e+01 | 5.188e-01 | 1.400e+00 |
| Zj1b | 5.880e+01 | 5.032e+00 | 5.481e+00 | 3.946e-01 | 8.907e-01 |
| Zj2b | 6.992e+01 | 6.915e+00 | 7.405e+00 | 6.539e-01 | 1.927e+00 |
| s_Top | 1.724e+03 | 2.801e+02 | 2.417e+02 | 2.087e+01 | 1.099e+02 |

Table C.6: The observed and predicted yields are given for the control regions for 1-lepton (μ) in 2016.

| Process | W+udcsg enr., med. $p_T(V)$ | W+udcsg enr., high $p_T(V)$ | W+udcsg enr., high $p_T(V)$, boosted |
|-----------|--------------------------------|--------------------------------|---|
| Data | 142862 | 32273 | 47879 |
| TT | 2.434e+04 | 5.758e+03 | 6.666e+03 |
| VVHF | 3.255e+01 | 1.459e+01 | 1.761e+01 |
| VVLF | 3.064e+03 | 1.098e+03 | 2.213e+03 |
| Wj0b_c | 2.029e+04 | 4.513e+03 | 5.586e+03 |
| Wj0b_udsg | 7.003e+04 | 1.729e+04 | 3.488e+04 |
| Wj1b | 1.483e+03 | 4.007e+02 | 5.024e+02 |
| Wj2b | 4.767e+02 | 1.902e+02 | 3.026e+02 |
| Zj0b_c | 8.355e+02 | 1.060e+02 | 1.540e+02 |
| Zj0b_udsg | 5.102e+03 | 6.679e+02 | 1.386e+03 |
| Zj1b | 1.789e+02 | 2.034e+01 | 2.862e+01 |
| Zj2b | 3.429e+01 | 6.691e+00 | 1.223e+01 |
| s_Top | 3.971e+03 | 1.084e+03 | 1.319e+03 |

| Process | W+bb enr., med. $p_T(V)$ | W+bb enr., high $p_T(V)$ | W+bb enr., high $p_T(V)$, boosted |
|-----------|-----------------------------|-----------------------------|---------------------------------------|
| Data | 28053 | 4545 | 2548 |
| TT | 1.985e+04 | 3.019e+03 | 1.848e+03 |
| VVHF | 7.078e+01 | 1.932e+01 | 8.710e+00 |
| VVLF | 1.010e+02 | 2.428e+01 | 1.191e+02 |
| Wj0b_c | 1.117e+03 | 2.483e+02 | 2.620e+02 |
| Wj0b_udsg | 1.069e+03 | 2.638e+02 | 2.446e+02 |
| Wj1b | 7.172e+02 | 1.623e+02 | 1.406e+02 |
| Wj2b | 1.082e+03 | 2.992e+02 | 3.850e+02 |
| Zj0b_c | 7.770e+01 | 6.280e+00 | 9.266e+00 |
| Zj0b_udsg | 9.657e+01 | 1.436e+01 | 2.218e+01 |
| Zj1b | 9.800e+01 | 8.921e+00 | 9.081e+00 |
| Zj2b | 1.165e+02 | 1.253e+01 | 1.402e+01 |
| s_Top | 2.764e+03 | 6.598e+02 | 3.339e+02 |

| Process | $t\bar{t}$ enr., med. $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$, boosted |
|-----------|-----------------------------------|-----------------------------------|---|
| Data | 50788 | 12017 | 4244 |
| TT | 4.647e+04 | 1.103e+04 | 5.029e+03 |
| VVHF | 8.711e+01 | 4.722e+01 | 4.302e+00 |
| VVLF | 6.077e+01 | 2.759e+01 | 9.766e+00 |
| Wj0b_c | 3.250e+02 | 1.275e+02 | 1.982e+01 |
| Wj0b_udsg | 3.234e+02 | 1.365e+02 | 9.878e+00 |
| Wj1b | 8.308e+02 | 3.153e+02 | 2.039e+01 |
| Wj2b | 1.030e+03 | 4.480e+02 | 7.608e+01 |
| Zj0b_c | 2.010e+01 | 4.969e+00 | 9.419e-01 |
| Zj0b_udsg | 2.642e+01 | 7.535e+00 | 1.095e+00 |
| Zj1b | 9.908e+01 | 1.860e+01 | 1.435e+00 |
| Zj2b | 1.073e+02 | 2.229e+01 | 5.295e+00 |
| s_Top | 3.127e+03 | 1.264e+03 | 3.906e+02 |

Table C.7: The observed and predicted yields are given for the signal regions for 2-lepton (e) in 2016.

| Process | Low $p_T(V)$ | Medium $p_T(V)$ 0J | Medium $p_T(V)$ GE1J | High $p_T(V)$ | High $p_T(V)$, boosted | Highest $p_T(V)$ | Highest $p_T(V)$, boosted |
|-------------------------|-----------------|-----------------------|----------------------------|------------------|-------------------------------|---------------------|----------------------------------|
| Data | 2478 | 234 | 292 | 108 | 52 | 12 | 18 |
| ZH 0_75 | 9.538e-02 | 0 | 0 | 0 | 0 | 0 | 0 |
| ZH 150_250 0J | 1.721e-01 | 4.048e+00 | 6.058e-01 | 2.573e-02 | 1.095e-03 | 0 | 0 |
| ZH 150_250 GE1J | 7.791e-02 | 5.432e-01 | 1.733e+00 | 9.020e-03 | 3.283e-03 | 0 | 0 |
| ZH 250_400 | 0 | 6.703e-02 | 3.716e-02 | 2.274e+00 | 3.555e-01 | 7.597e-03 | 4.489e-03 |
| ZH 75_150 | 1.312e+01 | 5.066e-02 | 2.020e-02 | 0 | 0 | 0 | 0 |
| ZH GT400 | 0 | 0 | 0 | 2.072e-02 | 2.095e-03 | 4.797e-01 | 2.361e-01 |
| ggZH 0_75 | 8.620e-03 | 0 | 0 | 0 | 0 | 0 | 0 |
| ggZH 150_250 0J | 5.764e-02 | 1.150e+00 | 1.833e-01 | 3.648e-03 | 0 | 0 | 0 |
| ggZH 150_250 GE1J | 5.569e-02 | 5.661e-01 | 1.371e+00 | 8.899e-03 | 5.241e-04 | 0 | 0 |
| ggZH 250_400 | 0 | 1.471e-02 | 1.984e-02 | 6.971e-01 | 1.250e-01 | 1.338e-03 | 5.944e-04 |
| ggZH 75_150 | 3.695e+00 | 2.109e-02 | 1.220e-02 | 0 | 0 | 0 | 0 |
| ggZH GT400 | 0 | 0 | 0 | 4.722e-03 | 2.156e-03 | 6.754e-02 | 3.801e-02 |
| TT | 5.685e+02 | 1.746e+01 | 2.116e+01 | 2.151e+00 | 6.171e-01 | 0 | 0 |
| VVHF | 3.261e+01 | 7.761e+00 | 6.748e+00 | 5.616e+00 | 1.486e+00 | 7.302e-01 | 4.118e-01 |
| VVLF | 1.509e+01 | 2.074e+00 | 1.875e+00 | 1.238e+00 | 2.246e+00 | 5.842e-02 | 8.934e-01 |
| Zj0b_c | 2.221e+02 | 2.743e+01 | 3.311e+01 | 1.333e+01 | 8.462e+00 | 2.051e+00 | 3.351e+00 |
| Zj0b_udsg | 3.374e+02 | 1.902e+01 | 4.046e+01 | 1.639e+01 | 1.204e+01 | 2.821e+00 | 4.843e+00 |
| Zj1b | 6.742e+02 | 5.491e+01 | 5.389e+01 | 1.795e+01 | 1.217e+01 | 2.492e+00 | 3.251e+00 |
| Zj2b | 5.994e+02 | 6.390e+01 | 6.629e+01 | 2.919e+01 | 1.791e+01 | 4.618e+00 | 8.243e+00 |
| s_Top | 2.582e+01 | 7.407e-01 | 1.750e+00 | 3.764e-01 | 0 | 0 | 0 |

Table C.8: The observed and predicted yields are given for the control regions for 2-lepton (e) in 2016.

| Process | Z+udcsg enr., low $p_T(V)$ | Z+udcsg enr., med. $p_T(V)$ | Z+udcsg enr., high $p_T(V)$ | Z+udcsg enr., high $p_T(V)$, boosted |
|-----------|----------------------------------|-----------------------------------|-----------------------------------|---|
| Data | 32439 | 6944 | 1325 | 843 |
| TT | 6.175e+01 | 5.378e+00 | 0 | 4.557e-01 |
| VVHF | 1.472e+00 | 5.498e-01 | 2.765e-01 | 1.024e+00 |
| VVLF | 2.073e+02 | 6.213e+01 | 1.521e+01 | 2.033e+01 |
| Zj0b_c | 1.764e+03 | 3.333e+02 | 8.301e+01 | 1.220e+02 |
| Zj0b_udsg | 3.223e+04 | 6.009e+03 | 1.211e+03 | 7.673e+02 |
| Zj1b | 3.540e+02 | 6.422e+01 | 1.221e+01 | 3.903e+01 |
| Zj2b | 2.945e+01 | 6.816e+00 | 1.885e+00 | 2.033e+01 |
| s_Top | 7.798e+00 | 1.109e+00 | 0 | 2.329e-01 |
| Process | Z+bb enr., low $p_T(V)$ | Z+bb enr., med. $p_T(V)$ | Z+bb enr., high $p_T(V)$ | Z+bb enr., high $p_T(V)$, boosted |
| Data | 3001 | 795 | 171 | 77 |
| TT | 3.387e+02 | 1.956e+01 | 7.755e-01 | 8.541e-01 |
| VVHF | 3.179e+01 | 1.263e+01 | 3.497e+00 | 1.382e+00 |
| VVLF | 1.355e+01 | 4.871e+00 | 1.454e+00 | 3.168e+00 |
| Zj0b_c | 3.448e+02 | 9.059e+01 | 2.615e+01 | 1.928e+01 |
| Zj0b_udsg | 4.584e+02 | 1.173e+02 | 3.146e+01 | 2.205e+01 |
| Zj1b | 9.570e+02 | 1.822e+02 | 3.439e+01 | 2.195e+01 |
| Zj2b | 8.844e+02 | 2.292e+02 | 5.217e+01 | 3.521e+01 |
| s_Top | 1.531e+01 | 2.523e+00 | 5.006e-01 | 3.363e-01 |
| Process | $t\bar{t}$ enr., low $p_T(V)$ | $t\bar{t}$ enr., med. $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$, boosted |
| Data | 9768 | 1011 | 84 | 42 |
| TT | 9.714e+03 | 8.896e+02 | 7.875e+01 | 1.396e+01 |
| VVHF | 7.954e+00 | 3.593e+00 | 1.106e+00 | 4.372e-01 |
| VVLF | 3.282e+00 | 9.188e-01 | 2.343e-01 | 1.889e+00 |
| Zj0b_c | 1.847e+01 | 4.388e+00 | 1.153e+00 | 5.458e+00 |
| Zj0b_udsg | 3.727e+01 | 4.997e+00 | 1.585e+00 | 5.493e+00 |
| Zj1b | 1.154e+02 | 2.219e+01 | 4.946e+00 | 4.049e+00 |
| Zj2b | 1.392e+02 | 3.314e+01 | 9.068e+00 | 7.211e+00 |
| s_Top | 3.775e+02 | 7.618e+01 | 1.605e+01 | 9.051e+00 |

Table C.9: The observed and predicted yields are given for the signal regions for 2-lepton (μ) in 2016.

| Process | Low $p_T(V)$ | Medium $p_T(V)$ 0J | Medium $p_T(V)$ GE1J | High $p_T(V)$ | High $p_T(V)$, boosted | Highest $p_T(V)$ | Highest $p_T(V)$, boosted |
|-------------------------|-----------------|-----------------------|----------------------------|------------------|-------------------------------|---------------------|----------------------------------|
| Data | 3842 | 332 | 374 | 135 | 73 | 12 | 36 |
| ZH 0_75 | 1.443e-01 | 0 | 0 | 0 | 0 | 0 | 0 |
| ZH 150_250 0J | 2.002e-01 | 5.410e+00 | 7.628e-01 | 5.285e-02 | 1.260e-04 | 0 | 0 |
| ZH 150_250 GE1J | 1.102e-01 | 7.763e-01 | 2.293e+00 | 4.025e-02 | 3.512e-03 | 0 | 0 |
| ZH 250_400 | 0 | 1.012e-01 | 5.866e-02 | 3.050e+00 | 4.853e-01 | 2.629e-02 | 7.796e-03 |
| ZH 75_150 | 1.872e+01 | 1.355e-01 | 5.663e-02 | 0 | 0 | 0 | 0 |
| ZH GT400 | 0 | 0 | 0 | 3.975e-02 | 1.540e-02 | 6.683e-01 | 4.154e-01 |
| ggZH 0_75 | 1.512e-02 | 0 | 0 | 0 | 0 | 0 | 0 |
| ggZH 150_250 0J | 7.090e-02 | 1.566e+00 | 2.368e-01 | 1.190e-02 | 0 | 0 | 0 |
| ggZH 150_250 GE1J | 7.971e-02 | 7.678e-01 | 1.855e+00 | 2.708e-02 | 2.043e-03 | 0 | 0 |
| ggZH 250_400 | 0 | 2.628e-02 | 3.569e-02 | 9.327e-01 | 1.640e-01 | 4.346e-03 | 1.996e-03 |
| ggZH 75_150 | 4.987e+00 | 4.276e-02 | 3.347e-02 | 0 | 0 | 0 | 0 |
| ggZH GT400 | 0 | 0 | 0 | 5.215e-03 | 3.877e-03 | 9.077e-02 | 5.374e-02 |
| TT | 8.450e+02 | 1.217e+01 | 2.144e+01 | 1.735e+00 | 3.223e+00 | 0 | 0 |
| VVHF | 4.729e+01 | 9.589e+00 | 9.398e+00 | 4.835e+00 | 1.620e+00 | 6.426e-01 | 6.430e-01 |
| VVLF | 2.178e+01 | 2.814e+00 | 4.348e+00 | 1.211e+00 | 3.049e+00 | 4.137e-01 | 1.680e+00 |
| Zj0b_c | 3.280e+02 | 3.884e+01 | 4.729e+01 | 1.787e+01 | 1.355e+01 | 2.589e+00 | 4.890e+00 |
| Zj0b_udsg | 4.631e+02 | 3.885e+01 | 7.390e+01 | 2.093e+01 | 1.528e+01 | 4.095e+00 | 6.173e+00 |
| Zj1b | 9.827e+02 | 7.556e+01 | 8.228e+01 | 2.659e+01 | 1.734e+01 | 3.787e+00 | 5.521e+00 |
| Zj2b | 8.415e+02 | 8.274e+01 | 9.380e+01 | 3.872e+01 | 2.588e+01 | 7.027e+00 | 1.208e+01 |
| s_Top | 2.933e+01 | 3.763e+00 | 4.099e+00 | 1.484e+00 | 3.822e-01 | 0 | 4.486e-01 |

Table C.10: The observed and predicted yields are given for the control regions for 2-lepton (μ) in 2016.

| Process | Z+udcsg enr., low $p_T(V)$ | Z+udcsg enr., med. $p_T(V)$ | Z+udcsg enr., high $p_T(V)$ | Z+udcsg enr., high $p_T(V)$, boosted |
|-----------|----------------------------------|-----------------------------------|-----------------------------------|---|
| Data | 48206 | 9545 | 1719 | 1163 |
| TT | 1.027e+02 | 1.326e+01 | 9.527e-01 | 6.617e+00 |
| VVHF | 2.069e+00 | 6.008e-01 | 1.863e-01 | 1.635e+00 |
| VVLF | 2.793e+02 | 8.190e+01 | 2.494e+01 | 2.857e+01 |
| Zj0b_c | 2.344e+03 | 4.663e+02 | 1.008e+02 | 1.809e+02 |
| Zj0b_udsg | 4.372e+04 | 8.170e+03 | 1.662e+03 | 1.078e+03 |
| Zj1b | 4.638e+02 | 8.890e+01 | 1.596e+01 | 5.479e+01 |
| Zj2b | 3.142e+01 | 8.627e+00 | 2.572e+00 | 2.852e+01 |
| s_Top | 1.009e+01 | 1.182e+00 | 2.228e-01 | 5.445e-01 |
| Process | Z+bb enr., low $p_T(V)$ | Z+bb enr., med. $p_T(V)$ | Z+bb enr., high $p_T(V)$ | Z+bb enr., high $p_T(V)$, boosted |
| Data | 4737 | 1125 | 234 | 128 |
| TT | 5.063e+02 | 2.992e+01 | 5.177e-01 | 1.444e+00 |
| VVHF | 4.581e+01 | 1.710e+01 | 4.736e+00 | 2.115e+00 |
| VVLF | 1.905e+01 | 6.396e+00 | 2.277e+00 | 3.906e+00 |
| Zj0b_c | 4.673e+02 | 1.288e+02 | 3.411e+01 | 2.739e+01 |
| Zj0b_udsg | 6.792e+02 | 1.690e+02 | 4.515e+01 | 3.105e+01 |
| Zj1b | 1.315e+03 | 2.466e+02 | 4.682e+01 | 3.048e+01 |
| Zj2b | 1.227e+03 | 3.127e+02 | 6.924e+01 | 5.268e+01 |
| s_Top | 2.254e+01 | 4.532e+00 | 3.693e-01 | 4.003e-01 |
| Process | $t\bar{t}$ enr., low $p_T(V)$ | $t\bar{t}$ enr., med. $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$, boosted |
| Data | 15396 | 1485 | 154 | 84 |
| TT | 1.362e+04 | 1.260e+03 | 1.138e+02 | 2.973e+01 |
| VVHF | 1.269e+01 | 4.208e+00 | 1.979e+00 | 4.603e-01 |
| VVLF | 4.902e+00 | 1.263e+00 | 1.123e-01 | 3.373e+00 |
| Zj0b_c | 4.336e+01 | 8.082e+00 | 2.100e+00 | 9.983e+00 |
| Zj0b_udsg | 4.271e+01 | 9.574e+00 | 4.269e+00 | 1.572e+01 |
| Zj1b | 2.004e+02 | 3.670e+01 | 8.129e+00 | 7.660e+00 |
| Zj2b | 2.276e+02 | 6.146e+01 | 1.810e+01 | 1.529e+01 |
| s_Top | 4.979e+02 | 1.125e+02 | 1.752e+01 | 1.490e+01 |

Table C.11: The observed and predicted yields are given for the signal regions for 0-lepton in 2017.

| Process | Medium $p_T(V)$ 0J | Medium $p_T(V)$ GE1J | High $p_T(V)$ | High $p_T(V)$, boosted | Highest $p_T(V)$ | Highest $p_T(V)$, boosted |
|-------------------------|-----------------------|----------------------------|------------------|-------------------------------|---------------------|----------------------------------|
| Data | 3517 | 3942 | 1787 | 2178 | 132 | 493 |
| WH 150_250 0J | 6.525e+00 | 1.151e+00 | 4.862e-01 | 1.102e-02 | 0 | 0 |
| WH 150_250 GE1J | 9.167e-01 | 1.757e+00 | 2.232e-01 | 1.422e-02 | 0 | 0 |
| WH 250_400 | 1.938e+00 | 1.079e+00 | 3.604e+00 | 6.879e-01 | 4.006e-02 | 2.263e-02 |
| WH 75_150 | 6.067e-01 | 1.454e-01 | 3.344e-03 | 0 | 0 | 0 |
| WH GT400 | 4.012e-02 | 1.552e-02 | 4.324e-01 | 2.015e-01 | 4.269e-01 | 2.856e-01 |
| ZH 0_75 | 0 | 0 | 7.024e-05 | 0 | 0 | 0 |
| ZH 150_250 0J | 2.230e+01 | 2.264e+00 | 2.374e+00 | 1.071e-02 | 0 | 0 |
| ZH 150_250 GE1J | 3.041e+00 | 5.752e+00 | 1.074e+00 | 7.932e-02 | 0 | 0 |
| ZH 250_400 | 5.077e+00 | 1.758e+00 | 1.759e+01 | 3.106e+00 | 5.365e-01 | 1.868e-01 |
| ZH 75_150 | 1.617e+00 | 4.150e-01 | 7.042e-03 | 0 | 0 | 0 |
| ZH GT400 | 1.911e-02 | 0 | 1.375e+00 | 5.896e-01 | 3.591e+00 | 2.809e+00 |
| ggZH 0_75 | 4.031e-04 | 0 | 0 | 0 | 0 | 0 |
| ggZH 150_250 0J | 6.024e+00 | 6.618e-01 | 5.750e-01 | 4.992e-03 | 0 | 0 |
| ggZH 150_250 GE1J | 2.357e+00 | 4.031e+00 | 7.207e-01 | 8.634e-02 | 1.669e-06 | 0 |
| ggZH 250_400 | 1.081e+00 | 7.692e-01 | 4.272e+00 | 8.838e-01 | 1.023e-01 | 4.760e-02 |
| ggZH 75_150 | 6.208e-01 | 2.435e-01 | 3.867e-03 | 4.688e-04 | 0 | 0 |
| ggZH GT400 | 1.303e-03 | 3.375e-04 | 1.871e-01 | 9.133e-02 | 3.556e-01 | 2.588e-01 |
| TT | 1.572e+03 | 2.401e+03 | 7.255e+02 | 7.945e+02 | 1.605e+01 | 8.150e+01 |
| VVHF | 9.968e+01 | 4.924e+01 | 7.134e+01 | 2.819e+01 | 8.717e+00 | 1.244e+01 |
| VVLF | 2.985e+01 | 2.112e+01 | 2.473e+01 | 4.020e+01 | 1.594e+00 | 9.143e+00 |
| Wj0b_c | 8.633e+01 | 9.060e+01 | 6.001e+01 | 9.490e+01 | 2.909e+00 | 1.726e+01 |
| Wj0b_udsg | 9.339e+01 | 7.690e+01 | 4.821e+01 | 9.944e+01 | 3.872e+00 | 1.943e+01 |
| Wj1b | 5.688e+01 | 5.159e+01 | 3.295e+01 | 4.947e+01 | 2.060e+00 | 9.091e+00 |
| Wj2b | 1.290e+02 | 1.201e+02 | 9.334e+01 | 9.753e+01 | 1.103e+01 | 2.187e+01 |
| Zj0b_c | 1.475e+02 | 9.531e+01 | 8.939e+01 | 1.253e+02 | 1.012e+01 | 4.077e+01 |
| Zj0b_udsg | 7.939e+01 | 6.434e+01 | 6.476e+01 | 1.336e+02 | 8.547e+00 | 4.703e+01 |
| Zj1b | 1.841e+02 | 9.792e+01 | 9.285e+01 | 1.278e+02 | 8.445e+00 | 3.107e+01 |
| Zj2b | 3.857e+02 | 2.463e+02 | 2.301e+02 | 1.556e+02 | 2.784e+01 | 6.055e+01 |
| s_Top | 1.840e+02 | 2.015e+02 | 7.780e+01 | 6.451e+01 | 1.952e+00 | 1.365e+01 |

Table C.12: The observed and predicted yields are given for the control regions for 0-lepton in 2017.

| Process | Z+udcsg enr., med. $p_T(V)$ | Z+udcsg enr., high $p_T(V)$ | Z+udcsg enr., high $p_T(V)$, boosted |
|-----------|--------------------------------|--------------------------------|---|
| Data | 9341 | 3286 | 79789 |
| TT | 1.415e+03 | 3.452e+02 | 3.154e+03 |
| VVHF | 1.691e+01 | 6.540e+00 | 3.202e+01 |
| VVLF | 1.556e+02 | 5.183e+01 | 1.690e+03 |
| Wj0b_c | 8.973e+02 | 2.041e+02 | 2.596e+03 |
| Wj0b_udsg | 2.231e+03 | 6.510e+02 | 2.380e+04 |
| Wj1b | 7.235e+01 | 2.001e+01 | 2.239e+02 |
| Wj2b | 4.304e+01 | 1.673e+01 | 1.101e+02 |
| Zj0b_c | 9.979e+02 | 3.476e+02 | 3.796e+03 |
| Zj0b_udsg | 1.812e+03 | 8.137e+02 | 3.586e+04 |
| Zj1b | 1.876e+02 | 6.596e+01 | 6.781e+02 |
| Zj2b | 9.290e+01 | 3.952e+01 | 2.117e+02 |
| s_Top | 1.801e+02 | 4.438e+01 | 5.380e+02 |

| Process | Z+bb enr., med. $p_T(V)$ | Z+bb enr., high $p_T(V)$ | Z+bb enr., high $p_T(V)$, boosted |
|-----------|-----------------------------|-----------------------------|---------------------------------------|
| Data | 13149 | 3508 | 2970 |
| TT | 5.779e+03 | 1.190e+03 | 8.650e+02 |
| VVHF | 1.100e+02 | 4.172e+01 | 2.812e+01 |
| VVLF | 4.393e+01 | 8.120e+00 | 4.627e+01 |
| Wj0b_c | 4.654e+02 | 1.100e+02 | 1.293e+02 |
| Wj0b_udsg | 4.761e+02 | 1.407e+02 | 1.832e+02 |
| Wj1b | 2.488e+02 | 6.411e+01 | 8.236e+01 |
| Wj2b | 5.602e+02 | 1.627e+02 | 1.927e+02 |
| Zj0b_c | 5.447e+02 | 1.899e+02 | 2.037e+02 |
| Zj0b_udsg | 3.570e+02 | 1.532e+02 | 2.578e+02 |
| Zj1b | 6.298e+02 | 1.963e+02 | 2.632e+02 |
| Zj2b | 1.360e+03 | 4.538e+02 | 3.382e+02 |
| s_Top | 6.969e+02 | 1.692e+02 | 9.035e+01 |

| Process | $t\bar{t}$ enr., med. $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$, boosted |
|-----------|-----------------------------------|-----------------------------------|---|
| Data | 3250 | 915 | 2250 |
| TT | 2.739e+03 | 6.866e+02 | 1.947e+03 |
| VVHF | 3.258e+00 | 1.133e+00 | 6.451e+00 |
| VVLF | 2.107e+00 | 8.234e-01 | 3.545e+00 |
| Wj0b_c | 3.273e+01 | 1.077e+01 | 1.672e+01 |
| Wj0b_udsg | 3.426e+01 | 9.288e+00 | 8.567e+00 |
| Wj1b | 1.597e+01 | 4.720e+00 | 1.332e+01 |
| Wj2b | 2.468e+01 | 9.192e+00 | 3.787e+01 |
| Zj0b_c | 1.942e+01 | 9.638e+00 | 1.380e+01 |
| Zj0b_udsg | 1.384e+01 | 8.046e+00 | 9.211e+00 |
| Zj1b | 1.988e+01 | 9.141e+00 | 2.597e+01 |
| Zj2b | 4.638e+01 | 2.100e+01 | 9.221e+01 |
| s_Top | 1.152e+02 | 3.578e+01 | 1.218e+02 |

Table C.13: The observed and predicted yields are given for the signal regions for 1-lepton (e) in 2017.

| Process | Medium $p_T(V)$ | High $p_T(V)$ | High $p_T(V)$, boosted | Highest $p_T(V)$ | Highest $p_T(V)$, boosted |
|------------|--------------------|---------------|----------------------------|---------------------|----------------------------------|
| Data | 13949 | 1826 | 2127 | 106 | 537 |
| WH 0_75 | 1.418e-02 | 0 | 0 | 0 | 0 |
| WH 150_250 | 2.659e+01 | 1.625e+00 | 6.539e-03 | 0 | 0 |
| 0J | | | | | |
| WH 150_250 | 9.725e+00 | 7.076e-01 | 1.236e-01 | 0 | 0 |
| GE1J | | | | | |
| WH 250_400 | 5.286e+00 | 1.265e+01 | 2.364e+00 | 4.084e-01 | 1.532e-01 |
| WH 75_150 | 7.872e+00 | 1.541e-03 | 2.247e-03 | 0 | 0 |
| WH GT400 | 1.401e-02 | 1.291e+00 | 6.061e-01 | 2.990e+00 | 2.533e+00 |
| ZH 0_75 | 5.364e-03 | 0 | 0 | 0 | 0 |
| ZH 150_250 | 6.486e-01 | 2.807e-02 | 4.764e-04 | 0 | 0 |
| 0J | | | | | |
| ZH 150_250 | 2.260e-01 | 1.036e-02 | 1.141e-03 | 0 | 0 |
| GE1J | | | | | |
| ZH 250_400 | 2.498e-01 | 1.672e-01 | 5.078e-02 | 2.242e-03 | 0 |
| ZH 75_150 | 3.428e-01 | 3.799e-05 | 0 | 0 | 0 |
| ZH GT400 | 8.527e-03 | 3.945e-02 | 2.457e-02 | 1.977e-02 | 2.104e-02 |
| TT | 1.158e+04 | 1.279e+03 | 1.696e+03 | 5.609e+01 | 2.835e+02 |
| VVHF | 8.274e+01 | 2.100e+01 | 1.055e+01 | 3.531e+00 | 2.584e+00 |
| VVLF | 5.138e+01 | 1.148e+01 | 2.908e+01 | 6.610e-01 | 1.252e+01 |
| Wj0b_c | 6.022e+02 | 9.819e+01 | 1.315e+02 | 1.500e+01 | 4.313e+01 |
| Wj0b_udsg | 4.274e+02 | 6.417e+01 | 1.075e+02 | 1.461e+01 | 3.442e+01 |
| Wj1b | 2.768e+02 | 5.659e+01 | 6.607e+01 | 6.229e+00 | 1.759e+01 |
| Wj2b | 4.191e+02 | 1.274e+02 | 1.445e+02 | 2.156e+01 | 5.506e+01 |
| Zj0b_c | 3.495e+01 | 2.973e+00 | 4.878e+00 | 3.237e-01 | 9.950e-01 |
| Zj0b_udsg | 3.561e+01 | 4.847e+00 | 4.773e+00 | 2.669e-02 | 9.319e-01 |
| Zj1b | 5.321e+01 | 2.422e+00 | 3.704e+00 | 1.224e-01 | 5.530e-01 |
| Zj2b | 5.692e+01 | 6.474e+00 | 5.370e+00 | 3.893e-01 | 1.332e+00 |
| s_Top | 1.632e+03 | 2.550e+02 | 2.426e+02 | 1.657e+01 | 8.558e+01 |

Table C.14: The observed and predicted yields are given for the control regions for 1-lepton (e) in 2017.

| Process | W+udcsg enr., med. $p_T(V)$ | W+udcsg enr., high $p_T(V)$ | W+udcsg enr., high $p_T(V)$, boosted |
|-----------|-----------------------------------|-----------------------------------|---|
| Data | 127417 | 27731 | 40956 |
| TT | 2.313e+04 | 5.286e+03 | 5.695e+03 |
| VVHF | 5.414e+01 | 1.822e+01 | 1.436e+01 |
| VVLF | 2.191e+03 | 7.324e+02 | 1.534e+03 |
| Wj0b_c | 2.650e+04 | 4.872e+03 | 4.599e+03 |
| Wj0b_udsg | 6.696e+04 | 1.498e+04 | 2.871e+04 |
| Wj1b | 1.181e+03 | 2.810e+02 | 2.661e+02 |
| Wj2b | 3.486e+02 | 1.339e+02 | 1.435e+02 |
| Zj0b_c | 8.695e+02 | 9.443e+01 | 1.049e+02 |
| Zj0b_udsg | 4.717e+03 | 5.236e+02 | 1.013e+03 |
| Zj1b | 1.762e+02 | 1.670e+01 | 1.799e+01 |
| Zj2b | 2.761e+01 | 4.001e+00 | 5.696e+00 |
| s_Top | 3.732e+03 | 9.610e+02 | 1.049e+03 |
| Process | W+bb enr., med. $p_T(V)$ | W+bb enr., high $p_T(V)$ | W+bb enr., high $p_T(V)$, boosted |
| Data | 24941 | 4103 | 2345 |
| TT | 1.943e+04 | 2.917e+03 | 1.709e+03 |
| VVHF | 7.643e+01 | 2.173e+01 | 1.162e+01 |
| VVLF | 8.407e+01 | 1.519e+01 | 6.308e+01 |
| Wj0b_c | 1.244e+03 | 2.301e+02 | 1.870e+02 |
| Wj0b_udsg | 1.027e+03 | 1.881e+02 | 1.798e+02 |
| Wj1b | 6.172e+02 | 1.174e+02 | 1.002e+02 |
| Wj2b | 9.861e+02 | 2.475e+02 | 2.542e+02 |
| Zj0b_c | 7.991e+01 | 7.877e+00 | 6.622e+00 |
| Zj0b_udsg | 7.274e+01 | 1.086e+01 | 9.457e+00 |
| Zj1b | 1.011e+02 | 8.157e+00 | 7.164e+00 |
| Zj2b | 1.112e+02 | 1.017e+01 | 8.770e+00 |
| s_Top | 2.716e+03 | 5.945e+02 | 2.660e+02 |
| Process | $t\bar{t}$ enr., med. $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$, boosted |
| Data | 45244 | 10894 | 4127 |
| TT | 4.431e+04 | 1.039e+04 | 4.966e+03 |
| VVHF | 6.674e+01 | 3.137e+01 | 3.626e+00 |
| VVLF | 4.447e+01 | 1.663e+01 | 4.968e+00 |
| Wj0b_c | 3.004e+02 | 8.725e+01 | 1.309e+01 |
| Wj0b_udsg | 2.719e+02 | 9.144e+01 | 7.953e+00 |
| Wj1b | 5.947e+02 | 1.945e+02 | 1.111e+01 |
| Wj2b | 7.838e+02 | 3.012e+02 | 4.308e+01 |
| Zj0b_c | 1.654e+01 | 3.480e+00 | 7.343e-01 |
| Zj0b_udsg | 2.267e+01 | 1.489e+01 | 2.604e-01 |
| Zj1b | 7.937e+01 | 1.313e+01 | 9.288e-01 |
| Zj2b | 8.508e+01 | 1.514e+01 | 2.818e+00 |
| s_Top | 2.883e+03 | 1.060e+03 | 3.409e+02 |

Table C.15: The observed and predicted yields are given for the signal regions for 1-lepton (μ) in 2017.

| Process | Medium $p_T(V)$ | High $p_T(V)$ | High $p_T(V)$, boosted | Highest $p_T(V)$ | Highest $p_T(V)$, boosted |
|------------|--------------------|---------------|----------------------------|---------------------|----------------------------------|
| Data | 18798 | 2464 | 2732 | 162 | 665 |
| WH 0_75 | 3.694e-02 | 0 | 0 | 0 | 0 |
| WH 150_250 | 3.311e+01 | 2.020e+00 | 1.831e-02 | 0 | 0 |
| 0J | | | | | |
| WH 150_250 | 1.134e+01 | 8.578e-01 | 9.079e-02 | 0 | 0 |
| GE1J | | | | | |
| WH 250_400 | 6.230e+00 | 1.534e+01 | 2.909e+00 | 5.658e-01 | 1.857e-01 |
| WH 75_150 | 1.014e+01 | 1.005e-02 | 0 | 0 | 0 |
| WH GT400 | 1.061e-02 | 1.434e+00 | 5.902e-01 | 3.441e+00 | 2.965e+00 |
| ZH 0_75 | 6.183e-03 | 0 | 0 | 0 | 0 |
| ZH 150_250 | 1.040e+00 | 4.830e-02 | 0 | 0 | 0 |
| 0J | | | | | |
| ZH 150_250 | 3.461e-01 | 1.816e-02 | 1.883e-03 | 0 | 0 |
| GE1J | | | | | |
| ZH 250_400 | 2.535e-01 | 2.538e-01 | 5.431e-02 | 4.154e-03 | 3.046e-03 |
| ZH 75_150 | 5.166e-01 | 9.071e-04 | 0 | 0 | 0 |
| ZH GT400 | 1.602e-02 | 3.079e-02 | 1.761e-02 | 1.983e-02 | 1.081e-02 |
| TT | 1.439e+04 | 1.584e+03 | 2.020e+03 | 5.080e+01 | 3.335e+02 |
| VVHF | 9.825e+01 | 2.551e+01 | 1.165e+01 | 3.004e+00 | 5.589e+00 |
| VVLF | 6.873e+01 | 1.513e+01 | 5.244e+01 | 6.777e+00 | 2.489e+01 |
| Wj0b_c | 8.628e+02 | 1.471e+02 | 1.632e+02 | 1.006e+01 | 5.310e+01 |
| Wj0b_udsg | 5.355e+02 | 1.099e+02 | 1.605e+02 | 7.281e+00 | 5.388e+01 |
| Wj1b | 3.299e+02 | 6.156e+01 | 8.005e+01 | 8.450e+00 | 1.986e+01 |
| Wj2b | 5.197e+02 | 1.652e+02 | 1.739e+02 | 2.595e+01 | 6.220e+01 |
| Zj0b_c | 4.129e+01 | 1.208e+00 | 5.689e+00 | 1.494e-01 | 1.688e+00 |
| Zj0b_udsg | 4.090e+01 | 6.197e+00 | 8.876e+00 | 8.134e-01 | 1.861e+00 |
| Zj1b | 6.614e+01 | 4.489e+00 | 5.209e+00 | 1.710e-01 | 8.871e-01 |
| Zj2b | 8.463e+01 | 8.702e+00 | 6.330e+00 | 8.084e-01 | 1.107e+00 |
| s_Top | 2.062e+03 | 3.213e+02 | 3.128e+02 | 2.509e+01 | 1.070e+02 |

Table C.16: The observed and predicted yields are given for the control regions for 1-lepton (μ) in 2017.

| Process | W+udcsg enr., med. $p_T(V)$ | W+udcsg enr., high $p_T(V)$ | W+udcsg enr., high $p_T(V)$, boosted |
|-----------|--------------------------------|--------------------------------|---|
| Data | 170543 | 36717 | 50184 |
| TT | 2.830e+04 | 6.347e+03 | 6.625e+03 |
| VVHF | 6.997e+01 | 2.199e+01 | 2.028e+01 |
| VVLF | 2.882e+03 | 9.800e+02 | 1.889e+03 |
| Wj0b_c | 3.302e+04 | 6.040e+03 | 5.453e+03 |
| Wj0b_udsg | 8.513e+04 | 1.893e+04 | 3.462e+04 |
| Wj1b | 1.518e+03 | 3.494e+02 | 3.185e+02 |
| Wj2b | 4.375e+02 | 1.646e+02 | 1.780e+02 |
| Zj0b_c | 1.159e+03 | 1.339e+02 | 1.344e+02 |
| Zj0b_udsg | 6.451e+03 | 7.281e+02 | 1.344e+03 |
| Zj1b | 2.369e+02 | 2.451e+01 | 2.396e+01 |
| Zj2b | 4.248e+01 | 7.277e+00 | 7.447e+00 |
| s_Top | 4.666e+03 | 1.153e+03 | 1.239e+03 |

| Process | W+bb enr., med. $p_T(V)$ | W+bb enr., high $p_T(V)$ | W+bb enr., high $p_T(V)$, boosted |
|-----------|-----------------------------|-----------------------------|---------------------------------------|
| Data | 34502 | 5370 | 2934 |
| TT | 2.404e+04 | 3.449e+03 | 1.916e+03 |
| VVHF | 9.896e+01 | 2.161e+01 | 1.411e+01 |
| VVLF | 1.123e+02 | 2.232e+01 | 6.996e+01 |
| Wj0b_c | 1.793e+03 | 2.914e+02 | 2.609e+02 |
| Wj0b_udsg | 1.136e+03 | 2.547e+02 | 2.299e+02 |
| Wj1b | 7.846e+02 | 1.467e+02 | 1.184e+02 |
| Wj2b | 1.255e+03 | 3.051e+02 | 3.072e+02 |
| Zj0b_c | 8.734e+01 | 7.739e+00 | 9.965e+00 |
| Zj0b_udsg | 1.137e+02 | 1.615e+01 | 1.455e+01 |
| Zj1b | 1.433e+02 | 1.006e+01 | 9.770e+00 |
| Zj2b | 1.633e+02 | 1.690e+01 | 1.108e+01 |
| s_Top | 3.424e+03 | 7.287e+02 | 3.197e+02 |

| Process | $t\bar{t}$ enr., med. $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$, boosted |
|-----------|-----------------------------------|-----------------------------------|---|
| Data | 60000 | 13808 | 5239 |
| TT | 5.412e+04 | 1.242e+04 | 5.786e+03 |
| VVHF | 8.366e+01 | 4.016e+01 | 3.656e+00 |
| VVLF | 6.860e+01 | 3.587e+01 | 2.761e+00 |
| Wj0b_c | 3.763e+02 | 1.110e+02 | 1.734e+01 |
| Wj0b_udsg | 3.443e+02 | 1.145e+02 | 1.058e+01 |
| Wj1b | 7.605e+02 | 2.414e+02 | 1.520e+01 |
| Wj2b | 9.595e+02 | 3.615e+02 | 5.273e+01 |
| Zj0b_c | 2.195e+01 | 3.310e+00 | 6.956e-01 |
| Zj0b_udsg | 2.987e+01 | 6.385e+00 | 7.106e-01 |
| Zj1b | 9.613e+01 | 1.569e+01 | 1.038e+00 |
| Zj2b | 1.114e+02 | 2.016e+01 | 4.148e+00 |
| s_Top | 3.508e+03 | 1.313e+03 | 4.174e+02 |

Table C.17: The observed and predicted yields are given for the signal regions for 2-lepton (e) in 2017.

| Process | Low $p_T(V)$ | Medium $p_T(V)$ 0J | Medium $p_T(V)$ GE1J | High $p_T(V)$ | High $p_T(V)$, boosted | Highest $p_T(V)$ | Highest $p_T(V)$, boosted |
|-------------------------|-----------------|-----------------------|----------------------------|------------------|-------------------------------|---------------------|----------------------------------|
| Data | 2761 | 273 | 327 | 130 | 78 | 13 | 29 |
| ZH 0_75 | 1.750e-01 | 0 | 0 | 0 | 0 | 0 | 0 |
| ZH 150_250 0J | 1.977e-01 | 5.088e+00 | 6.433e-01 | 3.382e-02 | 0 | 0 | 0 |
| ZH 150_250 GE1J | 7.274e-02 | 7.305e-01 | 2.136e+00 | 2.181e-02 | 1.854e-03 | 0 | 0 |
| ZH 250_400 | 0 | 6.883e-02 | 3.994e-02 | 2.878e+00 | 3.603e-01 | 1.202e-02 | 5.480e-03 |
| ZH 75_150 | 1.667e+01 | 1.189e-01 | 2.853e-02 | 0 | 0 | 0 | 0 |
| ZH GT400 | 0 | 0 | 0 | 2.573e-02 | 1.298e-02 | 6.193e-01 | 2.723e-01 |
| ggZH 0_75 | 1.366e-02 | 0 | 0 | 0 | 0 | 0 | 0 |
| ggZH 150_250 0J | 8.057e-02 | 1.792e+00 | 2.377e-01 | 9.407e-03 | 0 | 0 | 0 |
| ggZH 150_250 GE1J | 7.154e-02 | 7.384e-01 | 1.689e+00 | 1.463e-02 | 1.391e-03 | 0 | 0 |
| ggZH 250_400 | 0 | 1.667e-02 | 2.716e-02 | 9.354e-01 | 1.315e-01 | 3.634e-03 | 1.403e-03 |
| ggZH 75_150 | 5.049e+00 | 3.751e-02 | 2.720e-02 | 0 | 0 | 0 | 0 |
| ggZH GT400 | 0 | 0 | 0 | 5.431e-03 | 1.746e-03 | 8.830e-02 | 4.035e-02 |
| TT | 7.439e+02 | 1.228e+01 | 2.073e+01 | 0 | 3.314e-01 | 2.463e-01 | 0 |
| VVHF | 3.299e+01 | 6.787e+00 | 4.998e+00 | 3.336e+00 | 1.106e+00 | 7.033e-01 | 5.497e-01 |
| VVLF | 9.130e+00 | 1.564e+00 | 1.242e+00 | 6.538e-01 | 7.744e-01 | 1.564e-01 | 3.506e-01 |
| Zj0b_c | 2.957e+02 | 3.349e+01 | 3.968e+01 | 1.401e+01 | 1.245e+01 | 2.316e+00 | 4.094e+00 |
| Zj0b_udsg | 2.874e+02 | 3.464e+01 | 4.266e+01 | 2.008e+01 | 1.413e+01 | 1.887e+00 | 2.963e+00 |
| Zj1b | 7.915e+02 | 7.337e+01 | 6.230e+01 | 2.016e+01 | 1.451e+01 | 2.788e+00 | 3.169e+00 |
| Zj2b | 7.362e+02 | 8.954e+01 | 8.706e+01 | 4.075e+01 | 1.724e+01 | 4.978e+00 | 6.735e+00 |
| s_Top | 2.612e+01 | 1.715e+00 | 1.465e+00 | 3.041e-01 | 1.165e+00 | 0 | 0 |

Table C.18: The observed and predicted yields are given for the control regions for 2-lepton (e) in 2017.

| Process | Z+udcsg enr., low $p_T(V)$ | Z+udcsg enr., med. $p_T(V)$ | Z+udcsg enr., high $p_T(V)$ | Z+udcsg enr., high $p_T(V)$, boosted |
|-----------|----------------------------------|-----------------------------------|-----------------------------------|---|
| Data | 41833 | 8612 | 1717 | 1018 |
| TT | 7.976e+01 | 5.396e+00 | 4.966e-01 | 3.495e+00 |
| VVHF | 3.972e+00 | 1.234e+00 | 2.484e-01 | 6.720e-01 |
| VVLF | 2.119e+02 | 6.111e+01 | 1.494e+01 | 1.488e+01 |
| Zj0b_c | 2.249e+03 | 4.887e+02 | 8.662e+01 | 1.196e+02 |
| Zj0b_udsg | 4.349e+04 | 8.694e+03 | 1.673e+03 | 8.001e+02 |
| Zj1b | 4.056e+02 | 8.682e+01 | 1.695e+01 | 3.863e+01 |
| Zj2b | 2.705e+01 | 9.459e+00 | 2.251e+00 | 1.759e+01 |
| s_Top | 8.625e+00 | 1.129e+00 | 2.533e-01 | 4.488e-01 |
| Process | Z+bb enr., low $p_T(V)$ | Z+bb enr., med. $p_T(V)$ | Z+bb enr., high $p_T(V)$ | Z+bb enr., high $p_T(V)$, boosted |
| Data | 3537 | 932 | 222 | 123 |
| TT | 4.481e+02 | 2.518e+01 | 6.125e-01 | 1.104e+00 |
| VVHF | 2.867e+01 | 1.103e+01 | 3.215e+00 | 1.320e+00 |
| VVLF | 6.000e+00 | 2.292e+00 | 8.359e-01 | 9.310e-01 |
| Zj0b_c | 4.417e+02 | 1.406e+02 | 2.771e+01 | 1.995e+01 |
| Zj0b_udsg | 4.438e+02 | 1.241e+02 | 2.905e+01 | 2.419e+01 |
| Zj1b | 1.043e+03 | 2.227e+02 | 4.257e+01 | 2.518e+01 |
| Zj2b | 1.124e+03 | 3.056e+02 | 6.509e+01 | 3.870e+01 |
| s_Top | 1.726e+01 | 3.900e+00 | 0 | 1.860e-01 |
| Process | $t\bar{t}$ enr., low $p_T(V)$ | $t\bar{t}$ enr., med. $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$, boosted |
| Data | 12300 | 1240 | 118 | 87 |
| TT | 1.241e+04 | 1.123e+03 | 1.030e+02 | 2.702e+01 |
| VVHF | 6.536e+00 | 2.123e+00 | 8.447e-01 | 4.164e-01 |
| VVLF | 8.911e+00 | 7.944e-01 | 5.264e-01 | 3.886e+00 |
| Zj0b_c | 4.443e+01 | 1.025e+01 | 1.647e+00 | 5.752e+00 |
| Zj0b_udsg | 4.109e+01 | 8.556e+00 | 1.655e+00 | 7.094e+00 |
| Zj1b | 1.476e+02 | 2.621e+01 | 5.805e+00 | 5.939e+00 |
| Zj2b | 2.021e+02 | 4.878e+01 | 1.123e+01 | 8.915e+00 |
| s_Top | 4.456e+02 | 9.877e+01 | 1.452e+01 | 1.859e+01 |

Table C.19: The observed and predicted yields are given for the signal regions for 2-lepton (μ) in 2017.

| Process | Low $p_T(V)$ | Medium $p_T(V)$ 0J | Medium $p_T(V)$ GE1J | High $p_T(V)$ | High $p_T(V)$, boosted | Highest $p_T(V)$ | Highest $p_T(V)$, boosted |
|-------------------------|-----------------|-----------------------|----------------------------|------------------|-------------------------------|---------------------|----------------------------------|
| Data | 4133 | 354 | 413 | 174 | 106 | 16 | 31 |
| ZH 0_75 | 1.790e-01 | 0 | 0 | 0 | 0 | 0 | 0 |
| ZH 150_250 0J | 2.847e-01 | 6.542e+00 | 8.123e-01 | 6.978e-02 | 8.183e-09 | 0 | 0 |
| ZH 150_250 GE1J | 1.209e-01 | 9.774e-01 | 2.605e+00 | 3.429e-02 | 3.960e-03 | 0 | 0 |
| ZH 250_400 | 0 | 1.333e-01 | 6.798e-02 | 3.558e+00 | 3.957e-01 | 3.647e-02 | 7.580e-03 |
| ZH 75_150 | 2.210e+01 | 1.417e-01 | 5.190e-02 | 0 | 0 | 0 | 0 |
| ZH GT400 | 0 | 0 | 0 | 4.813e-02 | 1.798e-02 | 7.756e-01 | 3.945e-01 |
| ggZH 0_75 | 2.605e-02 | 0 | 0 | 0 | 0 | 0 | 0 |
| ggZH 150_250 0J | 1.103e-01 | 2.295e+00 | 3.094e-01 | 2.010e-02 | 0 | 0 | 0 |
| ggZH 150_250 GE1J | 1.014e-01 | 9.266e-01 | 2.210e+00 | 2.956e-02 | 1.416e-03 | 0 | 0 |
| ggZH 250_400 | 0 | 3.770e-02 | 4.257e-02 | 1.142e+00 | 1.650e-01 | 8.721e-03 | 2.215e-03 |
| ggZH 75_150 | 6.625e+00 | 6.998e-02 | 3.438e-02 | 0 | 0 | 0 | 0 |
| ggZH GT400 | 0 | 0 | 0 | 1.232e-02 | 3.953e-03 | 9.916e-02 | 5.060e-02 |
| TT | 1.006e+03 | 3.256e+01 | 3.746e+01 | 1.727e+00 | 1.919e+00 | 0 | 0 |
| VVHF | 4.349e+01 | 8.685e+00 | 6.474e+00 | 4.948e+00 | 1.465e+00 | 9.020e-01 | 7.292e-01 |
| VVLF | 1.046e+01 | 1.636e+00 | 1.911e+00 | 1.109e+00 | 1.473e+00 | 5.283e-02 | 7.281e-01 |
| Zj0b_c | 4.050e+02 | 4.059e+01 | 5.583e+01 | 2.061e+01 | 1.532e+01 | 2.350e+00 | 4.069e+00 |
| Zj0b_udsg | 4.158e+02 | 4.041e+01 | 5.926e+01 | 2.161e+01 | 1.635e+01 | 2.703e+00 | 4.642e+00 |
| Zj1b | 1.021e+03 | 9.766e+01 | 8.840e+01 | 2.952e+01 | 1.807e+01 | 3.040e+00 | 3.511e+00 |
| Zj2b | 1.032e+03 | 1.098e+02 | 1.082e+02 | 4.522e+01 | 2.311e+01 | 5.873e+00 | 8.465e+00 |
| s_Top | 4.079e+01 | 3.668e+00 | 2.479e+00 | 8.773e-01 | 2.453e-02 | 0 | 0 |

Table C.20: The observed and predicted yields are given for the control regions for 2-lepton (μ) in 2017.

| Process | Z+udcsg enr., low $p_T(V)$ | Z+udcsg enr., med. $p_T(V)$ | Z+udcsg enr., high $p_T(V)$ | Z+udcsg enr., high $p_T(V)$, boosted |
|-----------|----------------------------------|-----------------------------------|-----------------------------------|---|
| Data | 59966 | 11480 | 2213 | 1320 |
| TT | 1.112e+02 | 1.603e+01 | 6.773e-01 | 7.867e+00 |
| VVHF | 4.975e+00 | 1.565e+00 | 3.981e-01 | 8.253e-01 |
| VVLF | 2.748e+02 | 7.337e+01 | 1.975e+01 | 1.836e+01 |
| Zj0b_c | 2.875e+03 | 5.916e+02 | 1.084e+02 | 1.581e+02 |
| Zj0b_udsg | 5.657e+04 | 1.091e+04 | 2.105e+03 | 1.008e+03 |
| Zj1b | 5.628e+02 | 1.039e+02 | 1.945e+01 | 5.163e+01 |
| Zj2b | 3.505e+01 | 9.590e+00 | 2.400e+00 | 2.366e+01 |
| s_Top | 1.266e+01 | 1.712e+00 | 4.158e-01 | 1.176e+00 |
| Process | Z+bb enr., low $p_T(V)$ | Z+bb enr., med. $p_T(V)$ | Z+bb enr., high $p_T(V)$ | Z+bb enr., high $p_T(V)$, boosted |
| Data | 5029 | 1245 | 251 | 175 |
| TT | 6.372e+02 | 3.839e+01 | 8.654e-01 | 4.461e+00 |
| VVHF | 3.725e+01 | 1.388e+01 | 3.959e+00 | 2.373e+00 |
| VVLF | 7.647e+00 | 2.685e+00 | 1.054e+00 | 1.366e+00 |
| Zj0b_c | 5.644e+02 | 1.587e+02 | 3.810e+01 | 2.425e+01 |
| Zj0b_udsg | 6.259e+02 | 1.631e+02 | 4.301e+01 | 2.669e+01 |
| Zj1b | 1.420e+03 | 2.871e+02 | 5.238e+01 | 3.470e+01 |
| Zj2b | 1.506e+03 | 3.996e+02 | 7.789e+01 | 5.144e+01 |
| s_Top | 2.459e+01 | 4.882e+00 | 3.704e-01 | 1.821e-01 |
| Process | $t\bar{t}$ enr., low $p_T(V)$ | $t\bar{t}$ enr., med. $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$, boosted |
| Data | 18333 | 1717 | 169 | 122 |
| TT | 1.675e+04 | 1.533e+03 | 1.330e+02 | 5.020e+01 |
| VVHF | 9.141e+00 | 3.197e+00 | 1.148e+00 | 7.198e-01 |
| VVLF | 3.289e+00 | 5.260e+00 | 2.951e+00 | 4.437e+00 |
| Zj0b_c | 5.519e+01 | 1.730e+01 | 3.317e+00 | 1.180e+01 |
| Zj0b_udsg | 4.124e+01 | 1.213e+01 | 2.502e+00 | 1.588e+01 |
| Zj1b | 2.259e+02 | 4.693e+01 | 9.529e+00 | 1.246e+01 |
| Zj2b | 2.814e+02 | 7.289e+01 | 2.269e+01 | 1.586e+01 |
| s_Top | 5.785e+02 | 1.315e+02 | 2.160e+01 | 2.727e+01 |

Table C.21: The observed and predicted yields are given for the signal regions for 0-lepton in 2018.

| Process | Medium $p_T(V)$ 0J | Medium $p_T(V)$ GE1J | High $p_T(V)$ | High $p_T(V)$, boosted | Highest $p_T(V)$ | Highest $p_T(V)$, boosted |
|-------------------------|-----------------------|----------------------------|------------------|-------------------------------|---------------------|----------------------------------|
| Data | 4366 | 4602 | 2650 | 1685 | 215 | 347 |
| WH 150_250 0J | 7.915e+00 | 1.080e+00 | 4.422e-01 | 0 | 0 | 0 |
| WH 150_250 GE1J | 1.043e+00 | 2.077e+00 | 1.978e-01 | 1.931e-02 | 0 | 0 |
| WH 250_400 | 2.361e+00 | 1.459e+00 | 4.706e+00 | 7.740e-01 | 3.834e-02 | 1.119e-02 |
| WH 75_150 | 3.045e-01 | 1.219e-01 | 0 | 0 | 0 | 0 |
| WH GT400 | 2.061e-02 | 2.518e-02 | 6.051e-01 | 2.477e-01 | 5.549e-01 | 3.466e-01 |
| ZH 150_250 0J | 2.700e+01 | 2.604e+00 | 2.578e+00 | 9.690e-03 | 0 | 0 |
| ZH 150_250 GE1J | 3.626e+00 | 7.010e+00 | 1.117e+00 | 9.824e-02 | 0 | 0 |
| ZH 250_400 | 5.745e+00 | 2.078e+00 | 2.269e+01 | 3.465e+00 | 5.773e-01 | 1.735e-01 |
| ZH 75_150 | 1.083e+00 | 2.965e-01 | 0 | 0 | 0 | 0 |
| ZH GT400 | 1.081e-02 | 4.248e-03 | 1.897e+00 | 7.018e-01 | 5.076e+00 | 3.233e+00 |
| ggZH 150_250 0J | 7.161e+00 | 8.040e-01 | 5.466e-01 | 6.478e-03 | 0 | 0 |
| ggZH 150_250 GE1J | 2.755e+00 | 4.897e+00 | 6.555e-01 | 8.022e-02 | 0 | 0 |
| ggZH 250_400 | 1.339e+00 | 9.974e-01 | 5.723e+00 | 9.787e-01 | 1.041e-01 | 3.854e-02 |
| ggZH 75_150 | 4.532e-01 | 2.043e-01 | 8.407e-04 | 0 | 0 | 0 |
| ggZH GT400 | 3.370e-04 | 5.812e-04 | 2.378e-01 | 1.046e-01 | 5.053e-01 | 3.111e-01 |
| TT | 1.836e+03 | 3.001e+03 | 8.563e+02 | 7.961e+02 | 2.266e+01 | 8.071e+01 |
| VVHF | 1.099e+02 | 6.280e+01 | 8.554e+01 | 2.586e+01 | 1.202e+01 | 8.848e+00 |
| VVLF | 2.430e+01 | 1.812e+01 | 3.538e+01 | 2.419e+01 | 3.460e+00 | 1.202e+01 |
| Wj0b_c | 1.730e+02 | 1.197e+02 | 9.644e+01 | 3.831e+01 | 1.885e+00 | 8.100e+00 |
| Wj0b_udsg | 1.626e+02 | 1.528e+02 | 1.372e+02 | 2.192e+01 | 1.517e+01 | 1.382e+00 |
| Wj1b | 7.076e+01 | 5.824e+01 | 4.559e+01 | 4.634e+01 | 3.268e+00 | 6.576e+00 |
| Wj2b | 1.540e+02 | 1.497e+02 | 1.260e+02 | 1.070e+02 | 1.341e+01 | 2.332e+01 |
| Zj0b_c | 1.767e+02 | 1.287e+02 | 1.281e+02 | 7.226e+01 | 1.404e+01 | 1.945e+01 |
| Zj0b_udsg | 1.839e+02 | 1.240e+02 | 1.310e+02 | 2.404e+01 | 2.358e+01 | 6.320e+00 |
| Zj1b | 2.375e+02 | 1.283e+02 | 1.280e+02 | 1.230e+02 | 1.480e+01 | 3.053e+01 |
| Zj2b | 4.422e+02 | 2.845e+02 | 2.818e+02 | 1.635e+02 | 3.867e+01 | 6.779e+01 |
| s_Top | 2.330e+02 | 2.489e+02 | 9.822e+01 | 6.799e+01 | 3.088e+00 | 1.571e+01 |

Table C.22: The observed and predicted yields are given for the control regions for 0-lepton in 2018.

| Process | Z+udcsg enr., med. $p_T(V)$ | Z+udcsg enr., high $p_T(V)$ | Z+udcsg enr., high $p_T(V)$, boosted |
|-----------|-----------------------------------|-----------------------------------|---|
| Data | 12379 | 4990 | 95195 |
| TT | 1.587e+03 | 4.039e+02 | 4.111e+03 |
| VVHF | 1.466e+01 | 9.073e+00 | 4.532e+01 |
| VVLF | 2.011e+02 | 1.101e+02 | 2.118e+03 |
| Wj0b_c | 1.154e+03 | 2.958e+02 | 3.420e+03 |
| Wj0b_udsg | 3.383e+03 | 1.011e+03 | 3.105e+04 |
| Wj1b | 8.136e+01 | 2.965e+01 | 3.103e+02 |
| Wj2b | 4.687e+01 | 1.791e+01 | 1.862e+02 |
| Zj0b_c | 1.104e+03 | 4.583e+02 | 4.859e+03 |
| Zj0b_udsg | 2.777e+03 | 1.398e+03 | 4.649e+04 |
| Zj1b | 2.088e+02 | 8.279e+01 | 9.499e+02 |
| Zj2b | 9.193e+01 | 4.756e+01 | 3.538e+02 |
| s_Top | 2.053e+02 | 5.327e+01 | 6.468e+02 |
| Process | Z+bb enr., med. $p_T(V)$ | Z+bb enr., high $p_T(V)$ | Z+bb enr., high $p_T(V)$, boosted |
| Data | 15987 | 4868 | 2646 |
| TT | 7.015e+03 | 1.555e+03 | 9.006e+02 |
| VVHF | 1.357e+02 | 5.211e+01 | 4.142e+01 |
| VVLF | 2.933e+01 | 4.993e+01 | 3.335e+01 |
| Wj0b_c | 7.221e+02 | 1.563e+02 | 1.113e+02 |
| Wj0b_udsg | 1.067e+03 | 3.095e+02 | 4.882e+01 |
| Wj1b | 3.420e+02 | 9.114e+01 | 9.119e+01 |
| Wj2b | 6.378e+02 | 1.949e+02 | 2.111e+02 |
| Zj0b_c | 7.108e+02 | 2.549e+02 | 1.528e+02 |
| Zj0b_udsg | 7.362e+02 | 3.417e+02 | 7.935e+01 |
| Zj1b | 8.520e+02 | 2.848e+02 | 3.014e+02 |
| Zj2b | 1.542e+03 | 5.560e+02 | 3.863e+02 |
| s_Top | 8.532e+02 | 2.150e+02 | 9.127e+01 |
| Process | $t\bar{t}$ enr., med. $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$, boosted |
| Data | 3831 | 1152 | 2434 |
| TT | 3.488e+03 | 8.786e+02 | 2.212e+03 |
| VVHF | 4.540e+00 | 2.031e+00 | 7.929e+00 |
| VVLF | 3.735e+00 | 4.366e+00 | 3.684e+00 |
| Wj0b_c | 5.369e+01 | 1.926e+01 | 1.224e+01 |
| Wj0b_udsg | 7.749e+01 | 3.150e+01 | 3.779e+00 |
| Wj1b | 2.112e+01 | 7.183e+00 | 1.436e+01 |
| Wj2b | 2.933e+01 | 1.315e+01 | 4.750e+01 |
| Zj0b_c | 3.192e+01 | 1.352e+01 | 1.055e+01 |
| Zj0b_udsg | 3.032e+01 | 1.835e+01 | 3.877e+00 |
| Zj1b | 3.061e+01 | 1.428e+01 | 2.507e+01 |
| Zj2b | 5.670e+01 | 2.657e+01 | 1.103e+02 |
| s_Top | 1.634e+02 | 5.835e+01 | 1.153e+02 |

Table C.23: The observed and predicted yields are given for the signal regions for 1-lepton (e) in 2018.

| Process | Medium $p_T(V)$ | High $p_T(V)$ | High $p_T(V)$, boosted | Highest $p_T(V)$ | Highest $p_T(V)$, boosted |
|------------|--------------------|---------------|----------------------------|---------------------|----------------------------------|
| Data | 23105 | 2999 | 2362 | 235 | 526 |
| WH 0_75 | 1.643e-02 | 0 | 0 | 0 | 0 |
| WH 150_250 | 4.007e+01 | 2.263e+00 | 1.707e-02 | 0 | 0 |
| 0J | | | | | |
| WH 150_250 | 1.485e+01 | 9.443e-01 | 1.037e-01 | 0 | 0 |
| GE1J | | | | | |
| WH 250_400 | 7.017e+00 | 1.968e+01 | 3.252e+00 | 5.046e-01 | 1.492e-01 |
| WH 75_150 | 9.732e+00 | 1.254e-03 | 0 | 0 | 0 |
| WH GT400 | 4.116e-03 | 1.601e+00 | 5.061e-01 | 4.142e+00 | 3.202e+00 |
| ZH 0_75 | 2.667e-03 | 0 | 0 | 0 | 0 |
| ZH 150_250 | 1.064e+00 | 3.187e-02 | 0 | 0 | 0 |
| 0J | | | | | |
| ZH 150_250 | 3.764e-01 | 1.110e-02 | 2.182e-03 | 0 | 0 |
| GE1J | | | | | |
| ZH 250_400 | 3.517e-01 | 2.618e-01 | 6.275e-02 | 1.922e-03 | 9.850e-04 |
| ZH 75_150 | 4.193e-01 | 0 | 0 | 0 | 0 |
| ZH GT400 | 1.888e-02 | 7.269e-02 | 3.960e-02 | 4.452e-02 | 3.022e-02 |
| TT | 1.720e+04 | 1.930e+03 | 1.885e+03 | 8.391e+01 | 3.460e+02 |
| VVHF | 1.283e+02 | 3.503e+01 | 9.498e+00 | 1.983e+00 | 2.262e+00 |
| VVLF | 6.941e+01 | 1.886e+01 | 1.829e+01 | 7.554e+00 | 1.191e+01 |
| Wj0b_c | 1.043e+03 | 2.080e+02 | 8.942e+01 | 2.483e+01 | 2.735e+01 |
| Wj0b_udsg | 8.461e+02 | 2.037e+02 | 2.526e+01 | 3.201e+01 | 6.577e+00 |
| Wj1b | 4.374e+02 | 8.371e+01 | 6.410e+01 | 1.018e+01 | 1.773e+01 |
| Wj2b | 6.222e+02 | 2.029e+02 | 1.624e+02 | 3.434e+01 | 6.436e+01 |
| Zj0b_c | 4.593e+01 | 8.185e+00 | 2.330e+00 | 3.867e-01 | 6.397e-01 |
| Zj0b_udsg | 9.769e+01 | 1.024e+01 | 7.990e-01 | 5.467e-01 | 2.590e-01 |
| Zj1b | 9.190e+01 | 5.413e+00 | 4.972e+00 | 5.891e-01 | 7.025e-01 |
| Zj2b | 8.387e+01 | 8.816e+00 | 5.940e+00 | 7.114e-01 | 1.708e+00 |
| s_Top | 2.429e+03 | 4.121e+02 | 2.601e+02 | 3.611e+01 | 1.119e+02 |

Table C.24: The observed and predicted yields are given for the control regions for 1-lepton (e) in 2018.

| Process | W+udcsg enr., med. $p_T(V)$ | W+udcsg enr., high $p_T(V)$ | W+udcsg enr., high $p_T(V)$, boosted |
|-----------|--------------------------------|--------------------------------|---|
| Data | 193770 | 41572 | 57462 |
| TT | 2.880e+04 | 6.670e+03 | 8.374e+03 |
| VVHF | 6.427e+01 | 2.215e+01 | 2.635e+01 |
| VVLF | 3.082e+03 | 1.176e+03 | 2.157e+03 |
| Wj0b_c | 3.520e+04 | 6.690e+03 | 6.644e+03 |
| Wj0b_udsg | 1.040e+05 | 2.452e+04 | 4.129e+04 |
| Wj1b | 1.482e+03 | 3.708e+02 | 4.111e+02 |
| Wj2b | 4.340e+02 | 1.822e+02 | 2.725e+02 |
| Zj0b_c | 1.062e+03 | 1.163e+02 | 1.662e+02 |
| Zj0b_udsg | 6.537e+03 | 7.505e+02 | 1.606e+03 |
| Zj1b | 2.113e+02 | 2.135e+01 | 3.017e+01 |
| Zj2b | 2.967e+01 | 5.638e+00 | 1.102e+01 |
| s_Top | 4.710e+03 | 1.258e+03 | 1.570e+03 |

| Process | W+bb enr., med. $p_T(V)$ | W+bb enr., high $p_T(V)$ | W+bb enr., high $p_T(V)$, boosted |
|-----------|-----------------------------|-----------------------------|---------------------------------------|
| Data | 39928 | 6587 | 2565 |
| TT | 2.817e+04 | 4.350e+03 | 1.880e+03 |
| VVHF | 1.187e+02 | 3.467e+01 | 1.789e+01 |
| VVLF | 1.409e+02 | 5.294e+01 | 5.436e+01 |
| Wj0b_c | 2.453e+03 | 4.259e+02 | 1.857e+02 |
| Wj0b_udsg | 2.138e+03 | 4.823e+02 | 5.293e+01 |
| Wj1b | 9.698e+02 | 1.885e+02 | 1.152e+02 |
| Wj2b | 1.385e+03 | 3.772e+02 | 3.070e+02 |
| Zj0b_c | 9.179e+01 | 6.610e+00 | 9.481e+00 |
| Zj0b_udsg | 1.441e+02 | 2.227e+01 | 3.278e+00 |
| Zj1b | 1.454e+02 | 1.271e+01 | 1.042e+01 |
| Zj2b | 1.333e+02 | 1.608e+01 | 1.189e+01 |
| s_Top | 3.971e+03 | 8.775e+02 | 3.134e+02 |

| Process | $t\bar{t}$ enr., med. $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$, boosted |
|-----------|-----------------------------------|-----------------------------------|---|
| Data | 66941 | 15880 | 5432 |
| TT | 6.491e+04 | 1.531e+04 | 6.251e+03 |
| VVHF | 9.074e+01 | 4.928e+01 | 3.667e+00 |
| VVLF | 6.806e+01 | 4.098e+01 | 3.396e+00 |
| Wj0b_c | 4.724e+02 | 1.643e+02 | 1.100e+01 |
| Wj0b_udsg | 5.317e+02 | 1.664e+02 | 7.421e+00 |
| Wj1b | 9.104e+02 | 3.050e+02 | 1.347e+01 |
| Wj2b | 1.175e+03 | 4.533e+02 | 5.924e+01 |
| Zj0b_c | 3.027e+01 | 4.695e+00 | 6.660e-01 |
| Zj0b_udsg | 4.163e+01 | 8.545e+00 | 2.443e-01 |
| Zj1b | 1.101e+02 | 1.897e+01 | 1.634e+00 |
| Zj2b | 1.169e+02 | 2.323e+01 | 4.284e+00 |
| s_Top | 4.292e+03 | 1.636e+03 | 4.395e+02 |

Table C.25: The observed and predicted yields are given for the signal regions for 1-lepton (μ) in 2018.

| Process | Medium $p_T(V)$ | High $p_T(V)$ | High $p_T(V)$, boosted | Highest $p_T(V)$ | Highest $p_T(V)$, boosted |
|------------|--------------------|---------------|----------------------------|---------------------|----------------------------------|
| Data | 31579 | 4021 | 2960 | 294 | 599 |
| WH 0_75 | 2.262e-02 | 0 | 0 | 0 | 0 |
| WH 150_250 | 5.175e+01 | 2.568e+00 | 1.132e-02 | 0 | 0 |
| 0J | | | | | |
| WH 150_250 | 1.781e+01 | 1.101e+00 | 1.364e-01 | 0 | 0 |
| GE1J | | | | | |
| WH 250_400 | 8.456e+00 | 2.355e+01 | 3.771e+00 | 7.492e-01 | 1.761e-01 |
| WH 75_150 | 1.357e+01 | 3.760e-03 | 0 | 0 | 0 |
| WH GT400 | 1.499e-03 | 1.937e+00 | 7.714e-01 | 5.590e+00 | 3.732e+00 |
| ZH 0_75 | 2.724e-03 | 0 | 0 | 0 | 0 |
| ZH 150_250 | 1.726e+00 | 5.302e-02 | 7.135e-04 | 0 | 0 |
| 0J | | | | | |
| ZH 150_250 | 5.691e-01 | 2.210e-02 | 4.960e-03 | 0 | 0 |
| GE1J | | | | | |
| ZH 250_400 | 3.605e-01 | 3.943e-01 | 7.821e-02 | 4.068e-03 | 3.366e-04 |
| ZH 75_150 | 7.081e-01 | 0 | 0 | 0 | 0 |
| ZH GT400 | 1.129e-02 | 4.281e-02 | 2.207e-02 | 3.700e-02 | 2.988e-02 |
| TT | 2.256e+04 | 2.427e+03 | 2.312e+03 | 9.527e+01 | 4.177e+02 |
| VVHF | 1.619e+02 | 4.066e+01 | 1.123e+01 | 7.950e+00 | 8.388e+00 |
| VVLF | 1.393e+02 | 2.313e+01 | 8.864e+00 | 5.578e+00 | 7.667e+00 |
| Wj0b_c | 1.384e+03 | 2.563e+02 | 1.049e+02 | 2.844e+01 | 2.673e+01 |
| Wj0b_udsg | 1.348e+03 | 2.757e+02 | 1.980e+01 | 3.341e+01 | 1.150e+01 |
| Wj1b | 5.783e+02 | 1.126e+02 | 8.645e+01 | 1.518e+01 | 2.484e+01 |
| Wj2b | 7.780e+02 | 2.410e+02 | 2.095e+02 | 4.294e+01 | 8.545e+01 |
| Zj0b_c | 5.316e+01 | 7.564e+00 | 4.040e+00 | 4.491e-01 | 4.339e-01 |
| Zj0b_udsg | 1.592e+02 | 8.038e+00 | 1.700e+00 | 1.958e+00 | 1.576e-01 |
| Zj1b | 1.119e+02 | 7.940e+00 | 6.659e+00 | 4.733e-01 | 6.843e-01 |
| Zj2b | 1.115e+02 | 1.542e+01 | 9.168e+00 | 8.659e-01 | 1.446e+00 |
| s_Top | 3.203e+03 | 5.017e+02 | 3.558e+02 | 4.362e+01 | 1.380e+02 |

Table C.26: The observed and predicted yields are given for the control regions for 1-lepton (μ) in 2018.

| Process | W+udcsg enr., med. $p_T(V)$ | W+udcsg enr., high $p_T(V)$ | W+udcsg enr., high $p_T(V)$, boosted |
|-----------|--------------------------------|--------------------------------|---|
| Data | 260398 | 55094 | 69980 |
| TT | 3.728e+04 | 8.479e+03 | 1.010e+04 |
| VVHF | 8.490e+01 | 2.915e+01 | 2.662e+01 |
| VVLF | 4.182e+03 | 1.351e+03 | 2.737e+03 |
| Wj0b_c | 4.624e+04 | 8.475e+03 | 8.310e+03 |
| Wj0b_udsg | 1.414e+05 | 3.232e+04 | 5.274e+04 |
| Wj1b | 1.930e+03 | 4.689e+02 | 5.258e+02 |
| Wj2b | 5.624e+02 | 2.182e+02 | 3.405e+02 |
| Zj0b_c | 1.519e+03 | 1.611e+02 | 2.152e+02 |
| Zj0b_udsg | 9.581e+03 | 1.210e+03 | 2.167e+03 |
| Zj1b | 2.806e+02 | 3.073e+01 | 3.963e+01 |
| Zj2b | 5.045e+01 | 7.584e+00 | 1.454e+01 |
| s_Top | 6.177e+03 | 1.574e+03 | 1.927e+03 |

| Process | W+bb enr., med. $p_T(V)$ | W+bb enr., high $p_T(V)$ | W+bb enr., high $p_T(V)$, boosted |
|-----------|-----------------------------|-----------------------------|---------------------------------------|
| Data | 55450 | 8895 | 3283 |
| TT | 3.702e+04 | 5.489e+03 | 2.252e+03 |
| VVHF | 1.509e+02 | 4.565e+01 | 2.048e+01 |
| VVLF | 2.055e+02 | 5.641e+01 | 2.483e+01 |
| Wj0b_c | 3.287e+03 | 6.385e+02 | 2.670e+02 |
| Wj0b_udsg | 3.118e+03 | 6.437e+02 | 9.683e+01 |
| Wj1b | 1.317e+03 | 2.517e+02 | 1.480e+02 |
| Wj2b | 1.855e+03 | 4.605e+02 | 3.761e+02 |
| Zj0b_c | 1.474e+02 | 1.912e+01 | 7.336e+00 |
| Zj0b_udsg | 2.803e+02 | 3.444e+01 | 3.636e+00 |
| Zj1b | 1.958e+02 | 2.054e+01 | 1.189e+01 |
| Zj2b | 2.158e+02 | 2.033e+01 | 1.548e+01 |
| s_Top | 5.168e+03 | 1.149e+03 | 3.692e+02 |

| Process | $t\bar{t}$ enr., med. $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$, boosted |
|-----------|-----------------------------------|-----------------------------------|---|
| Data | 89639 | 20932 | 6748 |
| TT | 8.353e+04 | 1.917e+04 | 7.629e+03 |
| VVHF | 1.187e+02 | 6.113e+01 | 5.071e+00 |
| VVLF | 7.179e+01 | 4.845e+01 | 1.603e+00 |
| Wj0b_c | 5.961e+02 | 1.833e+02 | 1.732e+01 |
| Wj0b_udsg | 6.820e+02 | 2.045e+02 | 5.317e+00 |
| Wj1b | 1.166e+03 | 3.775e+02 | 1.699e+01 |
| Wj2b | 1.482e+03 | 5.833e+02 | 7.215e+01 |
| Zj0b_c | 3.751e+01 | 6.439e+00 | 6.920e-01 |
| Zj0b_udsg | 5.452e+01 | 1.295e+01 | 3.370e-01 |
| Zj1b | 1.458e+02 | 2.531e+01 | 1.742e+00 |
| Zj2b | 1.676e+02 | 3.272e+01 | 5.100e+00 |
| s_Top | 5.606e+03 | 2.052e+03 | 5.378e+02 |

Table C.27: The observed and predicted yields are given for the signal regions for 2-lepton (e) in 2018.

| Process | Low $p_T(V)$ | Medium $p_T(V)$ 0J | Medium $p_T(V)$ GE1J | High $p_T(V)$ | High $p_T(V)$, boosted | Highest $p_T(V)$ | Highest $p_T(V)$, boosted |
|-------------------------|-----------------|-----------------------|----------------------------|------------------|-------------------------------|---------------------|----------------------------------|
| Data | 4999 | 504 | 575 | 245 | 64 | 31 | 19 |
| ZH 0_75 | 2.289e-01 | 0 | 0 | 0 | 0 | 0 | 0 |
| ZH 150_250 0J | 3.373e-01 | 7.785e+00 | 9.678e-01 | 4.255e-02 | 8.011e-04 | 0 | 0 |
| ZH 150_250 GE1J | 1.493e-01 | 1.111e+00 | 3.269e+00 | 2.580e-02 | 3.602e-03 | 0 | 0 |
| ZH 250_400 | 0 | 1.249e-01 | 6.117e-02 | 4.417e+00 | 4.647e-01 | 1.842e-02 | 4.898e-03 |
| ZH 75_150 | 2.555e+01 | 1.220e-01 | 4.382e-02 | 0 | 0 | 0 | 0 |
| ZH GT400 | 0 | 0 | 0 | 3.598e-02 | 8.566e-03 | 8.884e-01 | 3.661e-01 |
| ggZH 0_75 | 2.594e-02 | 0 | 0 | 0 | 0 | 0 | 0 |
| ggZH 150_250 0J | 1.379e-01 | 2.646e+00 | 3.597e-01 | 9.620e-03 | 0 | 0 | 0 |
| ggZH 150_250 GE1J | 1.155e-01 | 1.113e+00 | 2.648e+00 | 1.633e-02 | 1.324e-03 | 0 | 0 |
| ggZH 250_400 | 0 | 3.244e-02 | 4.043e-02 | 1.483e+00 | 1.766e-01 | 2.691e-03 | 5.990e-04 |
| ggZH 75_150 | 7.824e+00 | 4.744e-02 | 3.488e-02 | 0 | 0 | 0 | 0 |
| ggZH GT400 | 0 | 0 | 0 | 8.888e-03 | 2.601e-03 | 1.293e-01 | 6.461e-02 |
| TT | 1.150e+03 | 2.658e+01 | 3.997e+01 | 1.226e+00 | 6.795e-01 | 0 | 2.631e-01 |
| VVHF | 5.039e+01 | 1.014e+01 | 8.163e+00 | 5.592e+00 | 6.454e-01 | 1.208e+00 | 6.631e-01 |
| VVLF | 1.260e+01 | 2.754e+00 | 2.359e+00 | 1.569e+00 | 1.225e+00 | 2.223e-01 | 4.583e-01 |
| Zj0b_c | 5.340e+02 | 6.238e+01 | 7.837e+01 | 2.303e+01 | 7.862e+00 | 3.571e+00 | 1.798e+00 |
| Zj0b_udsg | 7.475e+02 | 7.387e+01 | 1.212e+02 | 4.420e+01 | 1.816e+00 | 5.508e+00 | 6.458e-01 |
| Zj1b | 1.259e+03 | 1.188e+02 | 1.116e+02 | 3.898e+01 | 1.423e+01 | 3.843e+00 | 3.455e+00 |
| Zj2b | 1.167e+03 | 1.354e+02 | 1.368e+02 | 5.830e+01 | 2.181e+01 | 8.041e+00 | 7.796e+00 |
| s_Top | 4.040e+01 | 6.273e+00 | 3.225e+00 | 6.880e-01 | 0 | 0 | 0 |

Table C.28: The observed and predicted yields are given for the control regions for 2-lepton (e) in 2018.

| Process | Z+udcsg enr., low $p_T(V)$ | Z+udcsg enr., med. $p_T(V)$ | Z+udcsg enr., high $p_T(V)$ | Z+udcsg enr., high $p_T(V)$, boosted |
|-----------|----------------------------------|-----------------------------------|-----------------------------------|---|
| Data | 57826 | 11576 | 2368 | 1509 |
| TT | 9.427e+01 | 9.152e+00 | 2.976e-01 | 6.309e+00 |
| VVHF | 4.364e+00 | 1.404e+00 | 4.813e-01 | 1.243e+00 |
| VVLF | 2.698e+02 | 7.876e+01 | 2.135e+01 | 2.603e+01 |
| Zj0b_c | 3.314e+03 | 6.992e+02 | 1.307e+02 | 1.898e+02 |
| Zj0b_udsg | 6.430e+04 | 1.248e+04 | 2.380e+03 | 1.248e+03 |
| Zj1b | 5.998e+02 | 1.201e+02 | 2.142e+01 | 6.234e+01 |
| Zj2b | 3.336e+01 | 1.058e+01 | 3.179e+00 | 3.254e+01 |
| s_Top | 1.419e+01 | 2.919e+00 | 2.442e-01 | 1.097e+00 |
| Process | Z+bb enr., low $p_T(V)$ | Z+bb enr., med. $p_T(V)$ | Z+bb enr., high $p_T(V)$ | Z+bb enr., high $p_T(V)$, boosted |
| Data | 6249 | 1655 | 420 | 119 |
| TT | 7.093e+02 | 3.974e+01 | 7.926e-01 | 1.231e+00 |
| VVHF | 4.632e+01 | 1.621e+01 | 4.900e+00 | 2.484e+00 |
| VVLF | 9.093e+00 | 4.180e+00 | 1.246e+00 | 9.493e-01 |
| Zj0b_c | 8.230e+02 | 2.183e+02 | 5.327e+01 | 1.710e+01 |
| Zj0b_udsg | 1.066e+03 | 3.136e+02 | 8.774e+01 | 7.434e+00 |
| Zj1b | 1.748e+03 | 3.791e+02 | 7.215e+01 | 3.246e+01 |
| Zj2b | 1.655e+03 | 4.718e+02 | 1.008e+02 | 5.050e+01 |
| s_Top | 3.355e+01 | 6.414e+00 | 0 | 4.955e-01 |
| Process | $t\bar{t}$ enr., low $p_T(V)$ | $t\bar{t}$ enr., med. $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$, boosted |
| Data | 19584 | 1953 | 199 | 68 |
| TT | 1.964e+04 | 1.878e+03 | 1.694e+02 | 3.396e+01 |
| VVHF | 1.091e+01 | 4.021e+00 | 1.113e+00 | 6.643e-01 |
| VVLF | 6.835e+00 | 1.387e+00 | 3.596e+00 | 2.395e-01 |
| Zj0b_c | 7.226e+01 | 2.006e+01 | 2.272e+00 | 5.338e+00 |
| Zj0b_udsg | 6.501e+01 | 2.096e+01 | 2.974e+00 | 1.732e+00 |
| Zj1b | 2.739e+02 | 4.585e+01 | 8.711e+00 | 5.616e+00 |
| Zj2b | 2.836e+02 | 6.862e+01 | 1.788e+01 | 1.034e+01 |
| s_Top | 7.100e+02 | 1.608e+02 | 2.780e+01 | 1.576e+01 |

Table C.29: The observed and predicted yields are given for the signal regions for 2-lepton (μ) in 2018.

| Process | Low $p_T(V)$ | Medium $p_T(V)$ 0J | Medium $p_T(V)$ GE1J | High $p_T(V)$ | High $p_T(V)$, boosted | Highest $p_T(V)$ | Highest $p_T(V)$, boosted |
|-------------------------|-----------------|-----------------------|----------------------------|------------------|-------------------------------|---------------------|----------------------------------|
| Data | 7331 | 705 | 783 | 295 | 70 | 31 | 35 |
| ZH 0_75 | 2.889e-01 | 0 | 0 | 0 | 0 | 0 | 0 |
| ZH 150_250 0J | 4.431e-01 | 1.005e+01 | 1.257e+00 | 1.034e-01 | 2.004e-04 | 0 | 0 |
| ZH 150_250 GE1J | 1.992e-01 | 1.453e+00 | 4.195e+00 | 7.381e-02 | 3.357e-03 | 0 | 0 |
| ZH 250_400 | 0 | 1.758e-01 | 1.022e-01 | 5.524e+00 | 5.845e-01 | 5.370e-02 | 1.236e-02 |
| ZH 75_150 | 3.444e+01 | 2.071e-01 | 8.425e-02 | 0 | 0 | 0 | 0 |
| ZH GT400 | 0 | 0 | 0 | 6.876e-02 | 2.680e-02 | 1.108e+00 | 5.466e-01 |
| ggZH 0_75 | 3.529e-02 | 0 | 2.525e-04 | 0 | 0 | 0 | 0 |
| ggZH 150_250 0J | 1.599e-01 | 3.375e+00 | 4.952e-01 | 2.391e-02 | 3.038e-04 | 0 | 0 |
| ggZH 150_250 GE1J | 1.533e-01 | 1.427e+00 | 3.348e+00 | 4.348e-02 | 3.056e-03 | 0 | 0 |
| ggZH 250_400 | 0 | 5.089e-02 | 6.001e-02 | 1.751e+00 | 2.287e-01 | 1.206e-02 | 1.561e-03 |
| ggZH 75_150 | 1.022e+01 | 8.581e-02 | 5.159e-02 | 0 | 0 | 0 | 0 |
| ggZH GT400 | 0 | 0 | 0 | 1.904e-02 | 3.315e-03 | 1.617e-01 | 7.121e-02 |
| TT | 1.591e+03 | 4.376e+01 | 5.171e+01 | 3.347e+00 | 2.998e+00 | 0 | 0 |
| VVHF | 6.999e+01 | 1.453e+01 | 1.083e+01 | 6.557e+00 | 1.277e+00 | 1.198e+00 | 8.875e-01 |
| VVLF | 2.145e+01 | 2.783e+00 | 2.816e+00 | 1.973e+00 | 1.236e+00 | 1.780e-01 | 7.260e-01 |
| Zj0b_c | 7.340e+02 | 8.511e+01 | 8.904e+01 | 4.224e+01 | 8.671e+00 | 4.209e+00 | 3.089e+00 |
| Zj0b_udsg | 8.845e+02 | 9.877e+01 | 1.409e+02 | 5.813e+01 | 2.451e+00 | 8.230e+00 | 1.108e+00 |
| Zj1b | 1.815e+03 | 1.606e+02 | 1.462e+02 | 4.960e+01 | 1.742e+01 | 5.261e+00 | 4.598e+00 |
| Zj2b | 1.570e+03 | 1.734e+02 | 1.710e+02 | 7.151e+01 | 2.678e+01 | 1.001e+01 | 1.118e+01 |
| s_Top | 6.838e+01 | 5.039e+00 | 3.626e+00 | 0 | 0 | 0 | 0 |

Table C.30: The observed and predicted yields are given for the control regions for 2-lepton (μ) in 2018.

| Process | Z+udcsg enr., low $p_T(V)$ | Z+udcsg enr., med. $p_T(V)$ | Z+udcsg enr., high $p_T(V)$ | Z+udcsg enr., high $p_T(V)$, boosted |
|-----------|----------------------------------|-----------------------------------|-----------------------------------|---|
| Data | 81615 | 15434 | 3057 | 1976 |
| TT | 1.535e+02 | 1.872e+01 | 1.412e+00 | 1.317e+01 |
| VVHF | 6.382e+00 | 1.728e+00 | 2.966e-01 | 1.749e+00 |
| VVLF | 3.682e+02 | 1.020e+02 | 2.735e+01 | 2.809e+01 |
| Zj0b_c | 4.466e+03 | 8.822e+02 | 1.574e+02 | 2.460e+02 |
| Zj0b_udsg | 8.488e+04 | 1.594e+04 | 3.004e+03 | 1.597e+03 |
| Zj1b | 8.033e+02 | 1.509e+02 | 2.763e+01 | 7.948e+01 |
| Zj2b | 4.713e+01 | 1.298e+01 | 3.389e+00 | 4.048e+01 |
| s_Top | 2.289e+01 | 1.760e+00 | 7.158e-01 | 1.158e+00 |
| Process | Z+bb enr., low $p_T(V)$ | Z+bb enr., med. $p_T(V)$ | Z+bb enr., high $p_T(V)$ | Z+bb enr., high $p_T(V)$, boosted |
| Data | 9330 | 2258 | 495 | 149 |
| TT | 9.774e+02 | 6.429e+01 | 3.166e+00 | 3.249e+00 |
| VVHF | 6.060e+01 | 1.973e+01 | 6.293e+00 | 3.539e+00 |
| VVLF | 1.244e+01 | 4.716e+00 | 1.767e+00 | 1.391e+00 |
| Zj0b_c | 1.058e+03 | 2.945e+02 | 6.315e+01 | 2.417e+01 |
| Zj0b_udsg | 1.455e+03 | 3.920e+02 | 1.054e+02 | 9.642e+00 |
| Zj1b | 2.452e+03 | 4.951e+02 | 9.026e+01 | 4.420e+01 |
| Zj2b | 2.276e+03 | 6.077e+02 | 1.276e+02 | 6.493e+01 |
| s_Top | 4.292e+01 | 7.726e+00 | 2.201e-01 | 0 |
| Process | $t\bar{t}$ enr., low $p_T(V)$ | $t\bar{t}$ enr., med. $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$ | $t\bar{t}$ enr., high $p_T(V)$, boosted |
| Data | 28843 | 2891 | 333 | 112 |
| TT | 2.677e+04 | 2.490e+03 | 2.344e+02 | 3.955e+01 |
| VVHF | 1.455e+01 | 5.380e+00 | 1.617e+00 | 1.020e+00 |
| VVLF | 1.352e+01 | 4.143e-01 | 2.265e+00 | 6.160e-01 |
| Zj0b_c | 1.083e+02 | 2.056e+01 | 7.434e+00 | 8.186e+00 |
| Zj0b_udsg | 1.037e+02 | 2.296e+01 | 6.708e+00 | 5.635e+00 |
| Zj1b | 4.156e+02 | 7.519e+01 | 1.594e+01 | 1.082e+01 |
| Zj2b | 4.631e+02 | 1.288e+02 | 3.064e+01 | 2.091e+01 |
| s_Top | 9.704e+02 | 2.096e+02 | 3.764e+01 | 2.076e+01 |

Appendix D

Nuisance Parameter Impacts

The following figures show the impacts of the 60 most influential nuisance parameters on each STXS bin. There are 1878 nuisance parameters, but the influence of each on the final results quickly drops off. Notice the difference in the scale of each canvas in the figures.




Figure D-1: Impacts on WH $p_T(V) > 400$ GeV by the 60 most influential nuisance parameters.





Figure D-2: Impacts on WH $250 < p_T(V) < 400$ GeV by the 60 most influential nuisance parameters.





Figure D-3: Impacts on WH $150 < p_T(V) < 250$ GeV by the 60 most influential nuisance parameters.




Figure D-4: Impacts on ZH $p_T(V) > 400$ GeV by the 60 most influential nuisance parameters.





Figure D-5: Impacts on ZH $250 < p_T(V) < 400$ GeV by the 60 most influential nuisance parameters.




Figure D-6: Impacts on ZH 150 < $p_T(V)$ < 250 GeV with additional jets by the 60 most influential nuisance parameters.




Figure D-7: Impacts on ZH 150 < $p_T(V)$ < 250 GeV without additional jets by the 60 most influential nuisance parameters.





Figure D-8: Impacts on ZH $75 < p_T(V) < 150$ GeV by the 60 most influential nuisance parameters.

Bibliography

- [1] T. S. Kuhn, *The structure of scientific revolutions*. Chicago: University of Chicago Press, 1970.
- [2] J. Smith, W. L. van Neerven, and J. A. M. Vermaseren, “Transverse mass and width of the w boson,” *Phys. Rev. Lett.*, vol. 50, pp. 1738–1740, May 1983.
- [3] W. Dau, “Ua1 results from $p\bar{p}$ collisions at $\sqrt{s} = 540$ gev,” tech. rep., 198401126, 1983.
- [4] S. Chatrchyan, V. Khachatryan, A. Sirunyan, A. Tumasyan, W. Adam, E. Aguilo, T. Bergauer, M. Dragicevic, J. Erö, C. Fabjan, *et al.*, “Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC,” *Physics Letters B*, vol. 716, p. 30–61, Sep 2012.
- [5] G. Aad *et al.*, “Combined search for the Standard Model Higgs boson in pp collisions at $\sqrt{s}=7$ TeV with the ATLAS detector,” *Phys. Rev. D*, vol. 86, p. 032003, Aug 2012.
- [6] CMS Collaboration, “Observation of Higgs Boson Decay to Bottom Quarks,” *Physical Review Letters*, vol. 121, Sep 2018.
- [7] M. Aaboud, G. Aad, B. Abbott, O. Abdinov, B. Abeloos, D. Abhayasinghe, S. Abidi, O. AbouZeid, N. Abraham, H. Abramowicz, and *et al.*, “Observation of $h \rightarrow b\bar{b}$ decays and vh production with the atlas detector,” *Physics Letters B*, vol. 786, p. 59–86, Nov 2018.
- [8] M. Thomson, *Modern particle physics*. Cambridge: Cambridge University Press, 2013.
- [9] S. Brandt, *Measurement of W and Z boson production cross sections in proton-proton collisions at $\sqrt{s} = 5.02$ TeV and $\sqrt{s} = 13$ TeV*. PhD thesis, Massachusetts Institute of Technology, 2020.
- [10] D. J. Gross and F. Wilczek, “Ultraviolet behavior of non-abelian gauge theories,” *Phys. Rev. Lett.*, vol. 30, pp. 1343–1346, Jun 1973.
- [11] S. Weinberg, “A model of leptons,” *Phys. Rev. Lett.*, vol. 19, pp. 1264–1266, Nov 1967.

- [12] F. Englert and R. Brout, “Broken symmetry and the mass of gauge vector mesons,” *Phys. Rev. Lett.*, vol. 13, pp. 321–323, Aug 1964.
- [13] P. W. Higgs, “Broken symmetries and the masses of gauge bosons,” *Phys. Rev. Lett.*, vol. 13, pp. 508–509, Oct 1964.
- [14] G. S. Guralnik, C. R. Hagen, and T. W. B. Kibble, “Global conservation laws and massless particles,” *Phys. Rev. Lett.*, vol. 13, pp. 585–587, Nov 1964.
- [15] V. Fock, “Bemerkung zum Virialsatz,” *Zeitschrift für Physik*, vol. 63, pp. 855–858, Nov. 1930.
- [16] CMS Collaboration, *CMS Physics: Technical Design Report Volume 1: Detector Performance and Software*. Technical Design Report CMS, Geneva: CERN, 2006.
- [17] L. Evans and P. Bryant, “LHC machine,” *Journal of Instrumentation*, vol. 3, pp. S08001–S08001, aug 2008.
- [18] R. Bruce, G. Arduini, H. Bartosik, R. De Maria, M. Giovannozzi, G. Iadarola, J. Jowett, K. S. B. Li, M. Lamont, A. Lechner, E. Metral, D. Mirarchi, T. Pieloni, S. Redaelli, G. Rumolo, B. Salvant, R. Tomas Garcia, and J. Wenninger, “LHC Run 2: Results and challenges,” p. MOAM5P50. 7 p, Jul 2016.
- [19] D. Barney, “CMS Detector Slice.” CMS Collection., Jan 2016.
- [20] A. Dominguez, D. Abbaneo, K. Arndt, N. Bacchetta, A. Ball, E. Bartz, W. Bertl, G. M. Bilei, G. Bolla, H. W. K. Cheung, M. Chertok, S. Costa, N. Demaria, D. D. Vazquez, K. Ecklund, W. Erdmann, K. Gill, G. Hall, K. Harder, F. Hartmann, R. Horisberger, W. Johns, H. C. Kaestli, K. Klein, D. Kotlinski, S. Kwan, M. Pesaresi, H. Postema, T. Rohe, C. Schäfer, A. Starodumov, S. Streuli, A. Tricomi, P. Tropea, J. Troska, F. Vasey, and W. Zeuner, “CMS Technical Design Report for the Pixel Detector Upgrade,” Tech. Rep. CERN-LHCC-2012-016. CMS-TDR-11, Sep 2012. Additional contacts: Jeffrey Spalding, Fermilab, Jeffrey.Spalding@cern.ch Didier Contardo, Universite Claude Bernard-Lyon I, didier.claude.contardo@cern.ch.
- [21] A. Modak, “CMS Phase-1 Pixel Detector: Operational Experience, Performance and Lessons Learned,” Tech. Rep. CMS-CR-2019-283, CERN, Geneva, Nov 2019.
- [22] F. Monti, “CMS ECAL monitoring and calibration in LHC Run 2,” Tech. Rep. CMS-CR-2018-171, CERN, Geneva, Aug 2018.
- [23] N. Bartosik, “Performance of the CMS Electromagnetic Calorimeter in LHC Run2,” *PoS*, vol. LeptonPhoton2019, p. 126. 4 p, 2019.
- [24] M. Chadeeva and N. Lychkovskaya, “Calibration of the CMS hadron calorimeter in run 2,” *Journal of Instrumentation*, vol. 13, pp. C03025–C03025, mar 2018.

- [25] E. James, Y. Maravin, M. Mulders, and N. Neumeister, “Muon Identification in CMS,” Tech. Rep. CMS-NOTE-2006-010, CERN, Geneva, Jan 2006.
- [26] V. I. Klioukhine, D. Campi, B. Cure, A. Desirelli, S. Farinon, H. Gerwig, D. Green, J. P. Grillet, A. Herve, F. Kircher, B. Levesy, R. Loveless, and R. P. Smith, “3d magnetic analysis of the cms magnet,” *IEEE Transactions on Applied Superconductivity*, vol. 10, no. 1, pp. 428–431, 2000.
- [27] V. Klyukhin, N. Amapane, A. Ball, B. Curé, A. Gaddi, H. Gerwig, M. Mulders, V. Calvelli, A. Hervé, and R. Loveless, “Validation of the CMS Magnetic Field Map. Validation of the CMS Magnetic Field Map,” *J. Supercond. Novel Magn.*, vol. 28, pp. 701–704. 7 p, May 2016.
- [28] N. Pozzobon, “The CMS muon system: performance during the LHC run-2,” *Journal of Instrumentation*, vol. 14, pp. C11031–C11031, nov 2019.
- [29] A. Sirunyan, A. Tumasyan, W. Adam, E. Asilar, T. Bergauer, J. Brandstetter, E. Brondolin, M. Dragicevic, J. Erö, M. Flechl, and et al., “Particle-flow reconstruction and global event description with the cms detector,” *Journal of Instrumentation*, vol. 12, p. P10003–P10003, Oct 2017.
- [30] M. Tosi, “The CMS trigger in Run 2,” Tech. Rep. CMS-CR-2017-340, CERN, Geneva, Oct 2017.
- [31] L. Cadamuro, “The CMS level-1 trigger system for LHC run II,” *Journal of Instrumentation*, vol. 12, pp. C03021–C03021, mar 2017.
- [32] H. Sert, “CMS High Level Trigger Performance in Run 2,” Tech. Rep. CMS-CR-2019-278, CERN, Geneva, Nov 2019.
- [33] M. E. Peskin and D. V. Schroeder, *An introduction to quantum field theory*. Boulder, CO: Westview, 1995. Includes exercises.
- [34] C. Oleari, “The powheg box,” *Nuclear Physics B - Proceedings Supplements*, vol. 205-206, p. 36–41, Aug 2010.
- [35] V. Hirschi and O. Mattelaer, “Automated event generation for loop-induced processes,” 2015.
- [36] B. Andersson, G. Gustafson, G. Ingelman, and T. Sjöstrand, “Parton fragmentation and string dynamics,” *Physics Reports*, vol. 97, no. 2, pp. 31 – 145, 1983.
- [37] T. Sjöstrand, S. Ask, J. R. Christiansen, R. Corke, N. Desai, P. Ilten, S. Mrenna, S. Prestel, C. O. Rasmussen, and P. Z. Skands, “An introduction to pythia 8.2,” *Computer Physics Communications*, vol. 191, pp. 159 – 177, 2015.
- [38] S. Agostinelli *et al.*, “Geant4—a simulation toolkit,” *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment*, vol. 506, no. 3, pp. 250 – 303, 2003.

- [39] M. Hildreth, V. N. Ivanchenko, D. J. Lange, and M. J. Kortelainen, “CMS full simulation for run-2,” *Journal of Physics: Conference Series*, vol. 664, p. 072022, dec 2015.
- [40] V. Lefébure, S. Banerjee, and I. González, “CMS Simulation Software Using Geant4,” Tech. Rep. CMS-NOTE-1999-072, CERN, Geneva, Dec 1999.
- [41] K. Bloom, “Cms software and computing for lhcb run 2,” *Proceedings of 38th International Conference on High Energy Physics — PoS(ICHEP2016)*, Feb 2017.
- [42] N. Ratnikova, C.-H. Huang, A. Sanchez-Hernandez, T. Wildish, and X. Zhang, “CMS space monitoring,” *Journal of Physics: Conference Series*, vol. 513, p. 042036, jun 2014.
- [43] V. Innocente, L. Silvestris, and D. Stickland, “Cms software architecture: Software framework, services and persistency in high level trigger, reconstruction and analysis,” *Computer Physics Communications*, vol. 140, no. 1, pp. 31 – 44, 2001. CHEP2000.
- [44] K. Rose, “Deterministic annealing for clustering, compression, classification, regression, and related optimization problems,” *Proceedings of the IEEE*, vol. 86, no. 11, pp. 2210–2239, 1998.
- [45] CMS Collaboration, “Description and performance of track and primary-vertex reconstruction with the CMS tracker,” *Journal of Instrumentation*, vol. 9, pp. P10009–P10009, oct 2014.
- [46] J. R. and, “CMS electron and photon performance at 13 TeV,” *Journal of Physics: Conference Series*, vol. 1162, p. 012008, jan 2019.
- [47] M. Cacciari, G. P. Salam, and G. Soyez, “The anti-ktjet clustering algorithm,” *Journal of High Energy Physics*, vol. 2008, p. 063–063, Apr 2008.
- [48] V. Khachatryan, A. Sirunyan, A. Tumasyan, W. Adam, E. Asilar, T. Bergauer, J. Brandstetter, E. Brondolin, M. Dragicevic, J. Erö, and et al., “Jet energy scale and resolution in the cms experiment in pp collisions at 8 tev,” *Journal of Instrumentation*, vol. 12, p. P02014–P02014, Feb 2017.
- [49] A. Sirunyan, A. Tumasyan, W. Adam, F. Ambrogio, E. Asilar, T. Bergauer, J. Brandstetter, E. Brondolin, M. Dragicevic, J. Erö, and et al., “Identification of heavy-flavour jets with the cms detector in pp collisions at 13 tev,” *Journal of Instrumentation*, vol. 13, p. P05011–P05011, May 2018.
- [50] M. Abadi, A. Agarwal, P. Barham, E. Brevdo, Z. Chen, C. Citro, G. S. Corrado, A. Davis, J. Dean, M. Devin, S. Ghemawat, I. J. Goodfellow, A. Harp, G. Irving, M. Isard, Y. Jia, R. Józefowicz, L. Kaiser, M. Kudlur, J. Levenberg, D. Mané, R. Monga, S. Moore, D. G. Murray, C. Olah, M. Schuster, J. Shlens, B. Steiner, I. Sutskever, K. Talwar, P. A. Tucker, V. Vanhoucke, V. Vasudevan, F. B. Viégas,

- O. Vinyals, P. Warden, M. Wattenberg, M. Wicke, Y. Yu, and X. Zheng, “Tensorflow: Large-scale machine learning on heterogeneous distributed systems,” *CoRR*, vol. abs/1603.04467, 2016.
- [51] CMS Collaboration, “A deep neural network for simultaneous estimation of b jet energy and resolution,” 2019.
- [52] D. Abercrombie, “A search for dark matter via higgs decay using quark jet substructure,” 2014.
- [53] A. J. Larkoski, S. Marzani, G. Soyez, and J. Thaler, “Soft drop,” *Journal of High Energy Physics*, vol. 2014, May 2014.
- [54] The CMS collaboration, “Performance of the CMS missing transverse momentum reconstruction in pp data at $\sqrt{s} = 8$ TeV,” *Journal of Instrumentation*, vol. 10, pp. P02006–P02006, feb 2015.
- [55] A. M. Sirunyan, A. Tumasyan, W. Adam, F. Ambroggi, E. Asilar, T. Bergauer, J. Brandstetter, M. Dragicevic, J. Erö, and et al., “Search for production of higgs boson pairs in the four b quark final state using large-area jets in proton-proton collisions at $s = 13$ $\sqrt{s} = 13$ tev,” *Journal of High Energy Physics*, vol. 2019, Jan 2019.
- [56] The CMS Collaboration, *CMSSW Reference Manual*.
- [57] C. Kato, “Status and prospects of STXS measurements in ATLAS and CMS,” Tech. Rep. ATL-PHYS-PROC-2019-080, CERN, Geneva, Aug 2019.
- [58] “Muon reconstruction performance during Run II,” Jul 2019.
- [59] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H. S. Shao, T. Stelzer, P. Torrielli, and M. Zaro, “The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations,” *JHEP*, vol. 07, p. 079, 2014.
- [60] S. Kallweit, J. M. Lindert, P. Maierhöfer, S. Pozzorini, and M. Schönherr, “Nlo qcd+ew predictions for $v + \text{jets}$ including off-shell vector-boson decays and multi-jet merging,” *Journal of High Energy Physics*, vol. 2016, p. 1–51, Apr 2016.
- [61] D. E. De Florian Sabaris, C. Grojean, F. Maltoni, C. Mariotti, A. Nikitenko, M. Pieri, P. Savard, M. Schumacher, and R. Tanaka, “Handbook of LHC Higgs cross sections: 4. Deciphering the nature of the Higgs sector,” Sep 2016.
- [62] “CMS Luminosity Measurements for the 2016 Data Taking Period,” Tech. Rep. CMS-PAS-LUM-17-001, CERN, Geneva, 2017.
- [63] “CMS luminosity measurement for the 2017 data-taking period at $\sqrt{s} = 13$ TeV,” Tech. Rep. CMS-PAS-LUM-17-004, CERN, Geneva, 2018.

- [64] “CMS luminosity measurement for the 2018 data-taking period at $\sqrt{s} = 13$ TeV,” Tech. Rep. CMS-PAS-LUM-18-002, CERN, Geneva, 2019.
- [65] A. Sirunyan *et al.*, “Cross section measurement of t-channel single top quark production in pp collisions at $s=13$ tev,” *Physics Letters B*, vol. 772, pp. 752 – 776, 2017.
- [66] A. Sirunyan *et al.*, “Combination of searches for heavy resonances decaying to ww , wz , zz , wh , and zh boson pairs in proton–proton collisions at $s=8$ and 13 tev,” *Physics Letters B*, vol. 774, pp. 533 – 558, 2017.
- [67] R. Barlow and C. Beeston, “Fitting using finite monte carlo samples,” *Computer Physics Communications*, vol. 77, no. 2, pp. 219 – 228, 1993.
- [68] CMS Collaboration and others, “Documentation for the higgs combine tool.”
- [69] J. de Blas, M. Cepeda, J. D’Hondt, R. Ellis, C. Grojean, B. Heinemann, F. Maltoni, A. Nisati, E. Petit, R. Rattazzi, and et al., “Higgs boson studies at future particle colliders,” *Journal of High Energy Physics*, vol. 2020, Jan 2020.
- [70] “Search for the Standard Model Higgs Boson Decaying to Bottom Quarks and Produced in Association with a W or a Z Boson,” Tech. Rep. CMS-PAS-HIG-11-012, CERN, Geneva, 2011.
- [71] N. Berger, C. Bertella, T. P. Calvet, M. Calvetti, V. Dao, M. Delmastro, M. Duehrssen-Debling, P. Francavilla, Y. Haddad, O. Kivernyk, J. M. Langford, C. Li, G. Marchiori, P. Milenovic, C. E. Pandini, E. Scott, F. J. Tackmann, K. Tackmann, L. Viliani, and M. Xiao, “Simplified template cross sections - stage 1.1,” 2019.
- [72] M. Aaboud, G. Aad, B. Abbott, D. C. Abbott, O. Abdinov, A. Abed Abud, D. K. Abhayasinghe, S. H. Abidi, O. S. AbouZeid, and et al., “Measurement of VH , $H \rightarrow b\bar{b}$ production as a function of the vector-boson transverse momentum in 13 TeV pp collisions with the ATLAS detector,” *Journal of High Energy Physics*, vol. 2019, May 2019.
- [73] G. Apollinari, I. Béjar Alonso, O. Brüning, M. Lamont, and L. Rossi, “High-luminosity large hadron collider (hl-lhc) : Preliminary design report,” 12 2015.
- [74] T. C. S. Group, “Cepec conceptual design report: Volume 1 - accelerator,” 2018.
- [75] Y. Iiyama, B. Maier, D. Abercrombie, M. Goncharov, and C. Paus, “Dynamo – handling scientific data across sites and storage media,” 2020.
- [76] G. Van Rossum and F. L. Drake Jr, *Python reference manual*. Centrum voor Wiskunde en Informatica Amsterdam, 1995.
- [77] G. Van Rossum and F. L. Drake, *Python 3 Reference Manual*. Scotts Valley, CA: CreateSpace, 2009.

- [78] E. Laure, A. Edlund, F. Pacini, P. Buncic, M. Barroso, A. Di Meglio, F. Prelz, A. Frohner, O. Mulmo, A. Krenek, *et al.*, “Programming the grid with glite,” tech. rep., CERN, Geneva, Switzerland, 2006.
- [79] D. Abercrombie, *Dynamo Consistency*.
- [80] A. Dorigo, P. Elmer, F. Furano, and A. Hanushevsky, “Xrootd-a highly scalable architecture for data access,” *WSEAS Transactions on Computers*, vol. 1, no. 4.3, pp. 348–353, 2005.
- [81] D. Abercrombie and et al., *SmartDataProjects/dynamo-consistency*.
- [82] D. Abercrombie, *Site Admin Toolkit*.