

Série 3 Exercice 8

David Wiedemann

19 mars 2022

1

Indeed, let $\frac{a}{b} \in \mathbb{Q}$ in reduced form such that $\nu_p(\frac{a}{b}) = 0$. By the definition of p -adic, this means that we may suppose both a and b share no common factors with p , then $\frac{b}{a}$ also shares no common factor with p and hence $\nu_p(\frac{b}{a}) = 0$, implying $\frac{b}{a} \in R_\nu$.

Finally, $\frac{a}{b} \cdot \frac{b}{a} = \frac{1}{1}$ which finally implies that $\frac{a}{b}$ is invertible in R_ν .

2

First we show that all (p^n) are distinct ideals of R , indeed suppose there exists $a, b \in \mathbb{N}$ such that $(p^a) = (p^b)$, without loss of generality suppose $a < b$.

Hence, there exists an element $\frac{x}{y} \in \mathbb{Q}$ with $\nu_p(\frac{x}{y}) \geq 0$ such that $\frac{x}{y}p^b = p^a$.

As \mathbb{Q} is a field, this implies that $\frac{x}{y} = p^{a-b}$ which means $\frac{x}{y}$ has a negative valuation which contradicts our hypothesis.

Now we show that the ideals mentionned in the exercise are indeed all the ideals of R .

Let I be a non-zero ideal of R .

Define $a = \inf_{x \in I \setminus \{0\}} \{\nu(x)\}$. Since $\nu|_{I \setminus \{0\}}$ has codomain \mathbb{N} , this infimum exists and is attained by some element $y \in I$.

Note that we may write $y = p^a \frac{d}{c}$ where d and c are coprime to p .

By part 1, we know that $\frac{d}{c}$ is invertible, hence implying that (since I is an ideal) $p^a \in I$.

We pretend that $I = (p^a)$, to do this, we show the double inclusion.

First, note that, since by definition $p^a \in I$, we immediatly get that $(p^a) \subset I$ since (p^a) is the smallest ideal containing p^a .

Furthermore, let $x \in I$, then by definition of a , $\nu(x) \geq a$.

We may then write $x = p^{\nu(x)} \frac{d}{c} = p^a p^{\nu(x)-a} \frac{d}{c}$ where d and c are coprime to p , this implies that $x \in (p^a)$.

Hence, if I is a non-zero ideal, I is of the form p^n for some n and since these ideals are disjoint, we have characterised all of them.

3

Using the exercise of week 2, we know that $\mathbb{Z} \subset R$.

Hence consider the composition $\mathbb{Z} \xrightarrow{\iota} R \xrightarrow{q_R} R/(p^n)$ where ι is the inclusion morphism and q_R is the canonical projection morphism.

Furthermore define $q : \mathbb{Z} \rightarrow \mathbb{Z}/(p^n)$ to be the canonical projection.

We now pretend that $\ker(q_R \circ \iota) = \ker q = (p^n)$.

Indeed if $a \in \ker q = (p^n)$, then $p^n | a$ hence $p^n | \iota(a) \implies q_R(a) = 0$.

Similarly, if $r \in \ker(q_R \circ \iota)$, then $p^n | r$, ie. there exists $\frac{a}{b} \in R$ (in reduced form) such that $p^n \frac{a}{b} = r$ since $\nu(\frac{a}{b}) \geq 0$, in particular we may suppose b is coprime to p .

Hence, since $p^n \frac{a}{b}$ is an integer, $b | a$ implying $b = 1$.

Finally, this means that there exists an integer a such that $p^n a = r$ which means that $a \in (p^n) = \ker q$.

Hence applying the universal property of the quotient ring, we get an induced morphism as such :

$$\begin{array}{ccc} \mathbb{Z} & \xrightarrow{q_R \circ \iota} & R/(p^n) \\ q \downarrow & \nearrow \exists! \phi & \\ \mathbb{Z}/(p^n) & & \end{array}$$

We pretend that it is now sufficient to show that $q_R \circ \iota$ is surjective to show that ϕ is indeed an isomorphism.

Before showing that $q_R \circ \iota$ is surjective, we show how this implies ϕ is an isomorphism.

Indeed, suppose $q_R \circ \iota$ is surjective, since $\ker q_R \circ \iota = \ker q$ the universal property of the quotient implies that there exists a unique map $\psi : R/(p^n) \rightarrow \mathbb{Z}/(p^n)$ making the following diagram commute

$$\begin{array}{ccc} \mathbb{Z} & \xrightarrow{q} & \mathbb{Z}/(p^n) \\ q_R \circ \iota \downarrow & \nearrow \exists! \psi & \\ R/(p^n) & & \end{array}$$

Finally, we pretend that ψ is an inverse to ϕ , indeed, notice that the following

diagrams commute :

$$\begin{array}{ccc}
 R/(p^n) & & \mathbb{Z}/(p^n) \\
 \uparrow q_R \circ \iota & \nwarrow \phi & \\
 \mathbb{Z} & \xrightarrow{q} & \mathbb{Z}/(p^n) \\
 \downarrow q_R \circ \iota & \nearrow \psi & \\
 R/(p^n) & &
 \end{array}
 \qquad
 \begin{array}{ccc}
 \mathbb{Z}/(p^n) & & R/(p^n) \\
 \uparrow q & \nwarrow \psi & \\
 \mathbb{Z} & \xrightarrow{q_R \circ \iota} & R/(p^n) \\
 \downarrow q & \nearrow \phi & \\
 \mathbb{Z}/(p^n) & &
 \end{array}$$

Hence $q_R \circ \iota = \psi \circ \phi \circ q_R \circ \iota$ and a final application of the universal property of the quotient implies that $\psi \circ \phi = \text{Id}_{R/(p^n)}$.

This shows that $R/(p^n) \simeq \mathbb{Z}/(p^n)$.

We now show that $q_R \circ \iota$ is surjective.

Let $[p^i \frac{a}{b}] \in R/(p^n)$, where, as always, we have assumed $\frac{a}{b}$ is in reduced form and shares no factors with p .

To show this, we must find an integer $d \in \mathbb{Z}$ such that

$$\frac{a}{b} p^i - d = k p^n, \quad k \in R$$

where by abuse of notation, we regard d as included in R .

Indeed, then $q_R(\frac{a}{b} p^i) = q_R(d)$ which will imply that $q_R \circ \iota$ is surjective.

Using that p^n and b are coprime, choose x and d integers such that $x p^n + d b = a p^i$, such x and d always exist because of Bezout's theorem.

Now set $k = \frac{x}{b}$, note that $k \in R$ since b is coprime to p .

It is now immediatly verified that

$$\frac{a}{b} p^i - d = k p^n$$

Since

$$a p^i = k p^n b + d b = x p^n + d b$$

Hence $\frac{a}{b} p^i$ has a representative in \mathbb{Z} which in turn implies that $q_R \circ \iota$ is surjective, concluding our proof.

4

To show this, we proceed by contradiction.

So suppose there exist two different prime numbers $p \neq q$ such that $R_p \simeq R_q$. Note that an isomorphism of rings induces a bijection between the set of ideals which respects inclusion.

To declutter this proof, I prove this at the end of the document.

So let $\psi : R_p \rightarrow R_q$ be the isomorphism we assume to exist, using part 2, we know that all ideals of R_p are of the form (p^n) .

Furthermore, we notice that these ideals are nicely ordered :

$$(p) \supsetneq (p^2) \supsetneq \dots$$

Applying ϕ to this chain yields a chain of ideals in R_q

$$\phi((p)) \supsetneq \phi((p^2)) \supsetneq \dots$$

Using the result cited above, this immediatly implies that $\phi((p^i)) = (q^i)$. Since R_p and R_q are supposed isomorphic, we would need to have that $R_p/(p) \simeq R_q/(q)$ are isomorphic. Using part 3, this would imply that $\mathbb{Z}/(q) \simeq \mathbb{Z}/(p)$ which is a contradiction, since they are not even in bijection as sets.

We now show the result cited at the beginning of part 4.

So let $\phi : A \rightarrow B$ be an isomorphism of rings and let $I \subset A$ be an ideal, then $\phi(I)$ is an ideal since :

- Clearly $\phi(I)$ is an additive subgroup since $\phi(0) = 0 \in \phi(I)$ and $\phi(a) + \phi(b) = \phi(a + b) \in \phi(I)$.
- Let $\lambda \in B$ and $\phi(a) \in \phi(I)$, then $\lambda\phi(a) = \phi(\phi^{-1}(\lambda))\phi(a) = \phi(\underbrace{\phi^{-1}(\lambda)a}_{\in I}) \in \phi(I)$.

This correspondence is clearly a bijection between the ideals of A and B since ϕ is a bijection and the correspondence obviously preserves inclusions since maps of set preserve inclusions.