

Functional Analysis

David Wiedemann

Table des matières

| | | |
|----------|--|----------|
| 1 | Introduction | 3 |
| 1.1 | Topological Spaces | 3 |
| 1.2 | Metric spaces | 3 |
| 1.3 | Norms, Banach Spaces | 6 |
| 1.4 | Basis of a normed space | 7 |
| 1.5 | Scalar products and Hilbert spaces | 8 |

List of Theorems

| | | |
|----|---|---|
| 1 | Definition (Topological space) | 3 |
| 2 | Definition (Properties) | 3 |
| 3 | Definition (Sequences) | 3 |
| 4 | Definition (Continuity) | 3 |
| 5 | Definition (Metric space) | 3 |
| 6 | Definition | 4 |
| 1 | Lemma | 4 |
| 7 | Definition | 4 |
| 2 | Proposition | 4 |
| 3 | Theorem | 5 |
| 8 | Definition (Normed space) | 6 |
| 9 | Definition | 6 |
| 10 | Definition (Banach Space) | 6 |
| 4 | Proposition | 6 |
| 11 | Definition | 7 |
| 12 | Definition (Schauder Basis) | 7 |
| 13 | Definition (Equivalence of Norms) | 7 |
| 5 | Lemma | 7 |
| 14 | Definition | 8 |
| 7 | Proposition | 8 |
| 8 | Proposition | 8 |
| 15 | Definition (Hilbert Space) | 9 |

| | | |
|----|---|----|
| 9 | Lemma | 9 |
| 10 | Theorem | 9 |
| 11 | Lemma | 10 |
| 12 | Corollary | 10 |
| 16 | Definition (Orthonormal systemes) | 10 |
| 13 | Lemma | 10 |

1 Introduction

Lecture 1: Introduction

Wed 12 Oct

Main reference is "Functional Analysis" by H.W. Alt.

1.1 Topological Spaces

Definition 1 (Topological space)

Let X be a set, a topology is a subset $\tau \subset P(X)$ is a topology if

- $\emptyset, X \in \tau$
- any union of opens is open
- Finite intersections of opens are open.

Definition 2 (Properties)

For $A \subset X$, \bar{A} is the smallest closed set containing A and the interior A° is the biggest open set contained in A .

Finally, the boundary is $\partial A = \bar{A} \setminus A^\circ$.

X is separable if \exists a dense countable subset

Definition 3 (Sequences)

Let $x : \mathbb{N} \rightarrow X, \bar{x} \in X, \lim x_k = \bar{x} \iff$ any neighbourhood $U \in \tau$ of \bar{x} eventually contains x_k

Definition 4 (Continuity)

A function $f : X \rightarrow Y$ is continuous if $\forall U \in \tau_Y, f^{-1}(U) \in \tau_X$.

This is different from sequential continuity $x_n \rightarrow \bar{x} \implies f(x_n) \rightarrow f(\bar{x})$

.

f is continuous at $x \in X$ if $\forall V \in \tau_Y$ st $f(x) \in V \implies f^{-1}(V) \in \tau_X$

Lecture 2: More recaps

Fri 14 Oct

1.2 Metric spaces

Definition 5 (Metric space)

X a set, $d : X \times X \rightarrow [0, \infty)$ is a metric

Definition 6

X a set, d_1, d_2 metrics

1. d_1 is topologically stronger than d_2 if τ_{d_1} is finer.
2. d_1 is uniformly stronger than d_2 if $\exists C > 0$ such that $d_2 \leq C d_1$
3. d_1 is uniformly stronger than d_2 if $\exists C > 0$ such that $\frac{1}{C} d_1 \leq d_2 \leq C d_1$

Lemma 1

The following are equivalent

1. d_1 is topologically stronger than d_2
2. $\text{Id} : (X, \tau_{d_1}) \rightarrow (X, \tau_{d_2})$ is continuous
3. If $x_n \rightarrow \bar{x}$ in d_1 then $x_n \rightarrow \bar{x}$ in d_2
4. $\forall x \in X \forall \epsilon > 0 \exists \delta_{\epsilon, x} > 0$ such that

$$d(x, y) \leq \delta \implies d_2(x, y) < \epsilon$$

Definition 7

Let (X, d) be a metric space

1. $A \subset X$ is bounded if $\exists \bar{x} \in X$ such that $\sup_{y \in A} d(x, y) < \infty$ or $A = \emptyset$
2. x_n is Cauchy if

$$\lim_{n \rightarrow \infty} \sup_{i, j \geq n} d(x_i, x_j) = 0$$
3. X complete if x Cauchy $\implies x$ convergent.
4. (Y, e) is a metric, $f : X \rightarrow Y$ is uniformly continuous if $\forall \epsilon > 0 \exists \delta > 0$ such that $d(x, y) < \delta \implies e(f(x), f(y)) < \epsilon$.

Define $X = \{x : \mathbb{N} \rightarrow \mathbb{R} \text{ such that } \exists N \text{ such that } x_i = 0 \text{ eventually}\}$.

This space, with p -norm is not complete, so we construct the completion.

Proposition 2

Let (X, d) a metric space and (Y, e) a complete metric space, $A \subset X, \phi : A \rightarrow Y$ uniformly continuous.

Then \exists unique $\psi : \overline{A} \rightarrow Y$ such that ψ is uniformly continuous and $\phi = \psi|_A$.

Proof

If $x : \mathbb{N} \rightarrow A$ is Cauchy, then $\phi \circ x$ is also Cauchy.

To prove this, let $\epsilon > 0$ and $\delta_\epsilon > 0$ be such that $d(x, y) < \delta \implies e(\phi(x), \phi(y)) < \epsilon$.

Let $N = N_\delta^x$ be such that $i, j \geq N \implies d(x_i, x_j) < \delta$, then $e(\phi(x_i), \phi(x_j)) < \epsilon$

Now, let $a \in \overline{A}$, then $\exists x_k$ converging to a .

x is d -Cauchy and $\phi \circ x$ is e -cauchy.

\exists a limit $b^* = \lim \phi(x_k)$ So we define $\psi(a) = b^*$.

We now prove continuity/uniform continuity.

Let $a, b \in \overline{A}$, $x, y : \mathbb{N} \rightarrow A$ and $x_i \rightarrow b, y_j \rightarrow b$.

Then

$$e(\psi(a), \psi(b)) = \lim e(\phi(x_i), \phi(y_j))$$

Now, let $\epsilon > 0$, then $\exists \delta > 0$ such that $d(x, y) < \delta$.

Thus $e(\phi(x), \phi(y)) < \epsilon$

If $d(a, b) < \delta \exists N$ such that $d(x_i, y_j) < \delta \forall i, j > N$

$$e(\phi(x_i), \phi(y_j)) < \epsilon \implies e(\psi(a), \psi(b)) \leq \epsilon$$

□

Theorem 3

If (X, d) is a metric space, then there exists a complete metric space (Y, e) and an isometry $\phi : X \rightarrow Y$ such that $Y = \overline{\phi(X)}$.

Both are unique up to a bijective isometry.

Proof

Define $C_X := \{x : \mathbb{N} \rightarrow X, x \text{ Cauchy}\}$ and $x \tilde{y}$ if $\lim_{j \rightarrow \infty} d(x_i, y_j) = 0$.

Write $Y = C_X / \sim$.

For $x, y \in Y$, define $e(x, y) = \lim_{j \rightarrow \infty} d(x_i, x_j)$.

Is this well defined?

If $j, k \geq N$

$$|d(x_i, y_i) - d(x_k, y_k)| \leq d(x_i, x_k) + d(y_j, y_k)$$

And if $x \tilde{x}'$, then

$$\lim d(x_i, y_j) = \lim d(x'_j, y_j)$$

because

$$|d(x_i, y_j) - d(x'_j, y_j)| \leq d(x_j, x'_j) \rightarrow 0$$

To show that e is a metric, most properties are obvious.

We show that if $e(x, y) = 0$ then $\lim d(x_j, y_j) = 0 \implies x \tilde{y} \implies x = y$

Triangular equality holds because

$$e(x, y) = \lim d(x_j, y_j) \leq \limsup d(x_i, z_j) + d(z_j, y_j) = e(x, z) + e(z, y)$$

The isometry $\phi : X \rightarrow Y$ simply sends $x \mapsto [x]$.

We now show $[x] \in Y$, $\phi(x_k)$ is a sequence in Y , we want to show that

$\phi(x_k) \rightarrow [x]$.

$$\lim_{k \rightarrow \infty} e(\phi(x_k), [x]) = \lim_{k \rightarrow +\infty} \lim_{j \rightarrow \infty} d(x_k, x_j) = 0$$

Which shows $Y = \overline{\phi(X)}$ Let y^k Cauchy $\forall k \exists x_k \in X$ such that $e([y^k], \phi(x_k)) < 2^{-k}$.

We claim $[y^k] \rightarrow [x]$

$$d(x^k, x^h = e(\phi(x^k), \phi(x^h))) \leq 2^{-k} + 2^{-h+e([y^k], [y^h])}$$

Thus $x \in C_X$ $[x] \in Y$

$$e([y^k], [x]) = \lim d(y_j^k, x_j) \leq \lim d(U_j^k, x_k) + d(x_k, x_j) \leq 2^{-k}$$

Finally, to show uniqueness, if (Y, e) and (Y', e') are two completions.

Let $\psi = \phi \circ (\phi')^{-1} : \phi'(X) \rightarrow Y$.

ψ is an isometry so there is a unique extension $\psi : Y' \rightarrow Y$ and this is an isometry. \square

1.3 Norms, Banach Spaces

Throughout, $K = \mathbb{R}$ or \mathbb{C}

Definition 8 (Normed space)

$\|\cdot\| : X \rightarrow [0, \infty)$ is a norm if

- $\|x\| = 0 \iff x = 0$
- $\|\lambda x\| = |\lambda| \|x\|$
- $\|x + y\| \leq \|x\| + \|y\|$

Definition 9

c_0 is the space $c_0 = \{x : \mathbb{N} \rightarrow \mathbb{R} \text{ s.t. } \lim x_k = 0\}$ together with $\|x\|_{c_0} = \sup |x_k|$

For $p \in [1, \infty)$, $l_p = \{x : \mathbb{N} \rightarrow \mathbb{R} \text{ s.t. } \sum_{k \in \mathbb{N}} |x_k|^p < \infty\}$ with $\|x\|_{l_p} = (\sum |x_k|^p)^{\frac{1}{p}}$

Definition 10 (Banach Space)

A Banach space is a complete normed space.

Proposition 4

Any normed space has a completion which is Banach.

Proof

Let (Y, e) be the completion as above, define

$$[x] + [y] := [x + y] \text{ and } \lambda[x] := [\lambda x]$$

□

1.4 Basis of a normed space**Definition 11**

Let $A \subset X$.

A is linearly independent if $\forall N \in \mathbb{N}, \forall a_i \in A \forall \lambda_i \in K, \sum_i \lambda_i a_i = 0 \implies \lambda_i = 0$.

We define

$$\text{span}(A) = \left\{ \sum (i) \lambda_i a_i, \lambda_i \text{ as above} \right\}$$

A is a Hamel basis if A is linearly independent and $X = \text{span} A$

Definition 12 (Schauder Basis)

$e : \mathbb{N} \rightarrow X$ is a Schauder basis if $\forall x \in X$ there is a unique $\lambda : \mathbb{N} \rightarrow K$ such that $x = \sum_{i=0}^{\infty} \lambda_i e_i \iff \lim \left\| x - \sum^N \lambda_i e_i \right\| = 0$

Lecture 3: Projections onto Hilbert spaces

Wed 19 Oct

Definition 13 (Equivalence of Norms)

Let $\|\cdot\|_1$ and $\|\cdot\|_2$ be norms on a vector space X

1. $\|\cdot\|_1$ is stronger than $\|\cdot\|_2$ if the induced metrics are topologically stronger
2. $\|\cdot\|_1$ is equivalent to $\|\cdot\|_2$ if the induced metrics are equivalent

Lemma 5

1. If $\|\cdot\|_1$ is stronger than $\|\cdot\|_2 \implies \exists C > 0$ such that $\|x\|_2 \leq C \|x\|_1$
2. If $\|\cdot\|_1$ is equivalent to $\|\cdot\|_2 \implies \exists C > 0$ such that $\frac{1}{C} \|x\|_1 \leq \|x\|_2 \leq C \|x\|_1$

Proof

1. If not, $\forall k \in \mathbb{N}, \exists v_k \in X$ such that $\|v_k\|_2 > k \|v_k\|_1$.

Let $w_k = \frac{v_k}{\|v_k\|_2}$ then $1 = \|w_k\|_2 > k \|w_k\|_1$.

Thus $w_k \rightarrow 0 \in \|\cdot\|_1$, thus $w_k \rightarrow 0 \in \|\cdot\|_2$ which is a contradiction.

2. Follows from 1.

□

1.5 Scalar products and Hilbert spaces

Definition 14

Let H be a K -vector space.

A map $b : H \times H \rightarrow K$ is a scalar product if it satisfies

$$b(x, \lambda y + \mu z) = \lambda b(x, y) + \mu b(x, z)$$

$$b(\lambda x + \mu y, x) = \bar{\lambda} b(x, x) + \bar{\mu} b(y, x)$$

$$b(x, y) = \overline{b(y, x)} \text{ and } b(x, x) > 0.$$

(H, b) is a pre-Hilbert space

Example

1. K^d with the usual scalar product

2. $\ell^2(\mathbb{R})$

Proposition 7

1. $\|x\|_H = (x, x)^{\frac{1}{2}}$ is a norm on H
2. Cauchy-Schwarz : $|(x, y)| \leq \|x\| \|y\|$
3. $\|x + y\|^2 + \|x - y\|^2 = 2(\|x\|^2 + \|y\|^2)$

Proof

To show Cauchy-Schwarz, note that $(x + ty, x + ty) \geq 0 \forall t \in K$, thus

$$(x, x) + t((x, y) + (y, x)) + t^2(y, y) \geq 0 \quad \square$$

The middle term is $2t \operatorname{Re}(x, y)$, if the scalar product isn't real, we may rotate y to make it real

Proposition 8

Let $(X, \|\cdot\|)$ be a normed space.

If the parallelogram identity holds, then there is a scalar product b such that $\|x\| = b(x, x)^{\frac{1}{2}}$

Proof

Define $b(x, y) = \frac{1}{4} (\|x + y\|^2 - \|x - y\|^2)$.

We want to check $b(x, \lambda y + \mu z) = \lambda b(x, y) + \mu b(x, z)$.

First, check $b(x, y + y') + b(x, y - y') = 2b(x, y)$

$$\frac{1}{4} [\|x + y + y'\|^2 - \|x - y - y'\|^2 + \|x + y - y'\|^2 - \|x - y + y'\|^2] = \frac{1}{2} (\|x + y\|^2 - \|x - y\|^2)$$

From the parallelogram identity, we get that the left hand side is

$$\frac{1}{4} [2\|x + y\|^2 + 2\|y'\|^2 - 2\|x - y\|^2 - 2\|y'\|^2]$$

and thus the equality above holds.

If $y' = y \implies b(x, 2y) = 2b(x, y)$ and thus

$$y' = ny \quad b(x, (n+1)y) = 2b(x, y) - b(x, y - ny)$$

and we conclude by induction that $b(x, ny) = nb(x, y)$.

Thus $b(x, qy) = qb(x, y) \forall q \in \mathbb{Q}$ and by continuity, they agree on \mathbb{R} .

Pick $v, w \in X$ and $y = \frac{v+w}{2}, y' = \frac{v-w}{2}$ in the above equality, then

$$b(x, v) + b(x, w) = 2b(x, \frac{v+w}{2})$$

□

and we conclude from linearity.

For complex numbers, consider $s(x, y) = b(x, y) - ib(x, iy)$

Definition 15 (Hilbert Space)

(H, b) is a Hilbert space if it is a complete pre-Hilbert space.

Lemma 9

Every pre-Hilbert space has a completion, unique up to bijective isometry.

If $M \subset X$, then $p : X \rightarrow M$ is a projection if $p^2 = p$ and $p(X) = M$.

M is convex if $x, y \in M, t \in [0, 1]$, then $tx + (1-t)y \in M$

Theorem 10

Let H be a Hilbert space, $M \subset H$ non-empty, closed, convex, then \exists a unique map $p : H \rightarrow M$ such that

$$\|x - px\| = d(x, M)$$

Proof

If $x \in M, px = x$.

If $x \notin M$, let $d = d(x, M) > 0$.

If $y, z \in M$ are minimizers, then

$$\frac{1}{2} \|x - y\|^2 + \frac{1}{2} \|x - z\|^2 = \left\| x - \frac{y+z}{2} \right\|^2 + \left\| \frac{y-z}{2} \right\|^2$$

If $\|x - y\| = \|x - z\| = d$, then $\frac{y+z}{2} \in M \implies \|y - z\| \leq 0 \implies y = z \implies$
uniqueness.

To show existence, let $d = \inf_{y \in M} \|x - y\|$.

There is a sequence $y : \mathbb{N} \rightarrow M$ such that $\|x - y_k\| \rightarrow d$.

Thus

$$\frac{1}{2} \|x - y_h\|^2 + \frac{1}{2} \|x - y_k\|^2 = \left\| x - \frac{y_h + y_k}{2} \right\|^2 + \frac{1}{4} \|y_h - y_k\|^2$$

The LHS goes to d^2 and $\left\| x - \frac{y_h + y_k}{2} \right\| \geq d^2$. \square

Lemma 11

Let everything as above,, then $p : H \rightarrow M$ is an orthogonal projection

$$\iff \forall y \in M \forall x \in H, \operatorname{Re}(x - Px, y - px) \leq 0$$

Proof

Let $f(t) = \|x - (ty + (1 - t)px)\|^2$, thus $f(0) = \min f([0, 1])$, thus $f'(0) \geq 0$

$$f(t) = \|x - Px\|^2 - t[(x - px, y - px) + (y - px, x - px)] + t^2 \|y - px\|^2$$

Thus $f'(0) = -\operatorname{Re}(x - px, y - px) \geq 0$ \square

Corollary 12

If $M \subset H$ is a closed linear subspace, then

$$M^\perp = \{x \in H : (x, m) = 0 \forall m \in M\}$$

is a closed linear subspace, $M \cap M^\perp = \{0\}$, $H = M \oplus M^\perp$ and $p : H \rightarrow M$ satisfies $x - px \in M^\perp$ and p is linear

Definition 16 (Orthonormal systemes)

Let (X, b) be a pre-Hilbert space, the family $(e_i)_{i \in I}$ of vectors in X are orthonormal if $b(e_i, e_j) = \delta_{ij}$

Lemma 13

If the e_1, \dots, e_n are orthonormal, then $\forall x$

$$\sum_i |(e_i, x)|^2 \leq \|x\|^2$$

and if e_1, \dots, e_n are orthonormal, then $\forall x$

$$\sum_i |(e_i, x)|^2 \leq \|x\|^2$$

Proof

Let $y = \sum_i \lambda_i e_i \in \text{span} e_i$

$$g(\lambda) \left\| x - \sum_i \lambda_i e_i \right\|^2 = \|x\|^2 - \sum_i (\overline{\lambda_i}(e_i, x) + \lambda_i(x, e_i)) + \sum_i \lambda_i^2$$

Set $g(\lambda) \geq$

□