

RECEIVED: January 26, 2023 ACCEPTED: June 26, 2023 PUBLISHED: July 24, 2023

Model-independent search for the presence of new physics in events including $H\to\gamma\gamma$ with $\sqrt{s}=13$ TeV pp data recorded by the ATLAS detector at the LHC



The ATLAS collaboration

E-mail: atlas.publications@cern.ch

ABSTRACT: A model-independent search for new physics leading to final states containing a Higgs boson, with a mass of 125.09 GeV, decaying to a pair of photons is performed with $139\,\mathrm{fb^{-1}}$ of $\sqrt{s}=13\,\mathrm{TeV}$ pp collision data recorded by the ATLAS detector at the Large Hadron Collider at CERN. This search examines 22 final states categorized by the objects that are produced in association with the Higgs boson. These objects include isolated electrons or muons, hadronically decaying τ -leptons, additional photons, missing transverse momentum, and hadronic jets, as well as jets that are tagged as containing a b-hadron. No significant excesses above Standard Model expectations are observed and limits on the production cross section at 95% confidence level are set. Detector efficiencies are reported for all 22 signal regions, which can be used to convert detector-level cross-section limits reported in this paper to particle-level cross-section constraints.

Keywords: Hadron-Hadron Scattering, Higgs Physics

ARXIV EPRINT: 2301.10486

\mathbf{C}	ontents		
1	Introdu	ction	1
2	ATLAS	detector	3
3	Data ar	nd Monte Carlo samples	4
	3.1 Dat	ta sample	4
	3.2 Mo	nte Carlo samples	4
4	Object	and event selection	8
	4.1 Ob	ject definitions	8
	4.2 Eve	ent preselection	9
5	Signal r	regions	9
6	Signal a	and background modelling	12
	6.1 Mo	delling of BSM and SM Higgs boson production	12
	6.2 Cor	ntinuum background modelling	13
	6.3 Mo	delling of the background in the multilepton signal region	13
7	Systems	atic uncertainties	14
	7.1 Sys	tematic uncertainties in modelling $m_{\gamma\gamma}$ for SM and BSM Higgs boson	
	pro	duction	14
	7.2 Uno	certainty in the continuum background modelling	15
	7.3 Uno	certainties in resonant background yields	15
8	Results		15
	8.1 Up	per limits on BSM production cross sections	19
	8.2 Inte	erpretation with particle-level simulation	21
9	Conclus	sions	24
\mathbf{T}	he ATLA	S collaboration	32

1 Introduction

The discovery of the Higgs boson by the ATLAS and CMS experiments at the CERN Large Hadron Collider (LHC) [1, 2] opened up new avenues to search for physics beyond the Standard Model (BSM physics). The properties of the Higgs boson, including its inclusive production cross sections and decay rates, as well as its differential cross sections, may be altered by the presence of new interactions and new particles beyond the SM. In addition,

the Higgs boson may also be produced in BSM processes as a decay product of unknown new particles, which could give rise to deviations of Higgs boson production rates from SM predictions in exclusive final states. For example, the Higgs boson can be produced in the decays of supersymmetric particles predicted by weak-scale supersymmetry (SUSY) models [3–8], in exotic decays of top quarks in models such as those having flavour-changing neutral currents (FCNCs) [9–13], or in the decay of the partner of the top quark or bottom quark as predicted by models of vector-like quarks [14, 15]. So far, observations of Higgs boson production and decay properties at the LHC are consistent with SM predictions [16, 17], and direct searches targeting the aforementioned BSM models also reported no significant deviations from the SM expectation. Most of these analyses are designed with certain model assumptions, either targeting a specific SM Higgs boson production mode or optimized for a given BSM process involving the production of the Higgs boson.

This paper reports the results of a model-independent search for excesses in the production rate of the Higgs boson with a mass of 125.09 GeV in many distinct signal regions with the ATLAS detector at the LHC. The regions are defined by the presence and kinematic properties of objects produced in association with the Higgs boson. While a large fraction of such signals may populate signal regions probed by SM Higgs boson measurements, such as measurements of "simplified template cross sections" [18, 19] and fiducial and differential cross-section measurements [20], unknown BSM production of the Higgs boson could result in events that do not enter signal regions optimized for existing SM measurements. The signal regions in this search are chosen to cover a wide range of signatures defined by particles accompanying the production of the Higgs boson, and they are defined with simple and mostly inclusive selections in order to minimize model dependence.

In this search the Higgs boson is reconstructed via its diphoton decay channel, which has a predicted branching ratio of $(0.227\pm0.007)\%$ [18] in the SM. While the rate of $H\to\gamma\gamma$ decay is much lower than that of several other Higgs boson decay channels, the sensitivity of an analysis targeting this channel can be competitive with that of analyses targeting higher-rate channels. This is primarily because the diphoton background to the Higgs boson signal is relatively small and because of better experimental resolution for the reconstructed resonance mass, especially in comparison with hadronic decay modes of the Higgs boson. Another advantage of performing this model-independent search in the $H\to\gamma\gamma$ channel is that the same background modelling strategy can be applied to most signal regions; the signal and background can be determined by fitting analytic functions to the diphoton invariant mass $(m_{\gamma\gamma})$ distribution in data. This fit-based background estimation method breaks down only when the signal region does not have a sufficient number of events to constrain the analytic background function.

If data in the signal regions are found to be consistent with the SM expectations, including contributions from resonant Higgs boson production, limits are set on the BSM production cross section of the Higgs boson in detector-level signal regions. To enable the use of these detector-level cross-section limits in constraining BSM physics models, the detector efficiencies for different signal regions are evaluated. These efficiencies depend on the kinematics of the final-state particles and therefore vary between different BSM models. As a result, this paper reports a range of efficiency values for each signal region, evaluated from Monte Carlo simulation of several benchmark BSM models. Correlations in the observed limits between signal regions with a large number of events are also reported.

This paper is organized as follows. Section 2 gives a brief description of the ATLAS detector. Section 3 presents the simulated SM samples used to describe the backgrounds, and also the new-physics models used to determine the expected sensitivity of the search in each signal region. Section 4 describes the reconstructed data objects and subsequent selections used in the analysis. Section 5 discusses in detail the signal regions defined for the search. Section 6 describes the modelling of signal and background in each search region. Section 7 addresses systematic uncertainties related to the search. Section 8 presents the results of this search. Finally, section 9 summarizes the conclusions of the paper.

2 ATLAS detector

The ATLAS detector [21] at the LHC covers nearly the entire solid angle around the collision point.¹ It consists of an inner tracking detector surrounded by a thin superconducting solenoid, electromagnetic and hadronic calorimeters, and a muon spectrometer incorporating three large superconducting toroidal magnet systems. The inner-detector system is immersed in a 2T axial magnetic field and provides charged-particle tracking in the range $|\eta| < 2.5$.

The fine-granularity silicon pixel detector covers the vertex region and typically provides four measurements per track, the first hit normally being in the insertable B-layer installed before Run 2 [22, 23]. The pixel detector is followed by the silicon microstrip tracker, which usually provides eight measurements per track. These silicon detectors are complemented by the transition radiation tracker (TRT), which enables radially extended track reconstruction up to $|\eta| = 2.0$. The TRT also provides electron identification information based on the fraction of hits (typically 30 hits in total) above a higher energy-deposit threshold corresponding to transition radiation.

The calorimeter system covers the pseudorapidity range $|\eta| < 4.9$. Within the region $|\eta| < 3.2$, electromagnetic (EM) calorimetry is provided by barrel and endcap lead/liquid-argon (LAr) calorimeters, with an additional thin LAr presampler covering $|\eta| < 1.8$ to correct for energy loss in material upstream of the calorimeters. Hadronic calorimetry is provided by a steel/scintillator-tile calorimeter, segmented into three barrel structures within $|\eta| < 1.7$, and two copper/LAr hadronic endcap calorimeters. The solid angle coverage is completed with forward copper/LAr and tungsten/LAr calorimeter modules optimized for electromagnetic and hadronic measurements, respectively.

The muon spectrometer comprises separate trigger and high-precision tracking chambers measuring the deflection of muons in a magnetic field generated by the superconducting air-core toroids. The field integral of the toroids ranges between 2.0 and 6.0 T m across most of the detector. A set of precision chambers covers the region $|\eta| < 2.7$ with three layers of monitored drift tubes, complemented by cathode-strip chambers in the forward region,

¹ATLAS uses a right-handed coordinate system with its origin at the nominal interaction point (IP) in the centre of the detector and the z-axis along the beam pipe. The x-axis points from the IP to the centre of the LHC ring, and the y-axis points upwards. Cylindrical coordinates (r, ϕ) are used in the transverse plane, ϕ being the azimuthal angle around the z-axis. The pseudorapidity is defined in terms of the polar angle θ as $\eta = -\ln \tan(\theta/2)$. Angular distance is measured in units of $\Delta R \equiv \sqrt{(\Delta \eta)^2 + (\Delta \phi)^2}$.

where the background is highest. The muon trigger system covers the range $|\eta| < 2.4$ with resistive-plate chambers in the barrel, and thin-gap chambers in the endcap regions.

Interesting events are selected to be recorded by the first-level trigger system implemented in custom hardware, followed by selections made by algorithms implemented in software in the high-level trigger [24]. The first-level trigger accepts events from the 40 MHz bunch crossings at a rate below 100 kHz, which the high-level trigger reduces in order to record events to disk at about 1 kHz. An extensive software suite [25] is used in data simulation, in the reconstruction and analysis of real and simulated data, in detector operations, and in the trigger and data acquisition systems of the experiment.

3 Data and Monte Carlo samples

3.1 Data sample

This study uses the full Run-2 data set of $\sqrt{s}=13$ TeV proton-proton collisions recorded by the ATLAS detector during 2015–2018. After data quality requirements [26] are applied to ensure the good working condition of all detector components, the data set amounts to an integrated luminosity of $139.0 \pm 2.4 \, \text{fb}^{-1}$ [27, 28]. The mean number of interactions per bunch crossing was $\langle \mu \rangle = 33.7$ on average during Run 2.

The data sample was selected using a combination of diphoton and single-photon triggers. The transverse energy thresholds of the diphoton trigger were 35 GeV and 25 GeV for the leading and subleading photon candidates, respectively [29]. The diphoton trigger applies photon identification selections based on calorimeter shower shape variables. In 2015-2016, a loose photon identification requirement was used, and in 2017-2018, this requirement was tightened to cope with higher instantaneous luminosity. The single-photon trigger had a transverse energy threshold of 120 GeV for the leading photon in data collected between 2015 and 2017, and the threshold was increased to 140 GeV for data collected in 2018. The photon candidate used in the trigger decision is required to satisfy the loose photon identification criteria. The use of the single-photon trigger is found to improve the trigger efficiency for high- p_T Higgs boson candidates. Typically, the trigger efficiency is greater than 98% for events that pass the diphoton event selection described below.

3.2 Monte Carlo samples

Monte Carlo (MC) simulation samples are used to determine the SM Higgs boson event yield, test the continuum background modelling, and estimate detector efficiencies for various signal regions. These simulated samples include three categories of processes: SM Higgs boson production, SM diphoton continuum production, and BSM production of the Higgs boson with a mass of 125.09 GeV.

MC samples generated for SM Higgs boson production are summarized in table 1. The gluon-gluon fusion (ggF), vector-boson fusion (VBF), and VH production modes were generated using POWHEG NNLOPS or POWHEG BOX [30–38], with the PDF4LHC15 PDF set [39], and interfaced to PYTHIA 8 [40, 41] for simulation of parton showering, hadronization and the underlying event, using a set of data-tuned parameters called the AZNLO tune [42].

Process	Generator	PDF	PDF, Tune	Cross section [pb]
	(ME+PS)	(ME)	(PS, UE)	$(\sqrt{s} = 13 \text{ TeV})$
ggF	POWHEG NNLOPS + PYTHIA 8	PDF4LHC15	CTEQ6, AZNLO	48.5
VBF	POWHEG BOX + PYTHIA 8	PDF4LHC15	CTEQ6, AZNLO	3.78
WH	POWHEG BOX + PYTHIA 8	PDF4LHC15	CTEQ6, AZNLO	1.37
$q\bar{q}\to ZH$	POWHEG BOX + PYTHIA 8	PDF4LHC15	CTEQ6, AZNLO	0.76
$gg\to ZH$	POWHEG BOX + PYTHIA 8	PDF4LHC15	CTEQ6, AZNLO	0.12
$t \bar{t} H$	POWHEG BOX + PYTHIA 8	PDF4LHC15	NNPDF2.3, A14	0.51
$b ar{b} H$	POWHEG BOX + PYTHIA 8	PDF4LHC15	NNPDF2.3, A14	0.49
tHbj	aMC@NLO $+$ Pythia 8	NNPDF3.0	NNPDF2.3, A14	0.074
tWH	${ m aMC@NLO} + { m Pythia} 8$	NNPDF3.0	NNPDF2.3, A14	0.015

Table 1. Configuration of MC simulation of SM Higgs boson production and the predicted cross sections at $\sqrt{s} = 13$ TeV. The generator set-up includes the software programs used for the matrix element (ME) calculation and parton showering (PS). The parton distribution function (PDF) is given separately for the ME calculation and the parton showering, along with the set of tuned parameters (Tune) used in modelling the underlying event (UE).

The ggF simulation achieves NNLO accuracy for arbitrary inclusive $gg \to H$ observables by reweighting the Higgs boson rapidity spectrum in Hj-MiNLO [36, 43, 44] to that of HNNLO [45]. The transverse momentum spectrum of the Higgs boson obtained with this sample is found to be compatible with the fixed-order HNNLO calculation and the HRES 2.3 calculation [46, 47] performing resummation at next-to-next-to-leading-logarithm accuracy matched to a NNLO fixed-order calculation (NNLL+NNLO). The VBF production mode was simulated at next-to-leading-order (NLO) accuracy in QCD. The simulation of WHand $q\bar{q} \to ZH$ production is accurate to NLO in QCD with up to one extra jet in the event, while the simulation of the $gg \to ZH$ process was performed at leading order in QCD. The $t\bar{t}H$ and bbH processes were modelled using the POWHEG BOX v2 [31–33, 38, 48] generator, which provides matrix elements at NLO in the strong coupling constant α_s in the five-flavour scheme with the NNPDF2.3LO [49] PDF set. For the $t\bar{t}H$ sample, the functional form of the renormalization and factorization scale was set to $\sqrt[3]{m_{\rm T}(t) \cdot m_{\rm T}(\bar{t}) \cdot m_{\rm T}(H)^2}$ [18]. The generator was interfaced to Pythia 8.230 [41], which used the A14 tune [50] and the NNPDF2.3LO [49] PDF set. The decays of bottom and charm hadrons were simulated using the EVTGEN 1.6.0 program [51]. All the SM Higgs boson samples are normalized to cross-section values recommended by ref. [18], which are shown in table 1.

To study the modelling of continuum background, diphoton production event samples were simulated. Besides inclusive diphoton production, specific diphoton processes were also generated for signal regions where electroweak or top-associated production is enhanced. Inclusive diphoton production and diphoton production in association with a vector boson $(V\gamma\gamma)$ were simulated with the Sherpa 2.2.4 [52] generator. Matrix elements (MEs) for up to three additional partons were calculated at LO accuracy. For the inclusive diphoton

 $^{^2}m_{\rm T}$ denotes the transverse mass of a particle, defined as $m_{\rm T}=\sqrt{m^2+p_{\rm T}^2}$ where m and $p_{\rm T}$ are its mass and transverse momentum, respectively.

sample, NLO accuracy is reached for MEs including up to one additional parton. The Comix [53] and OpenLoops [54–56] libraries were used in these calculations, which were matched with the Sherpa parton shower [57] using the MEPS@NLO prescription [58–61] with a dynamic merging cut [62] of 10 GeV. A smooth-cone isolation requirement [63] was applied in the event generation. Samples were generated using the NNPDF3.0NNLO PDF set [49], along with the dedicated set of tuned parton-shower parameters developed by the Sherpa authors. Diphoton production in association with a top-quark pair $(t\bar{t}\gamma\gamma)$ was modelled by the Madgraph5_aMC@NLO 2.3.3 generator at LO with the NNPDF2.3LO [64] PDF set, and the events were interfaced with Pythia 8.212 for parton-shower and underlying-event simulation.

Several BSM processes were simulated to demonstrate the coverage of the signal regions and estimate the model dependence of the efficiency of the fiducial selections. These samples are summarized in table 2 and are described as follows.

- 1. BSM Higgs boson production characterized by the presence of additional charged or neutral leptons:
 - $\tilde{\chi}_1^{\pm} \tilde{\chi}_2^0 \to W/Z/H$: the production of a neutralino-chargino pair that decays to final states involving W, Z and Higgs bosons.
 - $\tilde{\chi}_1^{\pm} \tilde{\chi}_1^{\mp} \to H \ell^{\pm} H \ell^{\mp}$: the production of a chargino pair that decays to final states involving Higgs bosons and leptons through an R-parity-violating interaction.
- 2. BSM Higgs boson production characterized by the presence of additional heavy-flavour jets are represented by
 - $t\tilde{t}$, $b\tilde{b}$: stop-pair production and sbottom-pair production processes. In the subsequent decays of the stop or sbottom particles, the Higgs boson could be produced in the decays of next-to-lightest supersymmetric particles.
- 3. BSM Higgs boson production characterized by the presence of an additional top quark is represented by
 - $t\bar{t}$, $(t \to u/c + H) + \text{c.c.}$: pair-produced top quarks that decay into final states involving Higgs bosons, through a FCNC interaction.
- 4. BSM Higgs boson production characterized by the presence of an additional photon is represented by
 - $pp \to \chi \to H\gamma$: a massive neutral boson (χ) decaying into a photon and a Higgs boson, a process predicted by extensions of the Higgs sector or gauge sectors [65–67].

These BSM processes are studied in various dedicated ATLAS searches that may use Higgs boson decay channels other than $H \to \gamma \gamma$. The specific process, model parameters, expected cross section, and a reference to the dedicated search, are summarized in table 2. These BSM MC samples were generated with the same set-up as used in the dedicated searches, except that the Higgs boson is configured to decay into a pair of photons.

Process with $H \to \gamma \gamma$	Cianal maga	Cross section [pb]	Ref.
Process with $n \to \gamma \gamma$	Signal mass	Cross section [pb]	nei.
$ ilde{\chi}_1^{\pm} ilde{\chi}_2^0 \! o W/Z/H$	$m(\tilde{\chi}_1^{\pm}/\tilde{\chi}_2^0), m(\tilde{\chi}_1^0) = 150, 0.5 \text{ GeV}$	5.18	[69]
$\tilde{\chi}_1^{\pm} \tilde{\chi}_2^0 \rightarrow W/Z/H$	$m(\tilde{\chi}_1^{\pm}/\tilde{\chi}_2^0), m(\tilde{\chi}_1^0) = 300, 0.5 \text{ GeV}$	0.39	[69]
$\tilde{\chi}_1^{\pm} \tilde{\chi}_1^{\mp} \to H \ell^{\pm} H \ell^{\mp}$	$m(\tilde{\chi}_1^{\pm}) = 150 \text{ GeV}$	2.61	[70]
$\tilde{\chi}_1^{\pm} \tilde{\chi}_1^{\mp} \to H \ell^{\pm} H \ell^{\mp}$	$m(\tilde{\chi}_1^{\pm}) = 300 \text{ GeV}$	0.19	[70]
$\tilde{\chi}_1^{\pm} \tilde{\chi}_1^0, \tilde{\chi}_1^{\pm} \to H\ell^{\pm}, \tilde{\chi}_1^0 \to W\ell/Z\nu/H\nu$	$m(\tilde{\chi}_1^{\pm}/\tilde{\chi}_1^0) = 200 \text{ GeV}$	1.81	[70]
$\tilde{\chi}_1^{\pm} \tilde{\chi}_1^0, \tilde{\chi}_1^{\pm} \to H \ell^{\pm}, \tilde{\chi}_1^0 \to W \ell/Z \nu/H \nu$	$m(\tilde{\chi}_1^{\pm}/\tilde{\chi}_1^0) = 400 \text{ GeV}$	0.12	[70]
$\tilde{t}_2 \tilde{t}_2, \tilde{t}_2 \rightarrow \tilde{t}_1 H, \tilde{t}_1 \rightarrow \tilde{\chi}_1^0 b q \bar{q}/b \ell \nu$	$m(\tilde{t}_2), m(\tilde{t}_1), m(\tilde{\chi}_2^0) = 500, 340, 300 \text{ GeV}$	0.61	[71]
$\tilde{b}\tilde{b}, \tilde{b} \rightarrow \tilde{\chi}_2^0 b, \tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 H$	$m(\tilde{b}), m(\tilde{\chi}_2^0), m(\tilde{\chi}_1^0) = 500, 180, 50 \text{ GeV}$	0.61	[72]
$ ilde{b} ilde{b}, ilde{b} ightarrow ilde{\chi}_2^0 b, ilde{\chi}_2^0 ightarrow ilde{\chi}_1^0 H$	$m(\tilde{b}), m(\tilde{\chi}_2^0), m(\tilde{\chi}_1^0) = 1000, 205, 60 \text{ GeV}$	0.0068	[7 2]
$ ilde{b} ilde{b}, ilde{b} ightarrow ilde{\chi}_2^0 b, ilde{\chi}_2^0 ightarrow ilde{\chi}_1^0 H$	$m(\tilde{b}), m(\tilde{\chi}_2^0), m(\tilde{\chi}_1^0) = 1200, 205, 60 \text{ GeV}$	0.0017	[7 2]
$pp {\to} \chi {\to} H\gamma$	$m(\chi) = 200 \text{ GeV}$	_	[73]
$pp \to \chi \to H\gamma$	$m(\chi) = 500 \text{ GeV}$	_	[73]
$t\bar{t}, (t \rightarrow u/c + H) + \text{c.c.}$	_	0.00123	[74]

Table 2. Summary of processes, model parameters, and cross sections for the BSM samples used in this paper. A reference to their dedicated ATLAS search is also included. The cross sections quoted include the SM branching ratio for $H \to \gamma \gamma$. The $pp \to \chi \to H \gamma$ process is expected in multiple BSM models, so no particular model is used for the cross-section calculation.

The SUSY events were generated with up to two additional partons in the matrix element using Madgraph5_aMC@NLO 2.6.2 at LO in QCD with the NNPDF3.0LO PDF set and CKKW-L merging scheme. Parton showering and hadronization were handled by Pythia 8.230 with the A14 tune, using the NNPDF2.3LO PDF set.

MC samples were generated for the BSM process where a massive neutral resonance is produced and decays into a Higgs boson and a photon $(pp \to \chi \to H\gamma)$. This process could result in a final state with at least three photons. The event generation used the same set-up as for the SUSY samples, except that PYTHIA 8.235 was used to model the parton shower and underlying event.

A $t\bar{t}$ sample with FCNC-induced top-quark decays is also considered. In this process, one top quark decays into a Higgs boson and an up-type quark through a FCNC interaction, while the other top quark decays into a b-quark and a W boson. These FCNC samples were simulated using Madgraph5_aMC@NLO 2.4.3 [68] interfaced to Pythia 8.212 [40] with the A14 tune for the modelling of parton showers, hadronization and the underlying event. The FCNC samples were generated separately for the up-quark and charm-quark final states.

The effect of multiple interactions in the same and neighbouring bunch crossings (pile-up) was modelled by overlaying each simulated hard-scattering event with inelastic proton-proton (pp) events generated by Pythia 8.186 using the NNPDF2.3Lo set of PDFs and the A3 tune [75]. The generated Higgs boson events from table 1 were passed through a Geant4 [76] simulation of the ATLAS detector [77]. The remaining QCD diphoton events and BSM samples are passed through a fast simulation of the ATLAS detector to reduce the amount of CPU time needed to process the large numbers of generated events. All simulated events are reconstructed with the same software as used for the data [25].

4 Object and event selection

4.1 Object definitions

Events in this analysis are selected using the following procedure. First, reconstructed photon candidates are required to satisfy a set of preselection criteria. The two highest- $p_{\rm T}$ preselected photons along with the reconstructed vertex information in the event, are used as inputs to a neural-network algorithm trained on simulated events to determine the correct primary vertex, which is named the "diphoton vertex" [78]. The selected primary vertex is used to compute the properties of objects in the event. Finally, to be selected the photons are required to satisfy isolation and additional identification criteria. In addition to photons, jets (including b-jets), muons, electrons, and the amount of missing transverse momentum, $E_{\rm T}^{\rm miss}$, are used in the analysis to define model-independent signal regions. The specific requirements for objects used in this analysis are as follows.

Photons must satisfy $p_{\rm T} > 22$ GeV and $|\eta| < 2.37$, excluding the transition region $1.37 < |\eta| < 1.52$ between the barrel and endcap EM calorimeters. Photon candidates are separated from jet backgrounds using a tight identification criterion based on calorimeter shower shape variables [79]. The identification efficiency for reconstructed photons ranges from 84% at $p_{\rm T}=25$ GeV to 94% at $p_{\rm T}>100$ GeV. The final selection of photons includes both calorimeter- and track-based isolation requirements to further suppress jets misidentified as photons. The calorimeter isolation variable is defined as the energy in the EM calorimeter within a cone of size $\Delta R = 0.2$ around the photon candidate, excluding the energy in a fixed-size window that contains the photon shower; a correction is made for photon energy leakage from this window [79]. Contributions from pile-up and the underlying event are subtracted [79–83]. The calorimeter-based isolation must be less than 6.5% of each photon candidate's transverse energy. The track-based isolation variable is defined as the scalar sum of the transverse momenta of tracks within a $\Delta R = 0.2$ cone around the photon candidate. The tracks used in the isolation variable are restricted to those with $p_T > 1$ GeV that are associated with the selected diphoton vertex and not with a photon conversion vertex [79]. The track isolation must be less than 5% of each photon candidate's transverse energy.

Electron candidates must have $p_{\rm T}>10$ GeV and $|\eta|<2.47$, excluding the EM calorimeter transition region of $1.37<|\eta|<1.52$, and must satisfy a medium selection based on a likelihood discriminant using calorimeter shower shapes and track parameters [79]. Isolation criteria based on calorimeter- and track-based information are applied to electrons. The reconstructed track matched to the electron candidate must be consistent with the diphoton vertex by requiring its longitudinal impact parameter z_0 relative to the vertex to satisfy $|z_0 \sin \theta| < 0.5$ mm. In addition, the electron's transverse impact parameter d_0 with respect to the beam-spot must satisfy $|d_0|/\sigma_{d_0}<5$, where σ_{d_0} is the uncertainty in d_0 .

Muon candidates are required to have $p_{\rm T}>10$ GeV and $|\eta|<2.7$, and must satisfy the medium identification requirements [84]. Muons are required to satisfy calorimeter- and track-based isolation requirements that are 95%–97% efficient for muons with $p_{\rm T}\in[10,60]$ GeV and 99% efficient for $p_{\rm T}>60$ GeV. Muon tracks must satisfy $|z_0\sin\theta|<0.5$ mm and $|d_0|/\sigma_{d_0}<3$.

Jets are reconstructed using a particle-flow algorithm [85]. It improves the energy resolution by applying the anti- k_t algorithm [86, 87] with radius parameter R=0.4 to noise-suppressed positive-energy topological clusters [88] in the calorimeter after removing energy deposits associated with primary-vertex-matched tracks, and including the track momenta in the clustering instead. Jets must have $p_T > 25$ GeV and |y| < 4.4. To suppress jets from pile-up, a jet-vertex-tagger (JVT) multivariate discriminant [89] is applied to jets with $p_T < 60$ GeV and $|\eta| < 2.4$; in the range $|\eta| > 2.5$, a 'forward' version of the JVT [90] is applied to jets with $p_T < 120$ GeV.³ Jets with $|\eta| < 2.5$ containing b-hadrons are identified using the DL1r b-tagging algorithm and its 60%, 70%, 77% and 85% efficiency working points, with the outputs combined into a pseudo-continuous b-tagging score [91].

Hadronically decaying τ -leptons are used in this search. They are required to have $p_{\rm T} > 20$ GeV and $|\eta| < 2.7$. Furthermore, they must have either one or three charged tracks ("prongs") with a charge sum of \pm 1 in units of the elementary charge, and must satisfy the *medium* working point [92].

An overlap removal procedure is performed in order to avoid double-counting of objects. First, electrons overlapping with any of the two selected photons ($\Delta R < 0.4$) are removed. Jets overlapping with the selected photons ($\Delta R < 0.4$) and electrons ($\Delta R < 0.2$) are removed. Electrons overlapping with the remaining jets ($\Delta R < 0.4$) are removed to match the requirements imposed when measuring isolated-electron efficiencies. Finally, muons overlapping with photons or jets ($\Delta R < 0.4$) are removed.

The missing transverse momentum is defined as the negative vector sum of the transverse momenta of the selected photons, electrons, muons, and jets, as well as the transverse momenta of remaining low- $p_{\rm T}$ particles, estimated using tracks associated with the diphoton vertex but not with any selected object [93].

4.2 Event preselection

The two highest- $p_{\rm T}$ preselected photon candidates are required to satisfy the *tight* identification criteria and the isolation selection described above. Finally, the leading and subleading photon candidates are required to satisfy $p_{\rm T}^{\gamma 1} > 35$ GeV, $p_{\rm T}^{\gamma 1}/m_{\gamma\gamma} > 0.35$, and $p_{\rm T}^{\gamma 2} > 25$ GeV, $p_{\rm T}^{\gamma 2}/m_{\gamma\gamma} > 0.25$, respectively. Events that fail the tight identification or the isolation selection but pass the loose identification are used as a control sample for background estimation and modelling purposes.

The trigger, object, and event selection described above are used to define the events that are selected for further analysis. In total, about 1.2 million events are selected in this data set with a diphoton invariant mass between 105 and 160 GeV.

5 Signal regions

Preselected events are assigned to signal regions defined by the presence of additional reconstructed objects. These signal regions are designed with simple, inclusive selections, and the specific selections were not optimized for any particular BSM processes. Since

 $^{^{3}\}eta$ is used to select jets when applying the JVT and the *b*-tagging requirements, as the way these selections are defined is based on the jet position in the detector.

the selection criteria are not designed to be orthogonal, selected events can be assigned to multiple signal regions. The overlap and correlations of these signal regions are discussed in section 8. Based on the requirement of there being additional objects in the event, the signal regions can be broadly classified to six groups: events with heavy-flavour jets; events with high jet multiplicity or a large scalar sum of jet transverse momenta (H_T) ; events with large E_T^{miss} ; events with leptons; events with additional photons; and events with top quarks. As is discussed in section 8, each signal region, defined through selection criteria for detector-level objects, has a counterpart defined through equivalent criteria at the particle level, which enables the interpretation of the search results with particle-level simulation samples. These signal regions and their definitions are summarized in table 3 and described as follows.

Two "heavy-flavour" signal regions are defined. One requires the presence of at least three b-jets tagged at the 85% efficiency working point (WP), and the other requires the presence of at least four b-jets tagged at the same working point.

Six high-jet-activity signal regions are defined. Three of them are defined with inclusive jet-multiplicity requirements, using central jets with $|\eta| < 2.5$. The thresholds for the number of jets are four, six, and eight for these regions. Another three regions are defined by the $H_{\rm T}$. The inclusive $H_{\rm T}$ thresholds for these three regions are 500 GeV, 1000 GeV, and 1500 GeV.

Three high- $E_{\rm T}^{\rm miss}$ signal regions are defined using $E_{\rm T}^{\rm miss}$ thresholds of 100 GeV, 200 GeV, and 300 GeV.

Six leptonic signal regions are defined. The inclusive leptonic signal region ($\geq 1\ell$) requires the presence of at least one electron or muon in the event. Three dilepton signal regions are defined with electrons and muons: the inclusive dilepton signal region (2ℓ) requires the presence of exactly two leptons; the dilepton Z-veto (2ℓ –Z) region is a subset of the 2ℓ region, and it explicitly rejects events where the two leptons have the same flavour and their invariant mass is within 10 GeV of the PDG value of the Z boson mass; the same-sign dilepton (SS- 2ℓ) region is also a subset of the 2ℓ region and requires the electric charges of the two leptons to be the same; the multilepton signal region ($\geq 3\ell$) requires the presence of at least three leptons. The τ -lepton signal region requires the presence of at least two τ -lepton candidates.

Two triphoton signal regions are defined. Both require the presence of at least one additional photon in the preselected diphoton events. In the first photon signal region, the diphoton system that is considered as the Higgs boson candidate is built from the two leading photons in $p_{\rm T}$. The second photon region assumes the subleading and third-leading photons come from the Higgs boson decay. While the two photon signal regions mostly have the same definitions of the event selection criteria, the $m_{\gamma\gamma}$ observable is constructed differently. The $1\gamma-m_{\gamma\gamma}^{12}$ region targets exotic processes where the Higgs boson is accompanied by a photon with moderate $p_{\rm T}$, while the $1\gamma-m_{\gamma\gamma}^{23}$ region targets exotic processes in which a new state that is much more massive than the Higgs boson decays into a photon and a Higgs boson.

Three "top" signal regions are introduced. One inclusive top signal region (ℓb) requires the presence of at least one electron or one muon and at least one b-jet tagged at the 70% efficiency working point. One exclusive semileptonic top-quark signal region ($t_{\rm lep}$) requires the presence of exactly one lepton, and one b-jet tagged at the 70% efficiency working

Target	Region	Detector level	Particle level
	>3h	$n_{b\text{-iet}} \ge 3, 85\% \text{ WP}$	$n_{b ext{-jet}} \ge 3$
Heavy flavour	$\geq 4b$	$n_{b\text{-jet}} \ge 4,85\% \text{ WP}$	$n_{b ext{-jet}} \geq 4$
	≥4j	$n_{ m jet} \ge 4, \eta_{ m jet} < 2.5$	$n_{ m jet} \ge 4, \eta_{ m jet} < 2.5$
	≥6j	$n_{ m jet} \geq 6, \eta_{ m jet} < 2.5$	$n_{ m jet} \ge 6, \eta_{ m jet} < 2.5$
High jet	≥8j	$n_{ m jet} \geq 8, \eta_{ m jet} < 2.5$	$n_{ m jet}\geq 8, \eta_{ m jet} < 2.5$
activity	$H_{\rm T} > 500~{\rm GeV}$	$H_{\rm T} > 500~{ m GeV}$	$H_{\mathrm{T}} > 500 \mathrm{GeV}$
	$H_{\mathrm{T}} > 1000 \ \mathrm{GeV}$	$H_{\rm T} > 1000~{ m GeV}$	$H_{\mathrm{T}} > 1000 \mathrm{GeV}$
	$H_{\rm T} > 1500~{\rm GeV}$	$H_{\rm T} > 1500~{ m GeV}$	$H_{\rm T} > 1500~{ m GeV}$
	$E_{\mathrm{T}}^{\mathrm{miss}} > 100 \text{ GeV}$	$E_{\mathrm{T}}^{\mathrm{miss}} > 100 \text{ GeV}$	$E_{\mathrm{T}}^{\mathrm{miss,tru}} > 100 \text{ GeV}$
$E_{\mathrm{T}}^{\mathrm{miss}}$	$E_{\mathrm{T}}^{\mathrm{miss}} > 200 \text{ GeV}$	$E_{\mathrm{T}}^{\mathrm{miss}} > 200 \mathrm{GeV}$	$E_{\mathrm{T}}^{\mathrm{miss,tru}} > 200 \text{ GeV}$
	$E_{\mathrm{T}}^{\mathrm{miss}} > 300 \text{ GeV}$	$E_{\mathrm{T}}^{\mathrm{miss}} > 300 \text{ GeV}$	$E_{\mathrm{T}}^{\mathrm{miss,tru}} > 300 \text{ GeV}$
	ℓb	$n_{\ell=e,\mu} \ge 1, \ n_{b\text{-jet}} \ge 1, \ 70\% \text{ WP}$	$n_{\ell=e,\mu} \ge 1, \ n_{b\text{-jet}} \ge 1$
Тор	$t_{ m lep}$	$n_{\ell=e,\mu} = 1, n_{\rm jet} = n_{b ext{-jet}} = 1$ 70% WP	$n_{\ell=e,\mu} = 1, \ n_{\text{jet}} = n_{b\text{-jet}} = 1$
	$t_{ m had}$	$n_{\ell=e,\mu}=0,\ n_{\rm jet}=3,\ n_{b\text{-jet}}=1$ 70% WP, BDT $_{ m top}>0.9$, $n_{\ell=e,\mu} = 0$, $n_{\rm jet} = 3$, $n_{b\text{-jet}} = 1$
	$\geq 1\ell$	$n_{\ell=e,\mu} \ge 1$	$n_{\ell=e,\mu} \ge 1$
	2ℓ	$ee, \mu\mu$, or $e\mu$	$ee, \mu\mu$, or $e\mu$
Lepton	2ℓ – Z	$ee, \mu\mu, e\mu; m_{\ell\ell} - m_Z > 10$ GeV for same-flavour leptons	$ee, \mu\mu, e\mu; m_{\ell\ell} - m_Z > 10$ GeV for same-flavour leptons
	SS- 2ℓ	ee , $\mu\mu$, or $e\mu$ with same charge	ee , $\mu\mu$, or $e\mu$ with same charge
	$\geq 3\ell$	$n_{\ell=e,\mu} \ge 3$	$n_{\ell=e,\mu} \ge 3$
	$\geq 2\tau$	$n_{ au,\mathrm{had}} \geq 2$	$n_{ au} \ge 2$
Photon	$1\gamma - m_{\gamma\gamma}^{12}$	$n_{\gamma} \geq 3$, $m_{\gamma\gamma}$ defined with γ_1, γ_2	$n_{\gamma} \geq 3, m_{\gamma\gamma}$ defined with γ_1, γ_2
Photon	1γ – $m_{\gamma\gamma}^{23}$	$n_{\gamma} \geq 3$, $m_{\gamma\gamma}$ defined with γ_2, γ_3	$n_{\gamma} \geq 3, m_{\gamma\gamma}$ defined with γ_2, γ_3

Table 3. Signal region definitions at the detector level and particle level. Particle-level object definitions are detailed in section 8.2. The $H_{\rm T}$ variable is defined as the scalar sum of the transverse momenta of jets that have $|\eta| < 2.5$. WP stands for efficiency working point. At the detector level, the τ -lepton identification is optimized for hadronically decaying τ -leptons. The symbols γ_1 , γ_2 , and γ_3 denote the leading, subleading, and third-leading photons ordered in $p_{\rm T}$, respectively. The BDT_{top} variable is a boosted decision tree (BDT) score used to identify triplets of jets consistent with the hadronic-decay final states of top quarks, which was used in ref. [19].

point. One exclusive hadronic top-quark signal region (t_{had}) requires the presence of exactly three jets, one of which must be a b-jet tagged at the 70% efficiency working point. The hadronic top-quark signal region also vetoes events with at least one electron or one muon. To further suppress diphoton-plus-multijet background, a boosted decision tree (BDT) is used in top-quark reconstruction [94] performed with the three jets in the event, and the resulting BDT score is required to be greater than 0.9.

6 Signal and background modelling

This search uses the invariant mass of the diphoton system $(m_{\gamma\gamma})$ as the discriminating variable. The potential BSM signal is a narrow resonance in $m_{\gamma\gamma}$, and the background consists of a resonant component and a continuum component. The resonant component arises from the SM Higgs boson production processes, which include gluon-gluon fusion (ggF) and vector-boson fusion (VBF), and VH, $t\bar{t}H$, tH, and $b\bar{b}H$ production.

The continuum component arises from the production of two initial- or final-state photons, or from the misidentification of jets as either one or both of the photons selected at the detector level. In signal regions where lepton and/or b-jet requirements are applied, these three types of continuum background processes also involve the production of electroweak bosons or top quarks.

The $m_{\gamma\gamma}$ distribution in each signal region is described by a probability density function (pdf) where the signal and background components are modelled by analytic functions of $m_{\gamma\gamma}$ and are normalized to their expected yields. As in previous ATLAS $H\to\gamma\gamma$ measurements [19, 95], the analytic functions are defined in the range of 105 GeV $< m_{\gamma\gamma} <$ 160 GeV. In the statistical interpretation of the search results, the $m_{\gamma\gamma}$ pdfs corresponding to the signal-plus-background and background-only hypotheses are used to fit the observed $m_{\gamma\gamma}$ distributions.

The event yield in the multilepton signal region is not sufficient for performing a fit to the $m_{\gamma\gamma}$ distribution. In this case, an event-count analysis is performed, as is described in section 6.3.

6.1 Modelling of BSM and SM Higgs boson production

The modelling of the $m_{\gamma\gamma}$ distribution and its associated systematic uncertainties is done in a consistent way for the BSM and SM Higgs boson production processes. Their $m_{\gamma\gamma}$ distributions are both described by the same double-sided Crystal Ball (DSCB) function [96, 97], consisting of a Gaussian distribution in the region around the peak position, continued by power-law tails at lower and higher $m_{\gamma\gamma}$ values. The potential bias in the estimated signal yield due to an intrinsic shape difference between the DSCB function and signal $m_{\gamma\gamma}$ distribution is found to be much smaller than the MC statistical uncertainty of the signal yield and therefore is negligible [98]. The parameters of the DSCB function in each signal region are obtained by a fit to a mixture of the ggF, VBF. VH, tH, and bbH samples described in section 3, which are normalized according to their SM cross sections. A shift of 0.09 GeV is applied to the position of the signal peak to account for the difference between the reference Higgs boson mass used in this analysis ($m_H = 125.09 \,\text{GeV}$) [99] and the mass for which the samples were generated $(m_H = 125 \,\text{GeV})$. The uncertainty in the $m_{\gamma\gamma}$ modelling is discussed in section 7.1. There is a modest variation in the DSCB shape parameters between various SM and BSM Higgs boson production samples. However, the impact of this variation on the fitted number of signal events is much smaller than the statistical uncertainty of the fitted signal yield.

SM Higgs boson production is the resonant background to signals arising from BSM production. The cross sections of the resonant backgrounds are assumed to be the same as

the theoretical predictions, and their contributions to various signal regions are evaluated using the MC samples described in section 3. The recent Higgs boson measurements from the ATLAS and CMS experiments have shown that the data are consistent with the theory predictions within statistical and systematic uncertainties, which justifies the fixing of expected resonant background yields to their SM predictions. Potential interference effects between the SM and BSM Higgs boson production processes are not considered in the MC simulation. If the interference effect results in a significant deviation in the production rate from the SM prediction, this search would detect it as a signal.

6.2 Continuum background modelling

The $m_{\gamma\gamma}$ distribution of the continuum background is fully determined from a fit to data. The only systematic uncertainty associated with this method arises from the intrinsic difference between the true continuum background $m_{\gamma\gamma}$ shape and the shape of the chosen analytic pdf.

The modelling of the continuum background follows the same process used in ref. [98], which involves two main steps: first, a background $m_{\gamma\gamma}$ template is constructed from either the MC diphoton continuum background or a data control sample where at least one of the two leading photons fails the photon identification or isolation requirement. Between these two samples, the one with a higher number of events is selected to construct the $m_{\gamma\gamma}$ template. Second, a background function is selected from a number of candidate functions, using a procedure known as the *spurious-signal test* [98], with the goal of identifying an analytic function that is flexible enough to fit the $m_{\gamma\gamma}$ distribution in data and which results in a small potential bias compared to the background statistical uncertainty.

Several families of analytic functions are tested as candidates to model the $m_{\gamma\gamma}$ distribution for each signal region. They include power-law functions, Bernstein polynomials, and exponential functions of a polynomial. These functional forms can include up to five free parameters. The selected functional forms are mostly power-law and exponential functions with one free parameter. The coefficients of background functions are considered independent across signal regions, regardless of the functions chosen. In all cases, they are treated as free parameters in the fit to data.

6.3 Modelling of the background in the multilepton signal region

The expected continuum background event yield in the multilepton $(\geq 3\ell)$ signal region is so small that the $m_{\gamma\gamma}$ distribution may not be fitted with an analytic function. In this case, the event count in the $m_{\gamma\gamma}$ range from 123 GeV to 127 GeV is used for statistical interpretation. The expected background yield in this mass range is extrapolated from a control region in two steps. This control region is referred to as the non-tight-isolated region and is defined in the same way as the multilepton signal region, except that at least one of the two leading photons must fail to meet the photon quality criteria detailed in section 4. The expected background yield in the diphoton mass sidebands of the multilepton signal region is scaled from the event yield in the non-tight-isolated diphoton mass sidebands. The ratio of the event yield in the non-tight-isolated control region to that in the tight-isolated signal region is checked in the other leptonic signal regions and is found to be

consistent, which indicates that the scale factor derived from other leptonic signal regions can be applied to the multilepton signal region. The variations in the scale factor between various leptonic signal regions are considered as systematic uncertainties of the scale factor. Similarly, the expected background yield in the mass window of 123 GeV $< m_{\gamma\gamma} <$ 127 GeV is computed from the expected background yield in the diphoton mass sidebands, using scale factors evaluated in the non-tight-isolated control regions defined for various leptonic signal regions. These scale factors are consistent and their variations are considered as systematic uncertainties of the extrapolation. The expected continuum background yield in the multilepton signal region is found to be $0.25^{+0.59}_{-0.25}$ events, where the lower bound of the uncertainty is truncated so that the yield remains positive. The expected resonant background yield in this region is predicted by the MC simulation.

7 Systematic uncertainties

Systematic uncertainties considered in this analysis can be grouped into three categories: uncertainties in the modelling of the $m_{\gamma\gamma}$ distribution for the BSM and SM Higgs boson production processes, uncertainties in the modelling of the continuum background $m_{\gamma\gamma}$ distribution, and uncertainties in the expected resonant background yields in each signal region arising from either experimental or theoretical sources. These systematic uncertainties are incorporated into the analysis likelihood models as nuisance parameters.

7.1 Systematic uncertainties in modelling $m_{\gamma\gamma}$ for SM and BSM Higgs boson production

In general, the $m_{\gamma\gamma}$ modelling systematic uncertainties for a given region depend on the BSM signal considered. However, the signal shape variations associated with the properties of a specific BSM model are negligible relative to those associated with the experimental effects, and only the latter are therefore considered and estimated from the SM Higgs production processes. The extraction of a potential signal component from the data $m_{\gamma\gamma}$ distribution is subject to systematic uncertainties in the diphoton mass resolution and scale. While the diphoton mass resolution and scale uncertainties are also model-dependent, their variations are typically limited. The likelihood model considers only signal systematic uncertainties related to the diphoton mass scale and resolution, and does not include any systematic uncertainty for the signal yield, which is the parameter of interest in the search.

The diphoton mass scale uncertainty includes a contribution from the photon energy scale uncertainty, which is typically less than 1% of the diphoton mass, and also a systematic component of 240 MeV, corresponding to the measurement uncertainty of the Higgs boson mass from the combination of the ATLAS and CMS data [99]. The diphoton mass resolution uncertainty arises from the uncertainty of the photon energy resolution and is about 3%–15% relative to the reconstructed width of the diphoton resonance. The estimation and implementation of the photon energy scale and resolution uncertainties follow the procedure outlined in ref. [79].

7.2 Uncertainty in the continuum background modelling

The uncertainty in the modelling arises from an intrinsic shape difference between the continuum background $m_{\gamma\gamma}$ distribution and the chosen analytic function. The size of this uncertainty is estimated using the *spurious-signal* test mentioned in section 6.2 and is always smaller than 20% of the statistical uncertainty of the continuum background.

7.3 Uncertainties in resonant background yields

The definition of the signal regions includes selections based on photons, leptons, jets (b-jets), and missing transverse momentum. As such, the expected resonant background yield is subject to experimental systematic uncertainties in the reconstruction and identification efficiencies of these objects. The typical level of systematic uncertainty for object reconstruction and identification is less than 1% for photons [79], 1%–4% for electrons [79] and muons [100], 2%–16% for jets [101] and b-jets [91], and 2% for τ -leptons [102]. Uncertainties in the theoretical modelling of SM Higgs boson production processes are estimated by varying the scale of QCD renormalization and factorization, by changing the choice of parton distribution function set, and by comparing the predicted events from different parton showering implementations for a given Higgs boson production process. The theoretical systematic uncertainties typically range from 10% to 35% [18]. The uncertainty in the integrated luminosity of the data sample is 1.7% [27], obtained using the LUCID-2 detector [28] for the primary luminosity measurements. Other uncertainties in background event yields, such as the impact of pile-up on selection efficiencies [103], contribute < 1%.

8 Results

Figures 1–6 show the diphoton mass $m_{\gamma\gamma}$ distributions for all the signal regions considered in this paper, except the multilepton ($\geq 3\ell$) signal region. The result of a signal-plus-background fit as described in section 6 is shown for each signal region. The expected and observed event yields are determined from the signal-plus-background fit in each signal region and are summarized in table 4. Also included in this table are the expected and observed event counts in the multilepton signal region. No event is found in the multilepton signal region, while the sum of the continuum and resonant background expectations is $0.27^{+0.59}_{-0.27}$ events . A p_0 -value was evaluated for each of these signal regions, and the largest excess is in the $H_T > 1000$ GeV region and is less than two standard deviations above the expectation. The observed significance [104] is also given in table 4.

SR	Expected Background			Observed		
SIL	Resonant Higgs	Continuum	Total Background	Observed Yield	Excess Significance $[\sigma]$	
		1	Heavy flavour			
$\geq 3b$	6.47	23.4	29.9	30	-0.3	
$\geq 4b$	0.69	1.22	1.91	1	-0.2	
		H	igh jet activity			
≥4j	85.2	1330	1420	1404	-0.3	
≥6j	16.4	104	121	105	-1.3	
≥8j	2.44	6.37	8.81	6	-0.9	
$H_{\rm T} > 500 \; {\rm GeV}$	23.9	297	321	310	-0.6	
$H_{\rm T} > 1000 \; {\rm GeV}$	1.85	27	28.8	39	1.8	
$H_{\rm T} > 1500 \mathrm{GeV}$	0.264	3.9	4.17	4	0.1	
			$E_{\mathrm{T}}^{\mathrm{miss}}$			
$E_{\rm T}^{\rm miss} > 100 \text{ GeV}$	29	171	200	212	0.8	
$E_{\rm T}^{\rm miss} > 200 \text{ GeV}$	4.51	8.06	12.6	16	0.9	
$E_{\rm T}^{\rm miss} > 300 \text{ GeV}$	1.15	1.85	3	5	0.8	
			Top quark			
ℓb	14.9	27	41.9	34	-0.6	
t_{lep}	0.281	2.58	2.86	1	-0.7	
$t_{ m had}$	4.44	96.3	101	111	1.7	
			Lepton			
$\geq 1\ell$	38.8	183	222	237	1.4	
2ℓ	4.24	9.42	13.7	10	-0.5	
2ℓ – Z	1.95	7.35	9.3	10	0.7	
SS- 2ℓ	0.431	0.224	0.655	1	0.2	
$\geq 3\ell$	0.02	0.25	0.27	0	_	
$\geq 2\tau$	0.256	0.875	1.13	2	0.6	
			Photon			
1γ – $m_{\gamma\gamma}^{12}$	2.33	119	121	132	0.7	
$1\gamma - m_{\gamma\gamma}^{23}$	0.436	32.8	33.2	42	1.1	

Table 4. Expected background and observed event yields in the diphoton mass range 123 GeV < $m_{\gamma\gamma} < 127$ GeV. The resonant Higgs boson production and continuum backgrounds are shown separately. The observed significance is also shown for all signal regions except the $\geq 3\ell$ region, where less than one background event is expected and the observed number of events is zero. Note that the significance is calculated using the full mass spectrum of $105\,\mathrm{GeV} < m_{\gamma\gamma} < 160\,\mathrm{GeV}$, which may not be reproduced using yields reported in this table.




Figure 1. Diphoton mass distributions for heavy-flavour signal regions. Data are shown together with the signal-plus-background fit. The overall fitted signal-plus-background pdf is shown as a solid red curve, the contributions from the resonant Higgs boson and continuum backgrounds are shown as a dashed blue curve, while the contributions only from the continuum backgrounds are shown as a dashed green curve. The difference between the solid red and the dashed blue curves indicates the fitted deviation from the SM expectation.




Figure 2. Diphoton mass distributions for high-jet-activity signal regions. Data are shown together with the signal-plus-background fit. The overall fitted signal-plus-background pdf is shown as a solid red curve, the contributions from the resonant Higgs boson and continuum backgrounds are shown as a dashed blue curve, while the contributions only from the continuum backgrounds are shown as a dashed green curve. The difference between the solid red and the dashed blue curves indicates the fitted deviation from the SM expectation.




Figure 3. Diphoton mass distributions for $E_{\rm T}^{\rm miss}$ signal regions. Data are shown together with the signal-plus-background fit. The overall fitted signal-plus-background pdf is shown as a solid red curve, the contributions from the resonant Higgs boson and continuum backgrounds are shown as a dashed blue curve, while the contributions only from the continuum backgrounds are shown as a dashed green curve. The difference between the solid red and the dashed blue curves indicates the fitted deviation from the SM expectation.




Figure 4. Diphoton mass distributions for lepton signal regions. Data are shown together with the signal-plus-background fit. The overall fitted signal-plus-background pdf is shown as a solid red curve, the contributions from the resonant Higgs boson and continuum backgrounds are shown as a dashed blue curve, while the contributions only from the continuum backgrounds are shown as a dashed green curve. The difference between the solid red and the dashed blue curves indicates the fitted deviation from the SM expectation.




Figure 5. Diphoton mass distributions for top-quark signal regions. Data are shown together with the signal-plus-background fit. The overall fitted signal-plus-background pdf is shown as a solid red curve, the contributions from the resonant Higgs boson and continuum backgrounds are shown as a dashed blue curve, while the contributions only from the continuum backgrounds are shown as a dashed green curve. The difference between the solid red and the dashed blue curves indicates the fitted deviation from the SM expectation.




Figure 6. Diphoton mass distributions for photon signal regions. Data are shown together with the signal-plus-background fit. The overall fitted signal-plus-background pdf is shown as a solid red curve, the contributions from the resonant Higgs boson and continuum backgrounds are shown as a dashed blue curve, while the contributions only from the continuum backgrounds are shown as a dashed green curve. The difference between the solid red and the dashed blue curves indicates the fitted deviation from the SM expectation.

8.1 Upper limits on BSM production cross sections

Since no significant excess beyond the SM expectation was observed, the $m_{\gamma\gamma}$ distributions in the signal regions, as well as the event count in the multilepton $(\geq 3\ell)$ region, are used to set limits on the visible cross section of BSM production of the Higgs boson with a mass of 125.09 GeV in these regions. The 95% CL limits were derived using the modified frequentist CL_s method [105]. The limits are shown in table 5 and graphically in figure 7. The observed 95% CL limit on the visible cross section ranges from 0.05 fb $(\geq 3\ell)$ to 0.7 fb $(\geq 4j)$.

	95% CL limit [fb]					
SR	Observed	Median	$+1\sigma$	-1σ	$+2\sigma$	-2σ
	Н	leavy flavo	ur			
$\geq 3b$	0.12	0.13	0.18	0.093	0.25	0.069
$\geq 4b$	0.037	0.037	0.054	0.026	0.081	0.02
	Hi	gh jet acti	vity			
≥4j	0.73	0.81	1.1	0.58	1.5	0.43
≥6j	0.15	0.22	0.31	0.16	0.42	0.12
≥8j	0.047	0.065	0.094	0.047	0.13	0.035
$H_{\rm T} > 500~{\rm GeV}$	0.28	0.34	0.47	0.24	0.64	0.18
$H_{\mathrm{T}} > 1000 \; \mathrm{GeV}$	0.21	0.11	0.16	0.079	0.22	0.059
$H_{\rm T} > 1500~{\rm GeV}$	0.052	0.048	0.07	0.034	0.1	0.026
		$E_{\mathrm{T}}^{\mathrm{miss}}$				
$E_{\mathrm{T}}^{\mathrm{miss}} > 100 \text{ GeV}$	0.4	0.3	0.42	0.22	0.57	0.16
$E_{\mathrm{T}}^{\mathrm{miss}} > 200 \text{ GeV}$	0.11	0.075	0.11	0.054	0.15	0.04
$E_{\mathrm{T}}^{\mathrm{miss}} > 300 \text{ GeV}$	0.053	0.04	0.059	0.029	0.088	0.022
		Top				
ℓb	0.14	0.16	0.22	0.11	0.29	0.085
$t_{ m lep}$	0.034	0.043	0.062	0.031	0.092	0.023
$t_{ m had}$	0.39	0.23	0.32	0.17	0.44	0.12
		Lepton				
$\geq 1\ell$	0.5	0.32	0.45	0.23	0.61	0.17
2ℓ	0.069	0.08	0.11	0.057	0.16	0.043
2ℓ – Z'	0.089	0.068	0.098	0.049	0.14	0.037
SS- 2ℓ	0.028	0.026	0.04	0.019	0.061	0.014
$\geq 3\ell$	0.022	0.022	0.028	0.022	0.046	0.022
$\geq 2\tau$	0.036	0.031	0.046	0.022	0.07	0.016
Photon						
1γ – $m_{\gamma\gamma}^{12}$	0.44	0.35	0.49	0.25	0.66	0.19
1γ – $m_{\gamma\gamma}^{23}$	0.4	0.28	0.4	0.2	0.55	0.15

Table 5. Observed and expected 95% CL upper limits on the visible cross section of BSM Higgs boson production in each signal region. The $\pm 1\sigma$ and $\pm 2\sigma$ variations of the expected limit are also shown.

The observed limits reported here are correlated since the signal regions overlap. To enable the use of limits from multiple signal regions to constrain BSM processes, the correlation between the observed limits from any pair of signal regions considered in this paper is estimated using a pseudo-experiment-based bootstrapping procedure [106]. Readers of this paper could derive a combined limit on the cross section of a BSM signal using the observed limits in individual signal regions and their correlations. Only regions with at least 50 events in the $m_{\gamma\gamma}$ range from 105 GeV to 160 GeV were checked using this procedure. Correlations larger than 5% are observed for a few regions and are reported in table 6.




Figure 7. Observed and expected 95% CL limits on the visible cross section of BSM Higgs boson production in each signal region. The 1σ and 2σ variations of the expected limit are also shown as green and yellow bands, respectively.

8.2 Interpretation with particle-level simulation

This paper aims to present results that can be used without detailed knowledge of the ATLAS detector to derive constraints on the BSM processes of interest. For a specific BSM process,

Region	Region (Correlation)
≥4j	≥ 6 j (29%), $H_T > 500$ GeV (28%)
≥6j	$\geq 8j (25\%), H_T > 500 \text{ GeV } (26\%), H_T > 1000 \text{ GeV } (8\%)$
$H_{\mathrm{T}} > 500 \mathrm{GeV}$	$H_{\rm T} > 1000 \; {\rm GeV} \; (32\%)$
$E_{\mathrm{T}}^{\mathrm{miss}} > 200\mathrm{GeV}$	$E_{\rm T}^{\rm miss} > 100 {\rm GeV} (29\%)$
1ℓ	2ℓ (26%), ℓb (36%)
2ℓ	2ℓ – Z (97%)

Table 6. Correlations between observed 95% CL limits for pairs of signal regions. Only regions with a correlation greater than 5% are reported. Pairs of regions that appear in previous rows are not repeated.

Object	Requirements		
Photon	$p_{\rm T} > 22 {\rm GeV}, \eta < 2.5$		
Electron	$p_{\rm T} > 10 {\rm GeV}, \eta < 2.5$		
Muon	$p_{\rm T} > 10 \; {\rm GeV}, \; \eta < 2.7$		
Jet	$p_{\rm T} > 25 {\rm GeV}, \eta < 4.4$		
b-jet	$p_{\rm T} > 25 \; {\rm GeV}, \; \eta < 2.5, \; \Delta R(b\text{-quark, jet}) < 0.4$		
au-lepton	$p_{\rm T} > 20 \; {\rm GeV}, \; \eta < 2.5$		
Preselection			
$p_{\rm T}^{\gamma 1} > 35 \text{ GeV}, p_{\rm T}^{\gamma 2} > 25 \text{ GeV}, p_{\rm T}^{\gamma 1}/m_{\gamma \gamma} > 0.35, p_{\rm T}^{\gamma 2}/m_{\gamma \gamma} > 0.25$			

Table 7. Particle-level object definitions and event preselection.

its particle-level cross section, $\sigma_{\rm tru}$, in a given region can be derived from MC simulation with particle-level requirements as given in table 3. To determine if the BSM process is excluded or not, $\sigma_{\rm tru}$ should be compared with a particle-level 95% CL limit, $\sigma_{\rm tru}^{95}$, for the same region. Since this paper only reports a 95% CL limit on the visible cross section at the detector level, $\sigma_{\rm det}^{95}$, for any of the signal regions, a signal-region-specific detector efficiency, ϵ , must be provided to convert $\sigma_{\rm det}^{95}$ to $\sigma_{\rm tru}^{95}$. More specifically, one has $\sigma_{\rm tru}^{95} = \sigma_{\rm det}^{95}/\epsilon$.

The region-specific detector efficiency ϵ is defined as the ratio of the event yield in a signal region at the detector level to that in a signal region defined at the particle level. As such, for each signal region considered in this paper, a particle-level definition must be specified. Table 7 summarizes the particle-level object definitions. Particle-level jets are built from all MC generator-level "truth" particles, removing the contributions from muons and stable particles such as neutrinos and lightest supersymmetric particles, using the anti- k_t algorithm with a radius parameter of 0.4. The particle-level objects are required to be isolated by implementing the following overlap removal procedure: electrons, muons, photons, and τ -leptons are removed if there is another object of the same type within $\Delta R < 0.1$. Then any muon, jet or electron is removed if it is within $\Delta R = 0.4$ of any photon, any jet within $\Delta R = 0.2$ of any electron is removed, and finally any muon or electron within $\Delta R = 0.4$ of any jet is removed. In this paper, a particle-level b-jet is defined as a jet that is within $\Delta R = 0.4$ of a b-quark, and the b-quark should pass the requirements of $p_{\rm T} > 25$ GeV and $|\eta| < 2.5$. This requirement is not fully efficient due to the kinematic requirements on the b-quark. Depending on the specific signal process, the requirement has an efficiency of matching a particle-level jet to a b-quark at the level of 70% to 80%, for a b-quark with $p_T > 25$ GeV and $|\eta| < 2.5$. Then, the particle-level signal regions can be defined with the particle-level objects, and the definitions are detailed in table 3.

The detector efficiency, ϵ , is evaluated for all signal regions considered in this paper and is summarized in table 8. For each signal region, the value of ϵ is calculated for a few SM and BSM Higgs boson production processes, and the range of ϵ values is reported. Since the final-state object multiplicity and kinematics can differ between processes, variations in ϵ are expected. The reported ϵ values are subject to a small statistical uncertainty, which

is much smaller than the variation in ϵ between different physics processes. In general, the detector-level selection is always less efficient than the particle-level selection, except for the following two situations. First, when selecting a particle-level b-jet, the ΔR matching procedure may be less efficient than the detector-level b-tagging, for which a b-jet is defined as a jet containing a b-hadron. The adoption of the ΔR matching procedure is motivated by the convenience of implementation at the particle level. Second, when a signal region is defined by placing a cut at a value in the tail of a variable's distribution, such as for $H_{\rm T}$ or the number of jets, detector-level effects could lead to event migration and hence a higher detector-level selection efficiency. However, when all the detector effects are combined, as shown in table 8, ϵ is always less than 1. The value of ϵ is very small for the 2τ signal region because the particle-level τ -leptons include all τ -leptons regardless of their decay modes. This choice is also motivated by the convenience of selection implementation at the particle level.

As an example, an electroweak SUSY production model, corresponding to the one shown in the first row of table 2 is used to demonstrate how information reported in this section can be used to derive a constraint on a given BSM model. In this SUSY process, a chargino and neutralino pair $(\tilde{\chi}_1^{\pm}\tilde{\chi}_2^0)$ is produced in the pp collision, which results in a final state of a W boson, a Higgs boson, and two light neutralinos, $\tilde{\chi}_1^0$. The light neutralinos are undetectable, giving rise to large values of $E_{\rm T}^{\rm miss}$. The masses of the pair-produced chargino and neutralino are 150 GeV, and the light neutralino's mass is 0.5 GeV. For this model, a targeted search has already been carried out by the ATLAS Collaboration [69], which reported a 95% CL upper limit of 2 pb for the cross section at this specific signal point.

The $E_{\rm T}^{\rm miss}>100$ GeV signal region can be used to probe the same SUSY process, which predicts a particle-level cross section, $\sigma_{\rm tru}$, of 1.05 fb for this $E_{\rm T}^{\rm miss}>100$ GeV region. The observed 95% CL upper limit on the BSM cross section at the detector level is $\sigma_{\rm det}^{95}=0.4$ fb, for the $E_{\rm T}^{\rm miss}>100$ GeV region, and its corresponding ϵ is chosen, conservatively, to be 0.6, according to table 8. The particle-level 95% CL limit on the BSM cross section in the $E_{\rm T}^{\rm miss}>100$ GeV region is therefore $\sigma_{\rm tru}^{95}=0.67$ fb, which would exclude the SUSY model considered here. The observed limit in the $E_{\rm T}^{\rm miss}>100$ GeV region in the $H\to\gamma\gamma$ final state can be converted to a limit on the inclusive cross section for this SUSY model by taking into account the selection acceptance at the particle level and the $H\to\gamma\gamma$ branching ratio, and the resulting inclusive cross-section limit is 3.3 pb. This limit is of the same order of magnitude as the 2 pb upper limit in the dedicated search, but is 65% weaker.

SR	Relevant processes	Range of ϵ
	Heavy flavour	
$\geq 3b$	$ ilde{b} ilde{b}, ilde{b} o\chi_2^0 b, ilde{\chi}_2^0 o\chi_1^0 H, ilde{t}_2 ilde{t}_2, ilde{t}_2 o ilde{t}_1 H, ilde{t}_1 o ilde{\chi}_1^0 bqar{q}/b\ell u$	0.68 – 0.81
$\ge 4b$	$\tilde{b}\tilde{b},\tilde{b}\to\tilde{\chi}_2^0b,\tilde{\chi}_2^0\to\tilde{\chi}_1^0H,\tilde{t}_2\tilde{t}_2,\tilde{t}_2\to\tilde{t}_1H,\tilde{t}_1\to\tilde{\chi}_1^0bq\bar{q}/b\ell\nu$	0.64 – 0.97
	High jet activity	
≥4j	$t\bar{t}H,\tilde{b}\tilde{b},\tilde{b}\to\tilde{\chi}^0_2b,\tilde{\chi}^0_2\to\tilde{\chi}^0_1H,\tilde{t}_2\tilde{t}_2,\tilde{t}_2\to\tilde{t}_1H,\tilde{t}_1\to\tilde{\chi}^0_1bq\bar{q}/b\ell\nu,tWH$	0.60-0.70
≥6j	$t\bar{t}H,\ \tilde{b}\tilde{b},\tilde{b}\to\tilde{\chi}^0_2b,\tilde{\chi}^0_2\to\tilde{\chi}^0_1H,\ \tilde{t}_2\tilde{t}_2,\tilde{t}_2\to\tilde{t}_1H,\tilde{t}_1\to\tilde{\chi}^0_1bq\bar{q}/b\ell\nu,\ tWH$	0.64 – 0.80
≥8j	$t\bar{t}H,\tilde{b}\tilde{b},\tilde{b}\to\tilde{\chi}_2^0b,\tilde{\chi}_2^0\to\tilde{\chi}_1^0H,\tilde{t}_2\tilde{t}_2,\tilde{t}_2\to\tilde{t}_1H,\tilde{t}_1\to\tilde{\chi}_1^0bq\bar{q}/b\ell\nu$	0.65 – 0.90
$H_{\rm T} > 500 \; {\rm GeV}$	$t\bar{t}H,\tilde{b}\tilde{b},\tilde{b}\to\tilde{\chi}^0_2b,\tilde{\chi}^0_2\to\tilde{\chi}^0_1H,\tilde{t}_2\tilde{t}_2,\tilde{t}_2\to\tilde{t}_1H,\tilde{t}_1\to\tilde{\chi}^0_1bq\bar{q}/b\ell\nu,tWH$	0.52 – 0.66
$H_{\rm T} > 1000~{\rm GeV}$	$t\bar{t}H,\tilde{b}\tilde{b},\tilde{b}\to\tilde{\chi}_2^0b,\tilde{\chi}_2^0\to\tilde{\chi}_1^0H,\tilde{t}_2\tilde{t}_2,\tilde{t}_2\to\tilde{t}_1H,\tilde{t}_1\to\tilde{\chi}_1^0bq\bar{q}/b\ell\nu,tWH$	0.51 – 0.72
$H_{\rm T} > 1500 \; {\rm GeV}$	$t\bar{t}H,\tilde{b}\tilde{b},\tilde{b}\to\tilde{\chi}_2^0b,\tilde{\chi}_2^0\to\tilde{\chi}_1^0H,\tilde{t}_2\tilde{t}_2,\tilde{t}_2\to\tilde{t}_1H,\tilde{t}_1\to\tilde{\chi}_1^0bq\bar{q}/b\ell\nu$	0.41 – 0.73
	$E_{ m T}^{ m miss}$	
$E_{\mathrm{T}}^{\mathrm{miss}} > 100 \text{ GeV}$	$t\bar{t}H,tWH,WH,ZH,\tilde{\chi}_1^{\pm}\tilde{\chi}_2^0 \to W/Z/H$	0.60-0.78
$E_{\mathrm{T}}^{\mathrm{miss}} > 200 \text{ GeV}$	$tar{t}H,tWH,WH,ZH, ilde{\chi}_1^\pm ilde{\chi}_2^0\! ightarrow\!W/Z/H$	0.60 – 0.79
$E_{\mathrm{T}}^{\mathrm{miss}} > 300 \ \mathrm{GeV}$	$tar{t}H,tWH,WH,ZH, ilde{\chi}_1^\pm ilde{\chi}_2^0{ ightarrow}W/Z/H$	0.66 - 0.84
	Lepton	
≥1ℓ	$WH, t\bar{t}H, tWH, \text{FCNC}, \tilde{\chi}_1^{\pm} \tilde{\chi}_2^0 \rightarrow W/Z/H$	0.40-0.48
2ℓ	$ZH,t\bar{t}H,\tilde{t}_2\tilde{t}_2,\tilde{t}_2\rightarrow\tilde{t}_1H,\tilde{t}_1\rightarrow\tilde{\chi}^0_1bq\bar{q}/b\ell\nu,\tilde{\chi}^\pm_1\tilde{\chi}^\mp_1\rightarrow H\ell^\pm H\ell^\mp,\tilde{\chi}^\pm_1\tilde{\chi}^0_1,\tilde{\chi}^\pm_1\rightarrow H\ell^\pm,\tilde{\chi}^0_1\rightarrow W\ell/Z\nu/H\nu$	0.21 – 0.48
2ℓ – Z	$t\bar{t}H,\tilde{t}_2\tilde{t}_2,\tilde{t}_2\rightarrow\tilde{t}_1H,\tilde{t}_1\rightarrow\tilde{\chi}_1^0bq\bar{q}/b\ell\nu,\tilde{\chi}_1^\pm\tilde{\chi}_1^\mp\rightarrow H\ell^\pm H\ell^\mp,\tilde{\chi}_1^\pm\tilde{\chi}_1^0,\tilde{\chi}_1^\pm\rightarrow H\ell^\pm,\tilde{\chi}_1^0\rightarrow W\ell/Z\nu/H\nu$	0.20 – 0.46
$\geq 3\ell$	$\tilde{t}_2\tilde{t}_2,\tilde{t}_2\rightarrow \tilde{t}_1H,\tilde{t}_1\rightarrow \tilde{\chi}_1^0bq\bar{q}/b\ell\nu,~\tilde{\chi}_1^{\pm}\tilde{\chi}_1^{\mp}\rightarrow H\ell^{\pm}H\ell^{\mp},~\tilde{\chi}_1^{\pm}\tilde{\chi}_1^0,\tilde{\chi}_1^{\pm}\rightarrow H\ell^{\pm},\tilde{\chi}_1^0\rightarrow W\ell/Z\nu/H\nu$	0.18 – 0.33
SS- 2ℓ	$\tilde{t}_2\tilde{t}_2,\tilde{t}_2\rightarrow \tilde{t}_1H,\tilde{t}_1\rightarrow \tilde{\chi}_1^0bq\bar{q}/b\ell\nu,~\tilde{\chi}_1^{\pm}\tilde{\chi}_1^0,\tilde{\chi}_1^{\pm}\rightarrow H\ell^{\pm},\tilde{\chi}_1^0\rightarrow W\ell/Z\nu/H\nu$	0.29 – 0.49
$\geq 2\tau$	$ZH,\ \tilde{t}_2\tilde{t}_2,\tilde{t}_2\rightarrow\tilde{t}_1H,\tilde{t}_1\rightarrow\tilde{\chi}_1^0bq\bar{q}/b\ell\nu,\ \tilde{b}\tilde{b},\tilde{b}\rightarrow\tilde{\chi}_2^0b,\tilde{\chi}_2^0\rightarrow\tilde{\chi}_1^0H,\ \tilde{\chi}_1^\pm\tilde{\chi}_1^\mp\rightarrow H\ell^\pm H\ell^\mp,\ \tilde{\chi}_1^\pm\tilde{\chi}_1^0,\tilde{\chi}_1^\pm\rightarrow H\ell^\pm,\tilde{\chi}_1^0\rightarrow W\ell/Z\nu/H\nu$	0.04 – 0.09
	Top quark	
$t_{\rm lep}$	FCNC with semileptonically decaying top	0.32-0.36
$t_{\rm had}$	FCNC with hadronically decaying top	0.29 – 0.30
ℓb	$t\bar{t}H,tHjb,tWH,\mathrm{FCNC}$ with semileptonically decaying top	0.41 – 0.52
	Photon	
1γ - $m_{\gamma\gamma}^{12}$	$\tilde{t}_2\tilde{t}_2,\tilde{t}_2\to\tilde{t}_1H,\tilde{t}_1\to\tilde{\chi}_1^0bq\bar{q}/b\ell\nu,\tilde{b}\tilde{b},\tilde{b}\to\tilde{\chi}_2^0b,\tilde{\chi}_2^0\to\tilde{\chi}_1^0H$	0.23 - 0.33
$1\gamma - m_{\gamma\gamma}^{23}$	$ ilde{t}_2 ilde{t}_2, ilde{t}_2 ightarrow ilde{t}_1H, ilde{t}_1 ightarrow ilde{\chi}_1^0bqar{q}/b\ell u$	0.35 - 0.40

Table 8. Range of ϵ values for all SRs. The detector efficiency ϵ is only calculated for processes where the reconstruction-level yield in a given signal region is greater than 0.1 events for 139 fb⁻¹ of data.

9 Conclusions

This paper reports a model-independent examination of the diphoton final states where a Higgs boson is produced, using a 139 fb $^{-1}$ data sample of 13 TeV pp collisions collected by the ATLAS detector during Run 2 of the LHC. A total of 22 signal regions defined with various requirements on the additional reconstructed objects were examined, and no significant excess over the SM expectation was observed. Upper limits at 95% CL were set on the visible cross section of BSM Higgs boson production in each of these signal regions. To enable interpretation of the results as a constraint on other models, a set of detector efficiency factors is also reported.

Acknowledgments

We thank CERN for the very successful operation of the LHC, as well as the support staff from our institutions without whom ATLAS could not be operated efficiently.

We acknowledge the support of ANPCyT, Argentina; YerPhI, Armenia; ARC, Australia; BMWFW and FWF, Austria; ANAS, Azerbaijan; CNPq and FAPESP, Brazil; NSERC, NRC and CFI, Canada; CERN; ANID, Chile; CAS, MOST and NSFC, China; Minciencias, Colombia; MEYS CR, Czech Republic; DNRF and DNSRC, Denmark; IN2P3-CNRS

and CEA-DRF/IRFU, France; SRNSFG, Georgia; BMBF, HGF and MPG, Germany; GSRI, Greece; RGC and Hong Kong SAR, China; ISF and Benoziyo Center, Israel; INFN, Italy; MEXT and JSPS, Japan; CNRST, Morocco; NWO, Netherlands; RCN, Norway; MEiN, Poland; FCT, Portugal; MNE/IFA, Romania; MESTD, Serbia; MSSR, Slovakia; ARRS and MIZŠ, Slovenia; DSI/NRF, South Africa; MICINN, Spain; SRC and Wallenberg Foundation, Sweden; SERI, SNSF and Cantons of Bern and Geneva, Switzerland; MOST, Taiwan; TENMAK, Türkiye; STFC, United Kingdom; DOE and NSF, United States of America. In addition, individual groups and members have received support from BCKDF, CANARIE, Compute Canada and CRC, Canada; PRIMUS 21/SCI/017 and UNCE SCI/013, Czech Republic; COST, ERC, ERDF, Horizon 2020 and Marie Skłodowska-Curie Actions, European Union; Investissements d'Avenir Labex, Investissements d'Avenir Idex and ANR, France; DFG and AvH Foundation, Germany; Herakleitos, Thales and Aristeia programmes co-financed by EU-ESF and the Greek NSRF, Greece; BSF-NSF and MINERVA, Israel; Norwegian Financial Mechanism 2014-2021, Norway; NCN and NAWA, Poland; La Caixa Banking Foundation, CERCA Programme Generalitat de Catalunya and PROMETEO and GenT Programmes Generalitat Valenciana, Spain; Göran Gustafssons Stiftelse, Sweden; The Royal Society and Leverhulme Trust, United Kingdom.

The crucial computing support from all WLCG partners is acknowledged gratefully, in particular from CERN, the ATLAS Tier-1 facilities at TRIUMF (Canada), NDGF (Denmark, Norway, Sweden), CC-IN2P3 (France), KIT/GridKA (Germany), INFN-CNAF (Italy), NL-T1 (Netherlands), PIC (Spain), ASGC (Taiwan), RAL (U.K.) and BNL (U.S.A.), the Tier-2 facilities worldwide and large non-WLCG resource providers. Major contributors of computing resources are listed in ref. [107].

Open Access. This article is distributed under the terms of the Creative Commons Attribution License (CC-BY 4.0), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] ATLAS collaboration, Observation of a new particle in the search for the standard model Higgs boson with the ATLAS detector at the LHC, Phys. Lett. B 716 (2012) 1 [arXiv:1207.7214] [INSPIRE].
- [2] CMS collaboration, Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC, Phys. Lett. B 716 (2012) 30 [arXiv:1207.7235] [INSPIRE].
- [3] Y.A. Golfand and E.P. Likhtman, Extension of the algebra of Poincaré group generators and violation of p invariance, JETP Lett. 13 (1971) 323 [INSPIRE].
- [4] D.V. Volkov and V.P. Akulov, Is the neutrino a Goldstone particle?, Phys. Lett. B 46 (1973) 109 [INSPIRE].
- [5] J. Wess and B. Zumino, Supergauge transformations in four-dimensions, Nucl. Phys. B 70 (1974) 39 [INSPIRE].
- [6] J. Wess and B. Zumino, Supergauge invariant extension of quantum electrodynamics, Nucl. Phys. B 78 (1974) 1 [INSPIRE].

- [7] S. Ferrara and B. Zumino, Supergauge invariant Yang-Mills theories, Nucl. Phys. B 79 (1974)
 413 [INSPIRE].
- [8] A. Salam and J.A. Strathdee, Supersymmetry and non-Abelian gauges, Phys. Lett. B 51 (1974) 353 [INSPIRE].
- [9] J.A. Aguilar-Saavedra, Effects of mixing with quark singlets, Phys. Rev. D 67 (2003) 035003
 [Erratum ibid. 69 (2004) 099901] [hep-ph/0210112] [INSPIRE].
- [10] S. Bejar, J. Guasch and J. Sola, Loop induced flavor changing neutral decays of the top quark in a general two Higgs doublet model, Nucl. Phys. B 600 (2001) 21 [hep-ph/0011091] [INSPIRE].
- [11] J. Guasch and J. Sola, FCNC top quark decays: a door to SUSY physics in high luminosity colliders?, Nucl. Phys. B **562** (1999) 3 [hep-ph/9906268] [INSPIRE].
- [12] J.J. Cao et al., SUSY-induced FCNC top-quark processes at the Large Hadron Collider, Phys. Rev. D 75 (2007) 075021 [hep-ph/0702264] [INSPIRE].
- [13] G. Eilam et al., Top quark rare decay $t \to ch$ in R-parity violating SUSY, Phys. Lett. B **510** (2001) 227 [hep-ph/0102037] [INSPIRE].
- [14] F. del Aguila and M.J. Bowick, The possibility of new fermions with $\Delta I = 0$ mass, Nucl. Phys. B **224** (1983) 107 [INSPIRE].
- [15] J.A. Aguilar-Saavedra, Identifying top partners at LHC, JHEP 11 (2009) 030 [arXiv:0907.3155] [INSPIRE].
- [16] ATLAS collaboration, A detailed map of Higgs boson interactions by the ATLAS experiment ten years after the discovery, Nature 607 (2022) 52 [Erratum ibid. 612 (2022) E24] [arXiv:2207.00092] [INSPIRE].
- [17] CMS collaboration, A portrait of the Higgs boson by the CMS experiment ten years after the discovery, Nature 607 (2022) 60 [arXiv:2207.00043] [INSPIRE].
- [18] LHC Higgs Cross Section Working Group collaboration, Handbook of LHC Higgs cross sections: 4. Deciphering the nature of the Higgs sector, arXiv:1610.07922 [DOI:10.23731/CYRM-2017-002] [INSPIRE].
- [19] ATLAS collaboration, Measurement of the properties of Higgs boson production at $\sqrt{s} = 13 \text{ TeV}$ in the $H \to \gamma \gamma$ channel using 139 fb^{-1} of pp collision data with the ATLAS experiment, arXiv:2207.00348 [INSPIRE].
- [20] ATLAS collaboration, Measurement of the total and differential Higgs boson production cross-sections at $\sqrt{s} = 13$ TeV with the ATLAS detector by combining the $H \to ZZ^* \to 4\ell$ and $H \to \gamma\gamma$ decay channels, JHEP 05 (2023) 028 [arXiv:2207.08615] [INSPIRE].
- [21] ATLAS collaboration, The ATLAS experiment at the CERN Large Hadron Collider, 2008 JINST 3 S08003 [INSPIRE].
- [22] ATLAS collaboration, ATLAS Insertable B-Layer technical design report, CERN-LHCC-2010-013, CERN, Geneva, Switzerland (2010).
- [23] ATLAS IBL collaboration, Production and integration of the ATLAS Insertable B-Layer, 2018 JINST 13 T05008 [arXiv:1803.00844] [INSPIRE].
- [24] ATLAS collaboration, Performance of the ATLAS trigger system in 2015, Eur. Phys. J. C 77 (2017) 317 [arXiv:1611.09661] [INSPIRE].

- [25] ATLAS collaboration, *The ATLAS collaboration software and firmware*, ATL-SOFT-PUB-2021-001, CERN, Geneva, Switzerland (2021).
- [26] ATLAS collaboration, ATLAS data quality operations and performance for 2015–2018 data-taking, 2020 JINST 15 P04003 [arXiv:1911.04632] [INSPIRE].
- [27] ATLAS collaboration, Luminosity determination in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector at the LHC, ATLAS-CONF-2019-021, CERN, Geneva, Switzerland (2019).
- [28] G. Avoni et al., The new LUCID-2 detector for luminosity measurement and monitoring in ATLAS, 2018 JINST 13 P07017 [INSPIRE].
- [29] ATLAS collaboration, Performance of electron and photon triggers in ATLAS during LHC run 2, Eur. Phys. J. C 80 (2020) 47 [arXiv:1909.00761] [INSPIRE].
- [30] K. Hamilton, P. Nason, E. Re and G. Zanderighi, *NNLOPS simulation of Higgs boson production*, *JHEP* **10** (2013) 222 [arXiv:1309.0017] [INSPIRE].
- [31] P. Nason, A new method for combining NLO QCD with shower Monte Carlo algorithms, JHEP 11 (2004) 040 [hep-ph/0409146] [INSPIRE].
- [32] S. Frixione, P. Nason and C. Oleari, Matching NLO QCD computations with parton shower simulations: the POWHEG method, JHEP 11 (2007) 070 [arXiv:0709.2092] [INSPIRE].
- [33] S. Alioli, P. Nason, C. Oleari and E. Re, A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX, JHEP **06** (2010) 043 [arXiv:1002.2581] [INSPIRE].
- [34] P. Nason and C. Oleari, NLO Higgs boson production via vector-boson fusion matched with shower in POWHEG, JHEP 02 (2010) 037 [arXiv:0911.5299] [INSPIRE].
- [35] K. Mimasu, V. Sanz and C. Williams, Higher order QCD predictions for associated Higgs production with anomalous couplings to gauge bosons, JHEP 08 (2016) 039 [arXiv:1512.02572] [INSPIRE].
- [36] J.M. Campbell et al., NLO Higgs boson production plus one and two jets using the POWHEG BOX, MadGraph4 and MCFM, JHEP 07 (2012) 092 [arXiv:1202.5475] [INSPIRE].
- [37] G. Luisoni, P. Nason, C. Oleari and F. Tramontano, $HW^{\pm}/HZ + 0$ and 1 jet at NLO with the POWHEG BOX interfaced to GoSam and their merging within MiNLO, JHEP 10 (2013) 083 [arXiv:1306.2542] [INSPIRE].
- [38] H.B. Hartanto, B. Jäger, L. Reina and D. Wackeroth, Higgs boson production in association with top quarks in the POWHEG BOX, Phys. Rev. D 91 (2015) 094003 [arXiv:1501.04498] [INSPIRE].
- [39] J. Butterworth et al., PDF4LHC recommendations for LHC run II, J. Phys. G 43 (2016) 023001 [arXiv:1510.03865] [INSPIRE].
- [40] T. Sjöstrand, S. Mrenna and P.Z. Skands, A brief introduction to PYTHIA 8.1, Comput. Phys. Commun. 178 (2008) 852 [arXiv:0710.3820] [INSPIRE].
- [41] T. Sjöstrand et al., An introduction to PYTHIA 8.2, Comput. Phys. Commun. 191 (2015) 159 [arXiv:1410.3012] [INSPIRE].
- [42] ATLAS collaboration, Measurement of the Z/γ^* boson transverse momentum distribution in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector, JHEP **09** (2014) 145 [arXiv:1406.3660] [INSPIRE].

- [43] K. Hamilton, P. Nason and G. Zanderighi, MINLO: Multi-scale Improved NLO, JHEP 10 (2012) 155 [arXiv:1206.3572] [INSPIRE].
- [44] K. Hamilton, P. Nason, C. Oleari and G. Zanderighi, $Merging\ H/W/Z + 0$ and 1 jet at NLO with no merging scale: a path to parton shower + NNLO matching, $JHEP\ 05\ (2013)\ 082$ [arXiv:1212.4504] [INSPIRE].
- [45] S. Catani and M. Grazzini, An NNLO subtraction formalism in hadron collisions and its application to Higgs boson production at the LHC, Phys. Rev. Lett. 98 (2007) 222002 [hep-ph/0703012] [INSPIRE].
- [46] G. Bozzi, S. Catani, D. de Florian and M. Grazzini, Transverse-momentum resummation and the spectrum of the Higgs boson at the LHC, Nucl. Phys. B 737 (2006) 73 [hep-ph/0508068] [INSPIRE].
- [47] D. de Florian, G. Ferrera, M. Grazzini and D. Tommasini, Transverse-momentum resummation: Higgs boson production at the Tevatron and the LHC, JHEP 11 (2011) 064 [arXiv:1109.2109] [INSPIRE].
- [48] S. Frixione, P. Nason and G. Ridolfi, A positive-weight next-to-leading-order Monte Carlo for heavy flavour hadroproduction, JHEP 09 (2007) 126 [arXiv:0707.3088] [INSPIRE].
- [49] NNPDF collaboration, Parton distributions for the LHC run II, JHEP **04** (2015) 040 [arXiv:1410.8849] [INSPIRE].
- [50] ATLAS collaboration, ATLAS Pythia 8 tunes to 7 TeV data, ATL-PHYS-PUB-2014-021, CERN, Geneva, Switzerland (2014).
- [51] D.J. Lange, The EvtGen particle decay simulation package, Nucl. Instrum. Meth. A 462 (2001) 152 [INSPIRE].
- [52] Sherpa collaboration, Event generation with Sherpa 2.2, SciPost Phys. 7 (2019) 034 [arXiv:1905.09127] [INSPIRE].
- [53] T. Gleisberg and S. Höche, Comix, a new matrix element generator, JHEP 12 (2008) 039 [arXiv:0808.3674] [INSPIRE].
- [54] OPENLOOPS 2 collaboration, OpenLoops 2, Eur. Phys. J. C 79 (2019) 866 [arXiv:1907.13071] [INSPIRE].
- [55] F. Cascioli, P. Maierhöfer and S. Pozzorini, Scattering amplitudes with Open Loops, Phys. Rev. Lett. 108 (2012) 111601 [arXiv:1111.5206] [INSPIRE].
- [56] A. Denner, S. Dittmaier and L. Hofer, Collier: a fortran-based Complex One-Loop LIbrary in Extended Regularizations, Comput. Phys. Commun. 212 (2017) 220 [arXiv:1604.06792] [INSPIRE].
- [57] S. Schumann and F. Krauss, A parton shower algorithm based on Catani-Seymour dipole factorisation, JHEP 03 (2008) 038 [arXiv:0709.1027] [INSPIRE].
- [58] S. Höche, F. Krauss, M. Schönherr and F. Siegert, A critical appraisal of NLO+PS matching methods, JHEP 09 (2012) 049 [arXiv:1111.1220] [INSPIRE].
- [59] S. Höche, F. Krauss, M. Schonherr and F. Siegert, *QCD matrix elements + parton showers:* the NLO case, JHEP **04** (2013) 027 [arXiv:1207.5030] [INSPIRE].
- [60] S. Catani, F. Krauss, R. Kuhn and B.R. Webber, *QCD matrix elements + parton showers*, *JHEP* 11 (2001) 063 [hep-ph/0109231] [INSPIRE].

- [61] S. Höche, F. Krauss, S. Schumann and F. Siegert, *QCD matrix elements and truncated showers*, *JHEP* **05** (2009) 053 [arXiv:0903.1219] [INSPIRE].
- [62] F. Siegert, A practical guide to event generation for prompt photon production with Sherpa, J. Phys. G 44 (2017) 044007 [arXiv:1611.07226] [INSPIRE].
- [63] S. Frixione, Isolated photons in perturbative QCD, Phys. Lett. B 429 (1998) 369 [hep-ph/9801442] [INSPIRE].
- [64] R.D. Ball et al., Parton distributions with LHC data, Nucl. Phys. B 867 (2013) 244 [arXiv:1207.1303] [INSPIRE].
- [65] N. Arkani-Hamed, A.G. Cohen, E. Katz and A.E. Nelson, *The littlest Higgs*, *JHEP* **07** (2002) 034 [hep-ph/0206021] [INSPIRE].
- [66] K. Agashe, R. Contino and A. Pomarol, *The minimal composite Higgs model*, *Nucl. Phys. B* 719 (2005) 165 [hep-ph/0412089] [INSPIRE].
- [67] L. Randall and R. Sundrum, An alternative to compactification, Phys. Rev. Lett. 83 (1999) 4690 [hep-th/9906064] [INSPIRE].
- [68] J. Alwall et al., The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations, JHEP 07 (2014) 079 [arXiv:1405.0301] [INSPIRE].
- [69] ATLAS collaboration, Search for direct production of electroweakinos in final states with missing transverse momentum and a Higgs boson decaying into photons in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, JHEP 10 (2020) 005 [arXiv:2004.10894] [INSPIRE].
- [70] ATLAS collaboration, Search for trilepton resonances from chargino and neutralino pair production in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector, Phys. Rev. D 103 (2021) 112003 [arXiv:2011.10543] [INSPIRE].
- [71] ATLAS collaboration, Search for top squarks in events with a Higgs or Z boson using $139 \, fb^{-1}$ of pp collision data at $\sqrt{s} = 13 \, TeV$ with the ATLAS detector, Eur. Phys. J. C 80 (2020) 1080 [arXiv:2006.05880] [INSPIRE].
- [72] ATLAS collaboration, Search for bottom-squark pair production with the ATLAS detector in final states containing Higgs bosons, b-jets and missing transverse momentum, JHEP 12 (2019) 060 [arXiv:1908.03122] [INSPIRE].
- [73] ATLAS collaboration, Search for heavy resonances decaying into a photon and a hadronically decaying Higgs boson in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, Phys. Rev. Lett. 125 (2020) 251802 [arXiv:2008.05928] [inSPIRE].
- [74] ATLAS collaboration, Search for top quark decays $t \to qH$, with $H \to \gamma\gamma$, in $\sqrt{s} = 13$ TeV pp collisions using the ATLAS detector, JHEP 10 (2017) 129 [arXiv:1707.01404] [INSPIRE].
- [75] ATLAS collaboration, The Pythia 8 A3 tune description of ATLAS minimum bias and inelastic measurements incorporating the Donnachie-Landshoff diffractive model, ATL-PHYS-PUB-2016-017, CERN, Geneva, Switzerland (2016).
- [76] GEANT4 collaboration, GEANT4 a simulation toolkit, Nucl. Instrum. Meth. A 506 (2003) 250 [INSPIRE].
- [77] ATLAS collaboration, The ATLAS simulation infrastructure, Eur. Phys. J. C 70 (2010) 823 [arXiv:1005.4568] [INSPIRE].

- [78] ATLAS collaboration, Measurement of Higgs boson production in the diphoton decay channel in pp collisions at center-of-mass energies of 7 and 8 TeV with the ATLAS detector, Phys. Rev. D 90 (2014) 112015 [arXiv:1408.7084] [INSPIRE].
- [79] ATLAS collaboration, Electron and photon performance measurements with the ATLAS detector using the 2015–2017 LHC proton-proton collision data, 2019 JINST 14 P12006 [arXiv:1908.00005] [INSPIRE].
- [80] M. Cacciari, G.P. Salam and G. Soyez, The catchment area of jets, JHEP 04 (2008) 005 [arXiv:0802.1188] [INSPIRE].
- [81] M. Cacciari, G.P. Salam and S. Sapeta, On the characterisation of the underlying event, JHEP 04 (2010) 065 [arXiv:0912.4926] [INSPIRE].
- [82] ATLAS collaboration, Measurement of the photon identification efficiencies with the ATLAS detector using LHC run-1 data, Eur. Phys. J. C 76 (2016) 666 [arXiv:1606.01813] [INSPIRE].
- [83] ATLAS collaboration, Measurement of the inclusive isolated prompt photon cross section in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector, Phys. Rev. D 83 (2011) 052005 [arXiv:1012.4389] [INSPIRE].
- [84] ATLAS collaboration, Muon reconstruction performance of the ATLAS detector in proton-proton collision data at $\sqrt{s} = 13$ TeV, Eur. Phys. J. C **76** (2016) 292 [arXiv:1603.05598] [INSPIRE].
- [85] ATLAS collaboration, Jet reconstruction and performance using particle flow with the ATLAS detector, Eur. Phys. J. C 77 (2017) 466 [arXiv:1703.10485] [INSPIRE].
- [86] M. Cacciari, G.P. Salam and G. Soyez, The anti- k_t jet clustering algorithm, JHEP **04** (2008) 063 [arXiv:0802.1189] [INSPIRE].
- [87] M. Cacciari, G.P. Salam and G. Soyez, FastJet user manual, Eur. Phys. J. C 72 (2012) 1896 [arXiv:1111.6097] [INSPIRE].
- [88] ATLAS collaboration, Topological cell clustering in the ATLAS calorimeters and its performance in LHC run 1, Eur. Phys. J. C 77 (2017) 490 [arXiv:1603.02934] [INSPIRE].
- [89] ATLAS collaboration, Performance of pile-up mitigation techniques for jets in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector, Eur. Phys. J. C **76** (2016) 581 [arXiv:1510.03823] [INSPIRE].
- [90] ATLAS collaboration, Forward jet vertex tagging using the particle flow algorithm, ATL-PHYS-PUB-2019-026, CERN, Geneva, Switzerland (2019).
- [91] ATLAS collaboration, ATLAS b-jet identification performance and efficiency measurement with $t\bar{t}$ events in pp collisions at $\sqrt{s}=13$ TeV, Eur. Phys. J. C **79** (2019) 970 [arXiv:1907.05120] [INSPIRE].
- [92] ATLAS collaboration, *Identification of hadronic tau lepton decays using neural networks in the ATLAS experiment*, ATL-PHYS-PUB-2019-033, CERN, Geneva, Switzerland (2019).
- [93] ATLAS collaboration, Performance of missing transverse momentum reconstruction with the ATLAS detector using proton-proton collisions at $\sqrt{s} = 13$ TeV, Eur. Phys. J. C 78 (2018) 903 [arXiv:1802.08168] [INSPIRE].

- [94] ATLAS collaboration, CP properties of Higgs boson interactions with top quarks in the $t\bar{t}H$ and tH processes using $H\to\gamma\gamma$ with the ATLAS detector, Phys. Rev. Lett. 125 (2020) 061802 [arXiv:2004.04545] [INSPIRE].
- [95] ATLAS collaboration, Measurements of the Higgs boson inclusive and differential fiducial cross-sections in the diphoton decay channel with pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, JHEP 08 (2022) 027 [arXiv:2202.00487] [INSPIRE].
- [96] M. Oreglia, A study of the reactions $\psi' \to \gamma \gamma \psi$, Ph.D. thesis, Stanford University, Stanford, CA, U.S.A. (1980) [INSPIRE].
- [97] ATLAS collaboration, Search for scalar diphoton resonances in the mass range 65–600 GeV with the ATLAS detector in pp collision data at $\sqrt{s} = 8$ TeV, Phys. Rev. Lett. 113 (2014) 171801 [arXiv:1407.6583] [INSPIRE].
- [98] ATLAS collaboration, Measurements of Higgs boson properties in the diphoton decay channel with $36 \, fb^{-1}$ of pp collision data at $\sqrt{s} = 13 \, TeV$ with the ATLAS detector, Phys. Rev. D 98 (2018) 052005 [arXiv:1802.04146] [INSPIRE].
- [99] ATLAS and CMS collaborations, Combined measurement of the Higgs boson mass in pp collisions at $\sqrt{s} = 7$ and 8 TeV with the ATLAS and CMS experiments, Phys. Rev. Lett. 114 (2015) 191803 [arXiv:1503.07589] [INSPIRE].
- [100] ATLAS collaboration, Muon reconstruction and identification efficiency in ATLAS using the full run 2 pp collision data set at $\sqrt{s} = 13$ TeV, Eur. Phys. J. C 81 (2021) 578 [arXiv:2012.00578] [INSPIRE].
- [101] ATLAS collaboration, Jet energy scale and resolution measured in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, Eur. Phys. J. C 81 (2021) 689 [arXiv:2007.02645] [INSPIRE].
- [102] ATLAS collaboration, Measurement of the tau lepton reconstruction and identification performance in the ATLAS experiment using pp collisions at $\sqrt{s} = 13 \text{ TeV}$, ATLAS-CONF-2017-029, CERN, Geneva, Switzerland (2017).
- [103] ATLAS collaboration, Measurement of the inelastic proton-proton cross section at $\sqrt{s} = 13$ TeV with the ATLAS detector at the LHC, Phys. Rev. Lett. 117 (2016) 182002 [arXiv:1606.02625] [INSPIRE].
- [104] ATLAS collaboration, Formulae for estimating significance, ATL-PHYS-PUB-2020-025, CERN, Geneva, Switzerland (2020).
- [105] A.L. Read, Presentation of search results: the CL_s technique, J. Phys. G 28 (2002) 2693 [INSPIRE].
- [106] ATLAS collaboration, Evaluating statistical uncertainties and correlations using the bootstrap method, ATL-PHYS-PUB-2021-011, CERN, Geneva, Switzerland (2021).
- [107] ATLAS collaboration, ATLAS computing acknowledgements, ATL-SOFT-PUB-2021-003, CERN, Geneva, Switzerland (2021).

The ATLAS collaboration

```
G. Aad 10, B. Abbott 120, K. Abeling 15, S.H. Abidi 10, A. Aboulhorma 15, S.H. Abidi 10, A. Aboulhorma 15, A. Aboulhorma
H. Abramowicz <sup>151</sup>, H. Abreu <sup>150</sup>, Y. Abulaiti <sup>117</sup>, A.C. Abusleme Hoffman <sup>137a</sup>,
B.S. Acharya 669a,69b,o, C. Adam Bourdarios 64, L. Adamczyk 685a, L. Adamek 155,
S.V. Addepalli <sup>6</sup>26, J. Adelman <sup>6</sup>115, A. Adiguzel <sup>6</sup>21c, S. Adorni <sup>5</sup>6, T. Adye <sup>6</sup>134,
A.A. Affolder <sup>136</sup>, Y. Afik <sup>36</sup>, M.N. Agaras <sup>13</sup>, J. Agarwala <sup>73a,73b</sup>, A. Aggarwal <sup>100</sup>,
C. Agheorghiesei <sup>©</sup> <sup>27c</sup>, J.A. Aguilar-Saavedra <sup>©</sup> <sup>130f</sup>, A. Ahmad <sup>©</sup> <sup>36</sup>, F. Ahmadov <sup>©</sup> <sup>38,y</sup>,
W.S. Ahmed <sup>104</sup>, S. Ahuja <sup>95</sup>, X. Ai <sup>48</sup>, G. Aielli <sup>76a,76b</sup>, M. Ait Tamlihat <sup>35e</sup>,
B. Aitbenchikh ^{\circ} 35a, I. Aizenberg ^{\circ} 4.V. Akbiyik ^{\circ} 7.P.A. Åkesson ^{\circ} 8.V. Akimov ^{\circ} 37,
K. Al Khoury 641, G.L. Alberghi 23b, J. Albert 164, P. Albicocco 53, S. Alderweireldt 52,
M. Aleksa \mathbb{D}^{36}, I.N. Aleksandrov \mathbb{D}^{38}, C. Alexa \mathbb{D}^{27b}, T. Alexopoulos \mathbb{D}^{10}, A. Alfonsi \mathbb{D}^{114},
F. Alfonsi <sup>©</sup> <sup>23b</sup>, M. Alhroob <sup>©</sup> <sup>120</sup>, B. Ali <sup>©</sup> <sup>132</sup>, S. Ali <sup>©</sup> <sup>148</sup>, M. Aliev <sup>©</sup> <sup>37</sup>, G. Alimonti <sup>©</sup> <sup>71a</sup>,
W. Alkakhi <sup>65</sup>, C. Allaire <sup>66</sup>, B.M.M. Allbrooke <sup>6146</sup>, C.A. Allendes Flores <sup>6137</sup>f,
P.P. Allport <sup>©</sup> <sup>20</sup>, A. Aloisio <sup>©</sup> <sup>72a,72b</sup>, F. Alonso <sup>©</sup> <sup>90</sup>, C. Alpigiani <sup>©</sup> <sup>138</sup>, M. Alvarez Estevez <sup>©</sup> <sup>99</sup>,
A. Alvarez Fernandez <sup>100</sup>, M.G. Alviggi <sup>72a,72b</sup>, M. Aly <sup>101</sup>, Y. Amaral Coutinho <sup>82b</sup>,
A. Ambler 104, C. Amelung 6, M. Amerl 101, C.G. Ames 109, D. Amidei 101,
S.P. Amor Dos Santos 130a, K.R. Amos 162, V. Ananiev 125, C. Anastopoulos 139,
T. Andeen <sup>11</sup>, J.K. Anders <sup>36</sup>, S.Y. Andrean <sup>47a,47b</sup>, A. Andreazza <sup>71a,71b</sup>, S. Angelidakis <sup>9</sup>,
A. Angerami <sup>6</sup>41,aa, A.V. Anisenkov <sup>6</sup>37, A. Annovi <sup>6</sup>74a, C. Antel <sup>56</sup>, M.T. Anthony <sup>6</sup>139,
E. Antipov 145, M. Antonelli 53, D.J.A. Antrim 17a, F. Anulli 57a, M. Aoki 83, T. Aoki 15a,
J.A. Aparisi Pozo<sup>162</sup>, M.A. Aparo<sup>146</sup>, L. Aperio Bella<sup>48</sup>, C. Appelt<sup>18</sup>, N. Aranzabal<sup>36</sup>,
V. Araujo Ferraz <sup>682a</sup>, C. Arcangeletti <sup>53</sup>, A.T.H. Arce <sup>51</sup>, E. Arena <sup>92</sup>, J-F. Arguin <sup>108</sup>,
S. Argyropoulos <sup>54</sup>, J.-H. Arling <sup>948</sup>, A.J. Armbruster <sup>936</sup>, O. Arnaez <sup>94</sup>, H. Arnold <sup>9114</sup>,
Z.P. Arrubarrena Tame<sup>109</sup>, G. Artoni <sup>675</sup>a, H. Asada <sup>611</sup>1, K. Asai <sup>6118</sup>, S. Asai <sup>615</sup>3,
N.A. Asbah 661, J. Assahsah 535d, K. Assamagan 529, R. Astalos 528a, R.J. Atkin 533a,
M. Atkinson<sup>161</sup>, N.B. Atlay <sup>18</sup>, H. Atmani<sup>62b</sup>, P.A. Atmasiddha <sup>106</sup>, K. Augsten <sup>132</sup>,
S. Auricchio (10,72a,72b), A.D. Auriol (10,20), V.A. Austrup (10,170), G. Avner (10,150), G. Avolio (10,36),
K. Axiotis <sup>656</sup>, G. Azuelos <sup>6108,ac</sup>, D. Babal <sup>28a</sup>, H. Bachacou <sup>6135</sup>, K. Bachas <sup>152,q</sup>,
A. Bachiu <sup>6</sup> <sup>34</sup>, F. Backman <sup>6</sup> <sup>47a,47b</sup>, A. Badea <sup>61</sup>, P. Bagnaia <sup>675a,75b</sup>, M. Bahmani <sup>618</sup>,
A.J. Bailey <sup>162</sup>, V.R. Bailey <sup>161</sup>, J.T. Baines <sup>134</sup>, C. Bakalis <sup>10</sup>, O.K. Baker <sup>171</sup>,
E. Bakos 15, D. Bakshi Gupta 8, R. Balasubramanian 114, E.M. Baldin 37, P. Balek 133,
E. Ballabene <sup>671</sup>a, 71b, F. Balli <sup>135</sup>, L.M. Baltes <sup>63a</sup>, W.K. Balunas <sup>32</sup>, J. Balz <sup>100</sup>,
E. Banas <sup>686</sup>, M. Bandieramonte <sup>6129</sup>, A. Bandyopadhyay <sup>624</sup>, S. Bansal <sup>24</sup>, L. Barak <sup>5151</sup>,
E.L. Barberio <sup>105</sup>, D. Barberis <sup>57b,57a</sup>, M. Barbero <sup>102</sup>, G. Barbour <sup>96</sup>, K.N. Barends <sup>33a</sup>,
T. Barillari <sup>110</sup>, M-S. Barisits <sup>36</sup>, T. Barklow <sup>143</sup>, P. Baron <sup>122</sup>, D.A. Baron Moreno <sup>101</sup>,
A. Baroncelli 62a, G. Barone 29, A.J. Barr 6126, L. Barranco Navarro 47a,47b, F. Barreiro 99,
J. Barreiro Guimarães da Costa ^{\bullet 14a}, U. Barron ^{\bullet 151}, M.G. Barros Teixeira ^{\bullet 130a}, S. Barsov ^{\bullet 37},
F. Bartels 63a, R. Bartoldus 143, A.E. Barton 991, P. Bartos 28a, A. Basan 100, M. Baselga 49,
I. Bashta 677a,77b, A. Bassalat 666,ah, M.J. Basso 6155, C.R. Basson 1011, R.L. Bates 559,
S. Batlamous<sup>35e</sup>, J.R. Batley <sup>13</sup>, B. Batool <sup>14</sup>, M. Battaglia <sup>13</sup>, D. Battulga <sup>18</sup>,
M. Bauce • <sup>75a,75b</sup>, P. Bauer • <sup>24</sup>, J.B. Beacham • <sup>51</sup>, T. Beau • <sup>127</sup>, P.H. Beauchemin • <sup>158</sup>,
F. Becherer <sup>54</sup>, P. Bechtle <sup>24</sup>, H.P. Beck <sup>19,p</sup>, K. Becker <sup>166</sup>, A.J. Beddall <sup>21d</sup>,
```

```
V.A. Bednyakov <sup>38</sup>, C.P. Bee <sup>145</sup>, L.J. Beemster <sup>15</sup>, T.A. Beermann <sup>36</sup>, M. Begalli <sup>82d,82d</sup>,
M. Begel<sup>29</sup>, A. Behera<sup>145</sup>, J.K. Behr<sup>48</sup>, C. Beirao Da Cruz E Silva<sup>36</sup>, J.F. Beirer<sup>55,36</sup>,
F. Beisiegel\mathbb{D}^{24}, M. Belfkir\mathbb{D}^{116b}, G. Bella\mathbb{D}^{151}, L. Bellagamba\mathbb{D}^{23b}, A. Bellerive\mathbb{D}^{34},
P. Bellos <sup>©20</sup>, K. Beloborodov <sup>©37</sup>, N.L. Belyaev <sup>©37</sup>, D. Benchekroun <sup>©35a</sup>, F. Bendebba <sup>©35a</sup>,
Y. Benhammou<sup>151</sup>, M. Benoit<sup>29</sup>, J.R. Bensinger<sup>26</sup>, S. Bentvelsen<sup>114</sup>, L. Beresford<sup>36</sup>,
M. Beretta <sup>53</sup>, E. Bergeaas Kuutmann <sup>160</sup>, N. Berger <sup>4</sup>, B. Bergmann <sup>132</sup>, J. Beringer <sup>17a</sup>,
S. Berlendis \mathbb{D}^7, G. Bernardi \mathbb{D}^5, C. Bernius \mathbb{D}^{143}, F.U. Bernlochner \mathbb{D}^{24}, T. Berry \mathbb{D}^{95},
P. Berta 6 133, A. Berthold 5 50, I.A. Bertram 6 91, S. Bethke 1 10, A. Betti 7 7 5 a,75 b, A.J. Bevan 6 94,
M. Bhamjee ^{\odot 33c}, S. Bhatta ^{\odot 145}, D.S. Bhattacharya ^{\odot 165}, P. Bhattarai ^{\odot 26}, V.S. Bhopatkar ^{\odot 121},
R. Bi<sup>29,af</sup>, R.M. Bianchi<sup>129</sup>, O. Biebel<sup>109</sup>, R. Bielski<sup>123</sup>, M. Biglietti<sup>77a</sup>,
T.R.V. Billoud 0^{132}, M. Bindi 0^{55}, A. Bingul 0^{21b}, C. Bini 0^{75a,75b}, A. Biondini 0^{92},
C.J. Birch-sykes • 101, G.A. Bird • 20,134, M. Birman • 168, M. Biros • 133, T. Bisanz • 36,
E. Bisceglie <sup>6</sup>43b,43a, D. Biswas <sup>6</sup>169, A. Bitadze <sup>6</sup>101, K. Bjørke <sup>6</sup>125, I. Bloch <sup>6</sup>48, C. Blocker <sup>6</sup>26,
A. Blue <sup>59</sup>, U. Blumenschein <sup>94</sup>, J. Blumenthal <sup>100</sup>, G.J. Bobbink <sup>114</sup>, V.S. Bobrovnikov <sup>37</sup>,
M. Boehler <sup>54</sup>, D. Bogavac <sup>36</sup>, A.G. Bogdanchikov <sup>37</sup>, C. Bohm <sup>47a</sup>, V. Boisvert <sup>95</sup>,
P. Bokan 648, T. Bold 685a, M. Bomben 5, M. Bona 94, M. Boonekamp 135, C.D. Booth 95,
A.G. Borbély <sup>59</sup>, H.M. Borecka-Bielska <sup>108</sup>, L.S. Borgna <sup>96</sup>, G. Borissov <sup>91</sup>, D. Bortoletto <sup>126</sup>,
D. Boscherini <sup>©</sup> <sup>23b</sup>, M. Bosman <sup>©</sup> <sup>13</sup>, J.D. Bossio Sola <sup>©</sup> <sup>36</sup>, K. Bouaouda <sup>©</sup> <sup>35a</sup>, N. Bouchhar <sup>©</sup> <sup>162</sup>,
J. Boudreau <sup>129</sup>, E.V. Bouhova-Thacker <sup>91</sup>, D. Boumediene <sup>40</sup>, R. Bouquet <sup>5</sup>, A. Boveia <sup>119</sup>,
J. Boyd 636, D. Boye 629, I.R. Boyko 538, J. Bracinik 520, N. Brahimi 62d, G. Brandt 170,
O. Brandt 63, F. Braren 48, B. Brau 613, J.E. Brau 612, K. Brendlinger 64, R. Brener 616,
L. Brenner 114, R. Brenner 160, S. Bressler 168, D. Britton 59, D. Britzger 110, I. Brock 24,
G. Brooijmans <sup>14</sup>, W.K. Brooks <sup>137f</sup>, E. Brost <sup>29</sup>, L.M. Brown <sup>164</sup>, T.L. Bruckler <sup>126</sup>,
P.A. Bruckman de Renstrom 686, B. Brüers 648, D. Bruncko 628, A. Bruni 23b, G. Bruni 23b,
M. Bruschi • 23b, N. Bruscino • 75a,75b, T. Buanes • 16, Q. Buat • 138, A.G. Buckley • 59,
I.A. Budagov <sup>38,*</sup>, M.K. Bugge <sup>125</sup>, O. Bulekov <sup>37</sup>, B.A. Bullard <sup>143</sup>, S. Burdin <sup>92</sup>,
C.D. Burgard <sup>6</sup><sup>49</sup>, A.M. Burger <sup>6</sup><sup>40</sup>, B. Burghgrave <sup>6</sup><sup>8</sup>, J.T.P. Burr <sup>6</sup><sup>32</sup>, C.D. Burton <sup>6</sup><sup>11</sup>,
J.C. Burzynski <sup>142</sup>, E.L. Busch <sup>41</sup>, V. Büscher <sup>100</sup>, P.J. Bussey <sup>59</sup>, J.M. Butler <sup>25</sup>,
C.M. Buttar<sup>©</sup><sup>59</sup>, J.M. Butterworth<sup>©</sup><sup>96</sup>, W. Buttinger<sup>©</sup><sup>134</sup>, C.J. Buxo Vazquez<sup>107</sup>,
A.R. Buzykaev <sup>37</sup>, G. Cabras <sup>23b</sup>, S. Cabrera Urbán <sup>162</sup>, D. Caforio <sup>58</sup>, H. Cai <sup>129</sup>,
Y. Cai^{\bullet}<sup>14a,14d</sup>, V.M.M. Cairo^{\bullet}<sup>36</sup>, O. Cakir^{\bullet}<sup>3a</sup>, N. Calace^{\bullet}<sup>36</sup>, P. Calafiura^{\bullet}<sup>17a</sup>,
G. Calderini <sup>127</sup>, P. Calfayan <sup>68</sup>, G. Callea <sup>59</sup>, L.P. Caloba <sup>82b</sup>, D. Calvet <sup>40</sup>, S. Calvet <sup>40</sup>,
T.P. Calvet 10102, M. Calvetti 1074a,74b, R. Camacho Toro 10127, S. Camarda 1036,
D. Camarero Munoz <sup>©</sup> <sup>26</sup>, P. Camarri <sup>©</sup> <sup>76a,76b</sup>, M.T. Camerlingo <sup>©</sup> <sup>72a,72b</sup>, D. Cameron <sup>©</sup> <sup>125</sup>,
C. Camincher <sup>164</sup>, M. Campanelli <sup>96</sup>, A. Camplani <sup>42</sup>, V. Canale <sup>72a,72b</sup>, A. Canesse <sup>104</sup>,
M. Cano Bret 680, J. Cantero 6162, Y. Cao 6161, F. Capocasa 626, M. Capua 43b,43a,
A. Carbone ^{\circ}<sup>71a,71b</sup>, R. Cardarelli ^{\circ}<sup>76a</sup>, J.C.J. Cardenas ^{\circ}<sup>8</sup>, F. Cardillo ^{\circ}<sup>162</sup>, T. Carli ^{\circ}<sup>36</sup>,
G. Carlino <sup>72a</sup>, J.I. Carlotto <sup>13</sup>, B.T. Carlson <sup>129,r</sup>, E.M. Carlson <sup>164,156a</sup>,
L. Carminati <sup>1</sup>0<sup>71a,71b</sup>, M. Carnesale <sup>1</sup>0<sup>75a,75b</sup>, S. Caron <sup>1</sup>113, E. Carquin <sup>1</sup>137f, S. Carrá <sup>1</sup>11a,71b</sup>,
G. Carratta<sup>©</sup><sup>23b,23a</sup>, F. Carrio Argos<sup>©</sup><sup>33g</sup>, J.W.S. Carter<sup>©</sup><sup>155</sup>, T.M. Carter<sup>©</sup><sup>52</sup>,
M.P. Casado <sup>13,i</sup>, A.F. Casha <sup>155</sup>, M. Caspar <sup>48</sup>, E.G. Castiglia <sup>171</sup>, F.L. Castillo <sup>63a</sup>,
L. Castillo Garcia <sup>13</sup>, V. Castillo Gimenez <sup>162</sup>, N.F. Castro <sup>130a,130e</sup>, A. Catinaccio <sup>36</sup>,
J.R. Catmore <sup>125</sup>, V. Cavaliere <sup>29</sup>, N. Cavalli <sup>23b,23a</sup>, V. Cavasinni <sup>74a,74b</sup>, E. Celebi <sup>21a</sup>,
```

F. Celli ¹²⁶, M.S. Centonze ^{70a,70b}, K. Cerny ¹²², A.S. Cerqueira ^{82a}, A. Cerri ¹⁴⁶, L. Cerrito $^{\circ}76a,76b$, F. Cerutti $^{\circ}17a$, A. Cervelli $^{\circ}23b$, G. Cesarini $^{\circ}53$, S.A. Cetin $^{\circ}21d$, Z. Chadi 6^{35a}, D. Chakraborty 6¹¹⁵, M. Chala 6^{130f}, J. Chan 6¹⁶⁹, W.Y. Chan 6¹⁵³, J.D. Chapman ^{©32}, B. Chargeishvili ^{©149b}, D.G. Charlton ^{©20}, T.P. Charman ^{©94}, M. Chatterjee ¹⁹, S. Chekanov ⁶, S.V. Chekulaev ^{156a}, G.A. Chelkov ^{38,a}, A. Chen ¹⁰⁶, B. Chen 6 151, B. Chen 16 164, H. Chen 16 14c, H. Chen 1529, J. Chen 16 16c, J. Chen 1514, S. Chen 153, S.J. Chen 614c, X. Chen 62c, X. Chen 14b, ab, Y. Chen 62a, C.L. Cheng 169, H.C. Cheng 64a, S. Cheong ¹⁴³, A. Cheplakov ³⁸, E. Cheremushkina ⁴⁸, E. Cherepanova ¹¹⁴, R. Cherkaoui El Moursli^{©35e}, E. Cheu^{©7}, K. Cheung^{©65}, L. Chevalier^{©135}, V. Chiarella^{©53}. G. Chiarelli ^{1074a}, N. Chiedde ¹⁰², G. Chiodini ^{1070a}, A.S. Chisholm ¹⁰²⁰, A. Chitan ^{1027b}, M. Chitishvili ¹⁶², M.V. Chizhov ³⁸, K. Choi ¹¹, A.R. Chomont ^{75a,75b}, Y. Chou ¹⁰³, E.Y.S. Chow ¹¹⁴, T. Chowdhury ^{33g}, L.D. Christopher ^{33g}, K.L. Chu^{64a}, M.C. Chu ^{64a}, X. Chu 14a,14d, J. Chudoba 131, J.J. Chwastowski 86, D. Cieri 110, K.M. Ciesla 85a, V. Cindro 693, A. Ciocio 617a, F. Cirotto 72a,72b, Z.H. Citron 168,l, M. Citterio 71a, D.A. Ciubotaru^{27b}, B.M. Ciungu¹⁵⁵, A. Clark⁵⁶, P.J. Clark⁵², J.M. Clavijo Columbie⁴⁸, S.E. Clawson 101, C. Clement 47a,47b, J. Clercx 48, L. Clissa 23b,23a, Y. Coadou 112, M. Cobal 69a,69c, A. Coccaro 57b, R.F. Coelho Barrue 130a, R. Coelho Lopes De Sa 1013, S. Coelli ¹0^{71a}, H. Cohen ¹0¹⁵¹, A.E.C. Coimbra ¹0^{71a,71b}, B. Cole ¹0⁴¹, J. Collot ¹0⁶⁰, P. Conde Muiño 130a,130g, M.P. Connell 33c, S.H. Connell 33c, I.A. Connelly 59, E.I. Conroy ¹²⁶, F. Conventi ^{72a,ad}, H.G. Cooke ²⁰, A.M. Cooper-Sarkar ¹²⁶, F. Cormier ¹⁶³, L.D. Corpe 6 36, M. Corradi 6 75a,75b, F. Corriveau 6 104,w, A. Cortes-Gonzalez 6 18, M.J. Costa 6 162, F. Costanza • 4, D. Costanzo • 139, B.M. Cote • 119, G. Cowan • 55, J.W. Cowley • 32, K. Cranmer ¹¹⁷, S. Crépé-Renaudin ⁶⁰, F. Crescioli ¹²⁷, M. Cristinziani ¹⁴¹, M. Cristoforetti ⁶78a,78b,c, V. Croft ⁶114, G. Crosetti ⁶43b,43a, A. Cueto ⁶36, T. Cuhadar Donszelmann ¹⁵⁹, H. Cui ^{14a,14d}, Z. Cui ⁷, W.R. Cunningham ⁵⁹, F. Curcio 643b,43a, P. Czodrowski 636, M.M. Czurylo 63b, M.J. Da Cunha Sargedas De Sousa 662a, J.V. Da Fonseca Pinto ⁶82b, C. Da Via ⁶101, W. Dabrowski ⁶85a, T. Dado ⁶49, S. Dahbi ⁶33g, T. Dai 6 106, C. Dallapiccola 103, M. Dam 42, G. D'amen 29, V. D'Amico 109, J. Damp 100, J.R. Dandoy 128, M.F. Daneri 30, M. Danninger 142, V. Dao 36, G. Darbo 576, S. Darmora 6, S.J. Das 29, af, S. D'Auria 71a, 71b, C. David 156b, T. Davidek 133, B. Davis-Purcell⁶³⁴, I. Dawson⁶⁹⁴, K. De⁶⁸, R. De Asmundis^{672a}, N. De Biase⁶⁴⁸, S. De Castro 6^{23b,23a}, N. De Groot 1¹¹³, P. de Jong 1¹¹⁴, H. De la Torre 1¹⁰⁷, A. De Maria 1^{14c}, A. De Salvo^{675a}, U. De Sanctis^{676a,76b}, A. De Santo⁶¹⁴⁶, J.B. De Vivie De Regie⁶⁰, D.V. Dedovich³⁸, J. Degens¹¹⁴, A.M. Deiana⁴⁴, F. Del Corso^{23b,23a}, J. Del Peso⁹⁹, F. Del Rio 63a, F. Deliot 135, C.M. Delitzsch 49, M. Della Pietra 72a,72b, D. Della Volpe 56, A. Dell'Acqua ⁶³⁶, L. Dell'Asta ^{671a,71b}, M. Delmastro ⁶⁴, P.A. Delsart ⁶⁰, S. Demers ⁶¹⁷¹, M. Demichev ³⁸, S.P. Denisov ³⁷, L. D'Eramo ¹¹⁵, D. Derendarz ⁸⁶, F. Derue ¹²⁷, P. Dervan 692, K. Desch 624, K. Dette 6155, C. Deutsch 624, F.A. Di Bello 57b,57a, A. Di Ciaccio ^{6,76a,76b}, L. Di Ciaccio ^{6,4}, A. Di Domenico ^{6,75a,75b}, C. Di Donato ^{6,72a,72b}, A. Di Girolamo ³⁶, G. Di Gregorio ⁵, A. Di Luca ^{78a,78b}, B. Di Micco ^{77a,77b}, R. Di Nardo (10,77a,77b), C. Diaconu (10,102), F.A. Dias (11,114), T. Dias Do Vale (11,114), M.A. Diaz (13,7a,137b), F.G. Diaz Capriles ²⁴, M. Didenko ¹⁶², E.B. Diehl ¹⁰⁶, L. Diehl ⁵⁴, S. Díez Cornell ⁴⁸,

C. Diez Pardos 141, C. Dimitriadi 24,160, A. Dimitrievska 17a, J. Dingfelder 24, I-M. Dinu 27b,

```
S.J. Dittmeier <sup>63b</sup>, F. Dittus <sup>36</sup>, F. Djama <sup>102</sup>, T. Djobava <sup>149b</sup>, J.I. Djuvsland <sup>16</sup>,
C. Doglioni 6 101,98, J. Dolejsi 6 133, Z. Dolezal 6 133, M. Donadelli 6 82c, B. Dong 6 107,
J. Donini \mathbb{D}^{40}, A. D'Onofrio \mathbb{D}^{77a,77b}, M. D'Onofrio \mathbb{D}^{92}, J. Dopke \mathbb{D}^{134}, A. Doria \mathbb{D}^{72a},
M.T. Dova • 90, A.T. Doyle • 59, M.A. Draguet • 126, E. Drechsler • 142, E. Dreyer • 168,
I. Drivas-koulouris <sup>10</sup>, A.S. Drobac <sup>158</sup>, M. Drozdova <sup>56</sup>, D. Du <sup>62a</sup>, T.A. du Pree <sup>114</sup>,
F. Dubinin ^{\odot} 37, M. Dubovsky ^{\odot} 28a, E. Duchovni ^{\odot} 168, G. Duckeck ^{\odot} 109, O.A. Ducu ^{\odot} 27b,
D. Duda 110, A. Dudarev 36, E.R. Duden 26, M. D'uffizi 101, L. Duflot 66, M. Dührssen 36,
C. Dülsen 170, A.E. Dumitriu 27b, M. Dunford 63a, S. Dungs 49, K. Dunne 47a,47b,
A. Duperrin <sup>102</sup>, H. Duran Yildiz <sup>3a</sup>, M. Düren <sup>58</sup>, A. Durglishvili <sup>149b</sup>, B.L. Dwyer <sup>15</sup>,
G.I. Dyckes <sup>17a</sup>, M. Dyndal <sup>85a</sup>, S. Dysch <sup>101</sup>, B.S. Dziedzic <sup>86</sup>, Z.O. Earnshaw <sup>146</sup>,
B. Eckerova<sup>©</sup><sup>28a</sup>, S. Eggebrecht<sup>©</sup><sup>55</sup>, M.G. Eggleston<sup>51</sup>, E. Egidio Purcino De Souza<sup>©</sup><sup>127</sup>,
L.F. Ehrke • G. Eigen • K. Einsweiler • T. Ekelof • P.A. Ekman • 8,
Y. El Ghazali <sup>35b</sup>, H. El Jarrari <sup>35e,148</sup>, A. El Moussaouy <sup>35a</sup>, V. Ellajosyula <sup>160</sup>,
M. Ellert <sup>160</sup>, F. Ellinghaus <sup>170</sup>, A.A. Elliot <sup>194</sup>, N. Ellis <sup>36</sup>, J. Elmsheuser <sup>29</sup>, M. Elsing <sup>36</sup>,
D. Emeliyanov 134, Y. Enari 153, I. Ene 17a, S. Epari 13, J. Erdmann 149, P.A. Erland 186,
M. Errenst • <sup>170</sup>, M. Escalier • <sup>66</sup>, C. Escobar • <sup>162</sup>, E. Etzion • <sup>151</sup>, G. Evans • <sup>130a</sup>, H. Evans • <sup>68</sup>,
M.O. Evans <sup>146</sup>, A. Ezhilov <sup>37</sup>, S. Ezzarqtouni <sup>35a</sup>, F. Fabbri <sup>59</sup>, L. Fabbri <sup>23b,23a</sup>,
G. Facini ^{\circ} 96, V. Fadeyev ^{\circ} 136, R.M. Fakhrutdinov ^{\circ} 37, S. Falciano ^{\circ} 75a,
L.F. Falda Ulhoa Coelho 636, P.J. Falke 524, S. Falke 536, J. Faltova 5133, Y. Fan 514a,
Y. Fang 614a,14d, M. Fanti 71a,71b, M. Faraj 669a,69b, Z. Farazpay 7, A. Farbin 8, A. Farilla 77a,
T. Faroque 107, S.M. Farrington 52, F. Fassi 35e, D. Fassouliotis 9,
M. Faucci Giannelli <sup>6</sup>7<sup>6a,76b</sup>, W.J. Fawcett <sup>32</sup>, L. Fayard <sup>66</sup>, P. Federic <sup>133</sup>, P. Federicova <sup>131</sup>,
O.L. Fedin • 37, a, G. Fedotov • 37, M. Feickert • 169, L. Feligioni • 102, A. Fell • 139, D.E. Fellers • 123,
C. Feng 62b, M. Feng 14b, Z. Feng 114, M.J. Fenton 159, A.B. Fenyuk<sup>37</sup>, L. Ferencz 48,
R.A.M. Ferguson <sup>691</sup>, S.I. Fernandez Luengo <sup>6137f</sup>, J. Pretel <sup>54</sup>, J. Ferrando <sup>648</sup>, A. Ferrari <sup>6160</sup>,
P. Ferrari 114,113, R. Ferrari 73a, D. Ferrere 55, C. Ferretti 116, F. Fiedler 110, A. Filipčič 193,
E.K. Filmer <sup>1</sup>, F. Filthaut <sup>113</sup>, M.C.N. Fiolhais <sup>130a,130c,b</sup>, L. Fiorini <sup>162</sup>, F. Fischer <sup>141</sup>,
W.C. Fisher \mathbb{D}^{107}, T. Fitschen \mathbb{D}^{101}, I. Fleck \mathbb{D}^{141}, P. Fleischmann \mathbb{D}^{106}, T. Flick \mathbb{D}^{170},
L. Flores <sup>128</sup>, M. Flores <sup>33d,aj</sup>, L.R. Flores Castillo <sup>64a</sup>, F.M. Follega <sup>78a,78b</sup>, N. Fomin <sup>16</sup>,
J.H. Foo 155, B.C. Forland 8, A. Formica 135, A.C. Forti 101, E. Fortin 101, A.W. Fortman 161,
M.G. Foti^{17a}, L. Fountas^{9,j}, D. Fournier^{66}, H. Fox^{91}, P. Francavilla^{74a,74b},
S. Francescato 61, S. Franchellucci 55, M. Franchini 523b,23a, S. Franchino 53a, D. Francis 54,
L. Franco 113, L. Franconi 19, M. Franklin 161, G. Frattari 126, A.C. Freegard 194,
W.S. Freund ^{\odot 82b}, Y.Y. Frid ^{\odot 151}, N. Fritzsche ^{\odot 50}, A. Froch ^{\odot 54}, D. Froidevaux ^{\odot 36},
J.A. Frost <sup>126</sup>, Y. Fu <sup>62a</sup>, M. Fujimoto <sup>118</sup>, E. Fullana Torregrosa <sup>162,*</sup>, J. Fuster <sup>162</sup>,
A. Gabrielli <sup>6</sup><sup>23b,23a</sup>, A. Gabrielli <sup>6</sup><sup>155</sup>, P. Gadow <sup>6</sup><sup>48</sup>, G. Gagliardi <sup>6</sup><sup>57b,57a</sup>, L.G. Gagnon <sup>6</sup><sup>17a</sup>,
G.E. Gallardo <sup>126</sup>, E.J. Gallas <sup>126</sup>, B.J. Gallop <sup>134</sup>, R. Gamboa Goni <sup>94</sup>, K.K. Gan <sup>119</sup>,
S. Ganguly <sup>153</sup>, J. Gao <sup>62a</sup>, Y. Gao <sup>52</sup>, F.M. Garay Walls <sup>137a,137b</sup>, B. Garcia <sup>29,af</sup>,
C. García 162, J.E. García Navarro 162, M. Garcia-Sciveres 17a, R.W. Gardner 1639,
D. Garg <sup>680</sup>, R.B. Garg <sup>6143,am</sup>, C.A. Garner <sup>155</sup>, S.J. Gasiorowski <sup>6138</sup>, P. Gaspar <sup>682b</sup>,
G. Gaudio <sup>673a</sup>, V. Gautam<sup>13</sup>, P. Gauzzi <sup>675a,75b</sup>, I.L. Gavrilenko <sup>637</sup>, A. Gavrilyuk <sup>637</sup>,
C. Gay 163, G. Gaycken 48, E.N. Gazis 10, A.A. Geanta 1627b,27e, C.M. Gee 136,
C. Gemme <sup>57b</sup>, M.H. Genest <sup>60</sup>, S. Gentile <sup>75a,75b</sup>, S. George <sup>95</sup>, W.F. George <sup>20</sup>,
```

```
T. Geralis <sup>6</sup>46, L.O. Gerlach<sup>55</sup>, P. Gessinger-Befurt <sup>6</sup>36, M.E. Geyik <sup>6</sup>170, M. Ghneimat <sup>6</sup>141,
K. Ghorbanian <sup>694</sup>, A. Ghosal <sup>6141</sup>, A. Ghosh <sup>6159</sup>, A. Ghosh <sup>67</sup>, B. Giacobbe <sup>623b</sup>,
S. Giagu^{075a,75b}, P. Giannetti^{074a}, A. Giannini^{062a}, S.M. Gibson^{095}, M. Gignac^{0136},
D.T. Gil 6856, A.K. Gilbert 685a, B.J. Gilbert 641, D. Gilberg 634, G. Gilles 6114,
N.E.K. Gillwald 648, L. Ginabat 6127, D.M. Gingrich 62, ac, M.P. Giordani 669a, 69c,
P.F. Giraud \mathbb{D}^{135}, G. Giugliarelli \mathbb{D}^{69a,69c}, D. Giugni \mathbb{D}^{71a}, F. Giuli \mathbb{D}^{36}, I. Gkialas \mathbb{D}^{9,j},
L.K. Gladilin \mathbb{D}^{37}, C. Glasman \mathbb{D}^{99}, G.R. Gledhill \mathbb{D}^{123}, M. Glisic \mathbb{D}^{123}, I. Gnesi \mathbb{D}^{43b,f}, Y. Go \mathbb{D}^{29,af},
M. Goblirsch-Kolb<sup>26</sup>, B. Gocke<sup>49</sup>, D. Godin<sup>108</sup>, B. Gokturk<sup>21a</sup>, S. Goldfarb<sup>105</sup>,
T. Golling ^{\bullet 56}, M.G.D. Gololo^{33g}, D. Golubkov ^{\bullet 37}, J.P. Gombas ^{\bullet 107}, A. Gomes ^{\bullet 130a,130b},
G. Gomes Da Silva 141, A.J. Gomez Delegido 162, R. Gonçalo 130a,130c, G. Gonella 123,
L. Gonella 620, A. Gongadze 38, F. Gonnella 20, J.L. Gonski 41, S. González de la Hoz 162,
S. Gonzalez Fernandez <sup>13</sup>, R. Gonzalez Lopez <sup>92</sup>, C. Gonzalez Renteria <sup>17a</sup>,
R. Gonzalez Suarez <sup>160</sup>, S. Gonzalez-Sevilla <sup>56</sup>, G.R. Gonzalvo Rodriguez <sup>162</sup>,
R.Y. González Andana <sup>52</sup>, L. Goossens <sup>36</sup>, P.A. Gorbounov <sup>37</sup>, B. Gorini <sup>36</sup>,
E. Gorini <sup>6</sup>70a,70b, A. Gorišek <sup>6</sup>93, A.T. Goshaw <sup>6</sup>51, M.I. Gostkin <sup>6</sup>38, S. Goswami <sup>6</sup>121,
C.A. Gottardo 636, M. Gouighri 535, V. Goumarre 48, A.G. Goussiou 5138, N. Govender 53c,
I. Grabowska-Bold ^{6}8^{5a}, K. Graham ^{6}3^{4}, E. Gramstad ^{6}1^{25}, S. Grancagnolo ^{6}1^{8}, M. Grandi ^{6}1^{46},
V. Gratchev<sup>37,*</sup>, P.M. Gravila<sup>©</sup><sup>27f</sup>, F.G. Gravili<sup>©</sup><sup>70a,70b</sup>, H.M. Grav<sup>©</sup><sup>17a</sup>, M. Greco<sup>©</sup><sup>70a,70b</sup>,
C. Grefe 24, I.M. Gregor 48, P. Grenier 143, C. Grieco 13, A.A. Grillo 136, K. Grimm 131,m,
S. Grinstein ^{\bullet 13,t}, J.-F. Grivaz ^{\bullet 66}, E. Gross ^{\bullet 168}, J. Grosse-Knetter ^{\bullet 55}, C. Grud ^{106},
J.C. Grundy <sup>126</sup>, L. Guan <sup>106</sup>, W. Guan <sup>169</sup>, C. Gubbels <sup>163</sup>, J.G.R. Guerrero Rojas <sup>162</sup>,
G. Guerrieri 69a,69b, F. Guescini 110, R. Gugel 110, J.A.M. Guhit 110, A. Guida 48,
T. Guillemin <sup>6</sup> 4, E. Guilloton <sup>6</sup> <sup>166,134</sup>, S. Guindon <sup>636</sup>, F. Guo <sup>614a,14d</sup>, J. Guo <sup>62c</sup>, L. Guo <sup>66</sup>,
Y. Guo<sup>106</sup>, R. Gupta<sup>48</sup>, S. Gurbuz<sup>24</sup>, S.S. Gurdasani<sup>54</sup>, G. Gustavino<sup>36</sup>, M. Guth<sup>56</sup>,
P. Gutierrez D<sup>120</sup>, L.F. Gutierrez Zagazeta D<sup>128</sup>, C. Gutschow D<sup>96</sup>, C. Gwenlan D<sup>126</sup>,
C.B. Gwilliam • 22, E.S. Haaland • 125, A. Haas • 117, M. Habedank • 48, C. Haber • 17a,
H.K. Hadavand <sup>68</sup>, A. Hadef <sup>6100</sup>, S. Hadzic <sup>6110</sup>, E.H. Haines <sup>696</sup>, M. Haleem <sup>6165</sup>, J. Haley <sup>6121</sup>,
J.J. Hall^{0139}, G.D. Hallewell^{0102}, L. Halser^{019}, K. Hamano^{0164}, H. Hamdaoui^{035e},
M. Hamer ^{\odot 24}, G.N. Hamity ^{\odot 52}, E.J. Hampshire ^{\odot 95}, J. Han ^{\odot 62b}, K. Han ^{\odot 62a}, L. Han ^{\odot 14c},
L. Han 62a, S. Han 17a, Y.F. Han 155, K. Hanagaki 183, M. Hance 136, D.A. Hangal 141, aa,
H. Hanif • 142, M.D. Hank • 128, R. Hankache • 101, J.B. Hansen • 42, J.D. Hansen • 42,
P.H. Hansen (10<sup>42</sup>, K. Hara (10<sup>157</sup>, D. Harada (10<sup>56</sup>, T. Harenberg (10<sup>170</sup>, S. Harkusha (10<sup>37</sup>,
Y.T. Harris <sup>126</sup>, N.M. Harrison <sup>119</sup>, P.F. Harrison <sup>166</sup>, N.M. Hartman <sup>143</sup>, N.M. Hartmann <sup>109</sup>,
Y. Hasegawa <sup>140</sup>, A. Hasib <sup>52</sup>, S. Haug <sup>19</sup>, R. Hauser <sup>107</sup>, M. Havranek <sup>132</sup>,
C.M. Hawkes ^{\circ} R.J. Hawkings ^{\circ} S. Hayashida ^{\circ} 111, D. Hayden ^{\circ} C. Hayes ^{\circ} C.
R.L. Hayes 114, C.P. Hays 126, J.M. Hays 194, H.S. Hayward 192, F. He 162a, Y. He 154,
Y. He<sup>127</sup>, N.B. Heatley<sup>94</sup>, V. Hedberg<sup>98</sup>, A.L. Heggelund<sup>125</sup>, N.D. Hehir<sup>94</sup>,
C. Heidegger <sup>6</sup> <sup>54</sup>, K.K. Heidegger <sup>6</sup> <sup>54</sup>, W.D. Heidorn <sup>6</sup> <sup>81</sup>, J. Heilman <sup>6</sup> <sup>34</sup>, S. Heim <sup>6</sup> <sup>48</sup>,
T. Heim <sup>17a</sup>, J.G. Heinlein <sup>12a</sup>, J.J. Heinrich <sup>12a</sup>, L. Heinrich <sup>11a</sup>, J. Hejbal <sup>13a</sup>,
L. Helary <sup>148</sup>, A. Held <sup>169</sup>, S. Hellesund <sup>125</sup>, C.M. Helling <sup>163</sup>, S. Hellman <sup>147a,47b</sup>,
C. Helsens 6<sup>36</sup>, R.C.W. Henderson 1, L. Henkelmann 5<sup>32</sup>, A.M. Henriques Correia 6, H. Herde 5<sup>98</sup>,
Y. Hernández Jiménez <sup>145</sup>, L.M. Herrmann <sup>24</sup>, T. Herrmann <sup>50</sup>, G. Herten <sup>54</sup>,
R. Hertenberger <sup>109</sup>, L. Hervas <sup>36</sup>, N.P. Hessey <sup>156a</sup>, H. Hibi <sup>84</sup>, S.J. Hillier <sup>20</sup>,
```

```
F. Hinterkeuser <sup>©24</sup>, M. Hirose <sup>©124</sup>, S. Hirose <sup>©157</sup>, D. Hirschbuehl <sup>©170</sup>, T.G. Hitchings <sup>©101</sup>,
B. Hiti <sup>1</sup>
<sup>93</sup>, J. Hobbs <sup>145</sup>, R. Hobincu <sup>27e</sup>, N. Hod <sup>168</sup>, M.C. Hodgkinson <sup>139</sup>,
B.H. Hodkinson \mathbb{D}^{32}, A. Hoecker \mathbb{D}^{36}, J. Hofer \mathbb{D}^{48}, T. Holm \mathbb{D}^{24}, M. Holzbock \mathbb{D}^{110},
L.B.A.H. Hommels • 32, B.P. Honan • 101, J. Hong • 62c, T.M. Hong • 129, J.C. Honig • 54,
B.H. Hooberman (161), W.H. Hopkins (16), Y. Horii (111), S. Hou (148), A.S. Howard (193),
J. Howarth 59, J. Hoya 66, M. Hrabovsky 122, A. Hrynevich 48, T. Hryn'ova 4, P.J. Hsu 65,
S.-C. Hsu 138, Q. Hu 141, Y.F. Hu 14a,14d,ae, D.P. Huang 196, S. Huang 164b, X. Huang 164c,
Y. Huang 62a, Y. Huang 14a, Z. Huang 101, Z. Hubacek 132, M. Huebner 24, F. Huegging 24,
T.B. Huffman <sup>126</sup>, M. Huhtinen <sup>36</sup>, S.K. Huiberts <sup>16</sup>, R. Hulsken <sup>104</sup>, N. Huseynov <sup>12,a</sup>,
J. Huston 10<sup>107</sup>, J. Huth 16<sup>1</sup>, R. Hyneman 14<sup>13</sup>, G. Iacobucci 15<sup>16</sup>, G. Iakovidis 16<sup>29</sup>,
I. Ibragimov 141, L. Iconomidou-Fayard 66, P. Iengo 72a,72b, R. Iguchi 153, T. Iizawa 56,
Y. Ikegami \mathbb{D}^{83}, A. Ilg \mathbb{D}^{19}, N. Ilic \mathbb{D}^{155}, H. Imam \mathbb{D}^{35a}, T. Ingebretsen Carlson \mathbb{D}^{47a,47b},
G. Introzzi • <sup>73a,73b</sup>, M. Iodice • <sup>77a</sup>, V. Ippolito • <sup>75a,75b</sup>, M. Ishino • <sup>153</sup>, W. Islam • <sup>169</sup>,
C. Issever • 18,48, S. Istin • 21a,al, H. Ito • 167, J.M. Iturbe Ponce • 64a, R. Iuppa • 78a,78b,
A. Ivina 6168, J.M. Izen 45, V. Izzo 72a, P. Jacka 131,132, P. Jackson 1, R.M. Jacobs 48,
B.P. Jaeger <sup>142</sup>, C.S. Jagfeld <sup>109</sup>, P. Jain <sup>54</sup>, G. Jäkel <sup>170</sup>, K. Jakobs <sup>54</sup>, T. Jakoubek <sup>168</sup>,
J. Jamieson <sup>59</sup>, K.W. Janas <sup>85a</sup>, A.E. Jaspan <sup>92</sup>, M. Javurkova <sup>103</sup>, F. Jeanneau <sup>135</sup>,
L. Jeanty ^{\bullet} 123, J. Jejelava ^{\bullet} 149a,z, P. Jenni ^{\bullet} 54,g, C.E. Jessiman ^{\bullet} 34, S. Jézéquel ^{\bullet} 4, C. Jia 62b,
J. Jia 6145, X. Jia 61, X. Jia 14a,14d, Z. Jia 14c, Y. Jiang 2a, S. Jiggins 48,
J. Jimenez Pena^{\bullet 110}, S. Jin^{\bullet 14c}, A. Jinaru^{\bullet 27b}, O. Jinnouchi^{\bullet 154}, P. Johansson^{\bullet 139},
K.A. Johns<sup>17</sup>, J.W. Johnson<sup>136</sup>, D.M. Jones<sup>32</sup>, E. Jones<sup>166</sup>, P. Jones<sup>32</sup>, R.W.L. Jones<sup>91</sup>,
T.J. Jones ^{\odot 92}, R. Joshi ^{\odot 119}, J. Jovicevic ^{\odot 15}, X. Ju^{\odot 17a}, J.J. Junggeburth ^{\odot 36},
T. Junkermann 63a, A. Juste Rozas 13t, S. Kabana 137e, A. Kaczmarska 86, M. Kado 110,
H. Kagan 119, M. Kagan 143, A. Kahn 41, A. Kahn 128, C. Kahra 100, T. Kaji 167,
E. Kajomovitz <sup>150</sup>, N. Kakati <sup>168</sup>, C.W. Kalderon <sup>29</sup>, A. Kamenshchikov <sup>155</sup>,
S. Kanayama <sup>154</sup>, N.J. Kang <sup>136</sup>, D. Kar <sup>33g</sup>, K. Karava <sup>126</sup>, M.J. Kareem <sup>156b</sup>,
E. Karentzos <sup>654</sup>, I. Karkanias <sup>6152,e</sup>, S.N. Karpov <sup>638</sup>, Z.M. Karpova <sup>638</sup>, V. Kartvelishvili <sup>691</sup>,
A.N. Karyukhin 6<sup>37</sup>, E. Kasimi 6<sup>152,e</sup>, J. Katzy 6<sup>48</sup>, S. Kaur 6<sup>34</sup>, K. Kawade 6<sup>140</sup>,
T. Kawamoto 135, G. Kawamura 55, E.F. Kay 164, F.I. Kaya 158, S. Kazakos 13,
V.F. Kazanin 637, Y. Ke 6145, J.M. Keaveney 633a, R. Keeler 6164, G.V. Kehris 661,
J.S. Keller <sup>634</sup>, A.S. Kelly <sup>96</sup>, D. Kelsey <sup>6146</sup>, J.J. Kempster <sup>6146</sup>, K.E. Kennedy <sup>641</sup>,
P.D. Kennedy <sup>6</sup>100, O. Kepka <sup>6</sup>131, B.P. Kerridge <sup>6</sup>166, S. Kersten <sup>6</sup>170, B.P. Kerševan <sup>6</sup>93,
S. Keshri 666, L. Keszeghova 628a, S. Ketabchi Haghighat 6155, M. Khandoga 6127,
A. Khanov <sup>121</sup>, A.G. Kharlamov <sup>37</sup>, T. Kharlamova <sup>37</sup>, E.E. Khoda <sup>138</sup>, T.J. Khoo <sup>18</sup>,
G. Khoriauli <sup>165</sup>, J. Khubua <sup>149b</sup>, Y.A.R. Khwaira <sup>66</sup>, M. Kiehn <sup>36</sup>, A. Kilgallon <sup>123</sup>,
D.W. Kim • 47a,47b, E. Kim • 154, Y.K. Kim • 39, N. Kimura • 96, A. Kirchhoff • 55, C. Kirfel • 24,
J. Kirk 134, A.E. Kiryunin 110, T. Kishimoto 153, D.P. Kisliuk 55, C. Kitsaki 10,
O. Kivernyk • 24, M. Klassen • 63a, C. Klein • 34, L. Klein • 165, M.H. Klein • 106, M. Klein • 92,
S.B. Klein <sup>6</sup> <sup>56</sup>, U. Klein <sup>692</sup>, P. Klimek <sup>636</sup>, A. Klimentov <sup>629</sup>, F. Klimpel <sup>6110</sup>,
T. Klioutchnikova <sup>6</sup> <sup>36</sup>, P. Kluit <sup>6</sup> <sup>114</sup>, S. Kluth <sup>6</sup> <sup>110</sup>, E. Kneringer <sup>6</sup> <sup>79</sup>, T.M. Knight <sup>6</sup> <sup>155</sup>,
A. Knue<sup>654</sup>, R. Kobayashi<sup>87</sup>, M. Kocian<sup>6143</sup>, P. Kodyš<sup>6133</sup>, D.M. Koeck<sup>6146</sup>, P.T. Koenig<sup>624</sup>,
T. Koffas • 34, M. Kolb • 135, I. Koletsou • 4, T. Komarek • 122, K. Köneke • 4, A.X.Y. Kong • 1,
T. Kono <sup>118</sup>, N. Konstantinidis <sup>96</sup>, B. Konya <sup>98</sup>, R. Kopeliansky <sup>68</sup>, S. Koperny <sup>85a</sup>,
```

```
K. Korcyl 686, K. Kordas 6152,e, G. Koren 6151, A. Korn 696, S. Korn 655, I. Korolkov 613,
N. Korotkova <sup>©37</sup>, B. Kortman <sup>©114</sup>, O. Kortner <sup>©110</sup>, S. Kortner <sup>©110</sup>, W.H. Kostecka <sup>©115</sup>,
V.V. Kostyukhin <sup>141</sup>, A. Kotsokechagia <sup>135</sup>, A. Kotwal <sup>51</sup>, A. Koulouris <sup>36</sup>,
A. Kourkoumeli-Charalampidi <sup>673a,73b</sup>, C. Kourkoumelis <sup>69</sup>, E. Kourlitis <sup>6</sup>, O. Kovanda <sup>146</sup>,
R. Kowalewski <sup>164</sup>, W. Kozanecki <sup>135</sup>, A.S. Kozhin <sup>37</sup>, V.A. Kramarenko <sup>37</sup>,
G. Kramberger <sup>693</sup>, P. Kramer <sup>100</sup>, M.W. Krasny <sup>127</sup>, A. Krasznahorkay <sup>36</sup>, J.A. Kremer <sup>100</sup>,
T. Kresse <sup>150</sup>, J. Kretzschmar <sup>192</sup>, K. Kreul <sup>18</sup>, P. Krieger <sup>155</sup>, S. Krishnamurthy <sup>103</sup>,
M. Krivos 6 133, K. Krizka 6 20, K. Kroeninger 6 49, H. Kroha 6 110, J. Kroll 6 131, J. Kroll 6 128,
K.S. Krowpman <sup>107</sup>, U. Kruchonak <sup>38</sup>, H. Krüger <sup>24</sup>, N. Krumnack M.C. Kruse <sup>51</sup>,
J.A. Krzysiak <sup>686</sup>, O. Kuchinskaia <sup>637</sup>, S. Kuday <sup>3a</sup>, S. Kuehn <sup>636</sup>, R. Kuesters <sup>54</sup>, T. Kuhl <sup>48</sup>,
V. Kukhtin 638, Y. Kulchitsky 537,a, S. Kuleshov 5137d,137b, M. Kumar 533g, N. Kumari 5102,
A. Kupco<sup>131</sup>, T. Kupfer<sup>49</sup>, A. Kupich<sup>37</sup>, O. Kuprash<sup>54</sup>, H. Kurashige<sup>84</sup>,
L.L. Kurchaninov ^{\bullet} 156a, Y.A. Kurochkin ^{\bullet} 37, A. Kurova ^{\bullet} 37, M. Kuze ^{\bullet} 4.K. Kvam ^{\bullet} 103,
J. Kvita <sup>122</sup>, T. Kwan <sup>104</sup>, N.G. Kyriacou <sup>106</sup>, L.A.O. Laatu <sup>102</sup>, C. Lacasta <sup>162</sup>,
F. Lacava • <sup>75a,75b</sup>, H. Lacker • <sup>18</sup>, D. Lacour • <sup>127</sup>, N.N. Lad • <sup>96</sup>, E. Ladygin • <sup>38</sup>, B. Laforge • <sup>127</sup>,
T. Lagouri <sup>137e</sup>, S. Lai <sup>55</sup>, I.K. Lakomiec <sup>85a</sup>, N. Lalloue <sup>60</sup>, J.E. Lambert <sup>120</sup>,
S. Lammers 668, W. Lampl 7, C. Lampoudis 152,e, A.N. Lancaster 115, E. Lançon 29,
U. Landgraf <sup>654</sup>, M.P.J. Landon <sup>694</sup>, V.S. Lang <sup>654</sup>, R.J. Langenberg <sup>6103</sup>, A.J. Lankford <sup>6159</sup>,
F. Lanni <sup>36</sup>, K. Lantzsch <sup>24</sup>, A. Lanza <sup>73a</sup>, A. Lapertosa <sup>57b,57a</sup>, J.F. Laporte <sup>135</sup>,
T. Lari <sup>1</sup>0<sup>71a</sup>, F. Lasagni Manghi <sup>23b</sup>, M. Lassnig <sup>36</sup>, V. Latonova <sup>131</sup>, A. Laudrain <sup>100</sup>,
A. Laurier <sup>150</sup>, S.D. Lawlor <sup>95</sup>, Z. Lawrence <sup>101</sup>, M. Lazzaroni <sup>71a,71b</sup>, B. Le<sup>101</sup>,
E.M. Le Boulicaut <sup>51</sup>, B. Leban <sup>93</sup>, A. Lebedev <sup>81</sup>, M. LeBlanc <sup>36</sup>, F. Ledroit-Guillon <sup>60</sup>,
A.C.A. Lee^{96}, G.R. Lee^{16}, S.C. Lee^{148}, S. Lee^{47a,47b}, T.F. Lee^{92}, L.L. Leeuw^{33c},
H.P. Lefebvre <sup>95</sup>, M. Lefebvre <sup>164</sup>, C. Leggett <sup>17a</sup>, K. Lehmann <sup>142</sup>, G. Lehmann Miotto <sup>36</sup>,
M. Leigh • 56, W.A. Leight • 103, A. Leisos • 152, s, M.A.L. Leite • 82c, C.E. Leitgeb • 48,
R. Leitner <sup>133</sup>, K.J.C. Leney <sup>44</sup>, T. Lenz <sup>24</sup>, S. Leone <sup>74a</sup>, C. Leonidopoulos <sup>52</sup>,
A. Leopold <sup>144</sup>, C. Leroy <sup>108</sup>, R. Les <sup>107</sup>, C.G. Lester <sup>132</sup>, M. Levchenko <sup>137</sup>, J. Levêque <sup>14</sup>,
D. Levin 106, L.J. Levinson 168, M.P. Lewicki 186, D.J. Lewis 14, A. Li 15, B. Li 162, C. Li 162a,
C-Q. Li 62c, H. Li 62a, H. Li 62b, H. Li 62c, H. Li 62c, H. Li 62c, K. Li 62c, K. Li 62c,
M. Li 614a,14d, Q.Y. Li 62a, S. Li 614a,14d, S. Li 62d,62c,d, T. Li 62b, X. Li 6104, Z. Li 62b,
Z. Li<sup>1</sup>0<sup>126</sup>, Z. Li<sup>1</sup>0<sup>104</sup>, Z. Li<sup>1</sup>0<sup>92</sup>, Z. Li<sup>1</sup>0<sup>14a,14d</sup>, Z. Liang<sup>1</sup>0<sup>14a</sup>, M. Liberatore<sup>1</sup>0<sup>48</sup>, B. Liberti<sup>1</sup>0<sup>76a</sup>,
K. Lie<sup>64c</sup>, J. Lieber Marin<sup>82b</sup>, H. Lien<sup>68</sup>, K. Lin<sup>6107</sup>, R.A. Linck<sup>68</sup>, R.E. Lindley<sup>7</sup>,
J.H. Lindon <sup>©</sup><sup>2</sup>, A. Linss <sup>©</sup><sup>48</sup>, E. Lipeles <sup>©</sup><sup>128</sup>, A. Lipniacka <sup>©</sup><sup>16</sup>, A. Lister <sup>©</sup><sup>163</sup>, J.D. Little <sup>©</sup><sup>4</sup>,
B. \text{Liu}^{\bullet 14a}, B.X. \text{Liu}^{\bullet 142}, D. \text{Liu}^{\bullet 62d,62c}, J.B. \text{Liu}^{\bullet 62a}, J.K.K. \text{Liu}^{\bullet 32}, K. \text{Liu}^{\bullet 62d,62c},
M. Liu 62a, M.Y. Liu 62a, P. Liu 14a, Q. Liu 62d, 138,62c, X. Liu 62a, Y. Liu 14c,14d.
Y.L. Liu<sup>10106</sup>, Y.W. Liu<sup>1062a</sup>, J. Llorente Merino<sup>1142</sup>, S.L. Lloyd<sup>1094</sup>, E.M. Lobodzinska<sup>148</sup>,
P. Loch <sup>1</sup>0<sup>7</sup>, S. Loffredo <sup>1</sup>6<sup>6</sup>a, T. Lohse <sup>1</sup>8, K. Lohwasser <sup>1</sup>3<sup>9</sup>, E. Loiacono <sup>4</sup>8,
M. Lokajicek 131, J.D. Long 161, I. Longarini 159, L. Longo 170a, 70b, R. Longo 161,
I. Lopez Paz 6, A. Lopez Solis 4, J. Lorenz 10, N. Lorenzo Martinez 4, A.M. Lory 10, N. Lorenzo Martinez 4, A.M. Lory 10, N. Lorenzo Martinez 10, A.M. Lory 10, A.M. Lorenzo Martinez 11, A.M. Lorenz 
X. Lou 647a,47b, X. Lou 14a,14d, A. Lounis 666, J. Love 66, P.A. Love 991, G. Lu 14a,14d,
M. Lu<sup>®80</sup>, S. Lu<sup>®128</sup>, Y.J. Lu<sup>®65</sup>, H.J. Lubatti<sup>®138</sup>, C. Luci<sup>®75a,75b</sup>, F.L. Lucio Alves<sup>®14c</sup>,
A. Lucotte <sup>60</sup>, F. Luehring <sup>68</sup>, I. Luise <sup>145</sup>, O. Lukianchuk <sup>66</sup>, O. Lundberg <sup>144</sup>,
B. Lund-Jensen 144, N.A. Luongo 123, M.S. Lutz 151, D. Lynn 29, H. Lyons 2, R. Lysak 131,
```

```
E. Lytken <sup>698</sup>, V. Lyubushkin <sup>638</sup>, T. Lyubushkina <sup>638</sup>, M.M. Lyukova <sup>6145</sup>, H. Ma <sup>629</sup>,
L.L. Ma<sup>6</sup>62b, Y. Ma<sup>6</sup>96, D.M. Mac Donell<sup>6</sup>164, G. Maccarrone<sup>53</sup>, J.C. MacDonald<sup>139</sup>,
R. Madar • 40, W.F. Mader • 50, J. Maeda • 84, T. Maeno • 29, M. Maerker • 50, H. Maguire • 139,
A. Maio 6130a,130b,130d, K. Maj 85a, O. Majersky 648, S. Majewski 123, N. Makovec 666,
V. Maksimovic <sup>15</sup>, B. Malaescu <sup>127</sup>, Pa. Malecki <sup>86</sup>, V.P. Maleev <sup>37</sup>, F. Malek <sup>60</sup>,
D. Malito 643b,43a, U. Mallik 680, C. Malone 632, S. Maltezos 10, S. Malyukov 38, J. Mamuzic 613,
G. Mancini 0^{53}, G. Manco 0^{73a,73b}, J.P. Mandalia 0^{94}, I. Mandić 0^{93},
L. Manhaes de Andrade Filho <sup>682a</sup>, I.M. Maniatis <sup>6168</sup>, J. Manjarres Ramos <sup>6102,ai</sup>,
D.C. Mankad • 168, A. Mann • 109, B. Mansoulie • 135, S. Manzoni • 36, A. Marantis • 152, s,
G. Marchiori <sup>6</sup>, M. Marcisovsky <sup>6</sup>, C. Marcon <sup>6</sup>, M. Marinescu <sup>6</sup>, M. Marjanovic <sup>6</sup>,
E.J. Marshall <sup>91</sup>, Z. Marshall <sup>17a</sup>, S. Marti-Garcia <sup>162</sup>, T.A. Martin <sup>166</sup>, V.J. Martin <sup>52</sup>,
B. Martin dit Latour <sup>16</sup>, L. Martinelli <sup>75a,75b</sup>, M. Martinez <sup>13,t</sup>, P. Martinez Agullo <sup>162</sup>,
V.I. Martinez Outschoorn <sup>103</sup>, P. Martinez Suarez <sup>13</sup>, S. Martin-Haugh <sup>134</sup>, V.S. Martoiu <sup>27b</sup>,
A.C. Martyniuk <sup>696</sup>, A. Marzin <sup>636</sup>, S.R. Maschek <sup>6110</sup>, D. Mascione <sup>678a,78b</sup>, L. Masetti <sup>6100</sup>,
T. Mashimo <sup>153</sup>, J. Masik <sup>101</sup>, A.L. Maslennikov <sup>37</sup>, L. Massa <sup>23b</sup>, P. Massarotti <sup>72a,72b</sup>,
P. Mastrandrea <sup>1</sup>0<sup>74a,74b</sup>, A. Mastroberardino <sup>1</sup>43b,43a</sup>, T. Masubuchi <sup>1</sup>53, T. Mathisen <sup>1</sup>60,
N. Matsuzawa<sup>153</sup>, J. Maurer <sup>©</sup><sup>27b</sup>, B. Maček <sup>©</sup><sup>93</sup>, D.A. Maximov <sup>©</sup><sup>37</sup>, R. Mazini <sup>©</sup><sup>148</sup>,
I. Maznas • 152,e, M. Mazza • 1017, S.M. Mazza • 136, C. Mc Ginn • 29, J.P. Mc Gowan • 104,
S.P. Mc Kee <sup>106</sup>, E.F. McDonald <sup>105</sup>, A.E. McDougall <sup>114</sup>, J.A. Mcfayden <sup>146</sup>,
G. Mchedlidze \mathbb{D}^{149b}, R.P. Mckenzie \mathbb{D}^{33g}, T.C. Mclachlan \mathbb{D}^{48}, D.J. Mclaughlin \mathbb{D}^{96},
K.D. McLean 16, S.J. McMahon 134, P.C. McNamara 15, C.M. Mcpartland 19,
R.A. McPherson • 164,w, T. Megy • 40, S. Mehlhase • 109, A. Mehta • 92, D. Melini • 150,
B.R. Mellado Garcia <sup>33g</sup>, A.H. Melo <sup>55</sup>, F. Meloni <sup>48</sup>, A.M. Mendes Jacques Da Costa <sup>101</sup>,
H.Y. Meng 6155, L. Meng 691, S. Menke 6110, M. Mentink 636, E. Meoni 6436,43a,
C. Merlassino ^{\bullet 126}, L. Merola ^{\bullet 72a,72b}, C. Meroni ^{\bullet 71a}, G. Merz<sup>106</sup>, O. Meshkov ^{\bullet 37},
J. Metcalfe 6, A.S. Mete 6, C. Meyer 68, J-P. Meyer 135, R.P. Middleton 134, L. Mijović 52,
G. Mikenberg <sup>168</sup>, M. Mikestikova <sup>131</sup>, M. Mikuž <sup>93</sup>, H. Mildner <sup>139</sup>, A. Milic <sup>36</sup>,
C.D. Milke • 44, D.W. Miller • 39, L.S. Miller • 34, A. Milov • 168, D.A. Milstead 47a, 47b, T. Min 14c,
A.A. Minaenko <sup>637</sup>, I.A. Minashvili <sup>6149b</sup>, L. Mince <sup>659</sup>, A.I. Mincer <sup>6117</sup>, B. Mindur <sup>685a</sup>,
M. Mineev • 38, Y. Mino • 87, L.M. Mir • 13, M. Miralles Lopez • 162, M. Mironova • 126,
M.C. Missio ^{\odot 113}, T. Mitani ^{\odot 167}, A. Mitra ^{\odot 166}, V.A. Mitsou ^{\odot 162}, O. Miu ^{\odot 155},
P.S. Miyagawa <sup>694</sup>, Y. Miyazaki<sup>89</sup>, A. Mizukami <sup>683</sup>, T. Mkrtchyan <sup>63a</sup>, M. Mlinarevic <sup>696</sup>,
T. Mlinarevic <sup>696</sup>, M. Mlynarikova <sup>636</sup>, S. Mobius <sup>655</sup>, K. Mochizuki <sup>6108</sup>, P. Moder <sup>648</sup>,
P. Mogg <sup>109</sup>, A.F. Mohammed <sup>14a,14d</sup>, S. Mohapatra <sup>14</sup>, G. Mokgatitswane <sup>33g</sup>,
B. Mondal<sup>141</sup>, S. Mondal<sup>132</sup>, K. Mönig<sup>148</sup>, E. Monnier<sup>102</sup>, L. Monsonis Romero<sup>162</sup>,
J. Montejo Berlingen <sup>683</sup>, M. Montella <sup>6119</sup>, F. Monticelli <sup>690</sup>, N. Morange <sup>66</sup>,
A.L. Moreira De Carvalho <sup>130a</sup>, M. Moreno Llácer <sup>162</sup>, C. Moreno Martinez <sup>56</sup>,
P. Morettini <sup>657b</sup>, S. Morgenstern <sup>6166</sup>, M. Morii <sup>61</sup>, M. Morinaga <sup>6153</sup>, A.K. Morley <sup>636</sup>,
F. Morodei <sup>575</sup>a, T. Morvaj <sup>536</sup>, P. Moschovakos <sup>36</sup>, B. Moser <sup>536</sup>, M. Mosidze <sup>149b</sup>,
T. Moskalets ^{\odot} 54, P. Moskvitina ^{\odot} 113, J. Moss ^{\odot} 31,^n, E.J.W. Moyse ^{\odot} 103, O. Mtintsilana ^{\odot} 33^g,
S. Muanza <sup>10</sup> J. Mueller <sup>129</sup>, D. Muenstermann <sup>191</sup>, R. Müller <sup>19</sup>, G.A. Mullier <sup>160</sup>,
J.J. Mullin<sup>128</sup>, D.P. Mungo <sup>155</sup>, J.L. Munoz Martinez <sup>13</sup>, D. Munoz Perez <sup>162</sup>,
```

F.J. Munoz Sanchez 1010, M. Murin 1010, W.J. Murray 166,134, A. Murrone 171a,71b,

```
J.M. Muse 120, M. Muškinja 17a, C. Mwewa 29, A.G. Myagkov 37,a, A.J. Myers 8,
A.A. Myers<sup>129</sup>, G. Myers<sup>68</sup>, M. Myska<sup>132</sup>, B.P. Nachman<sup>17a</sup>, O. Nackenhorst<sup>49</sup>,
A. Nago 50, K. Nagai 126, K. Nagano 83, J.L. Nagle 29, af, E. Nagy 102, A.M. Nairz 36,
Y. Nakahama <sup>683</sup>, K. Nakamura <sup>683</sup>, H. Nanjo <sup>6124</sup>, R. Narayan <sup>644</sup>, E.A. Narayanan <sup>6112</sup>,
I. Naryshkin \mathbb{D}^{37}, M. Naseri \mathbb{D}^{34}, C. Nass \mathbb{D}^{24}, G. Navarro \mathbb{D}^{22a}, J. Navarro-Gonzalez \mathbb{D}^{162},
R. Nayak 151, A. Nayaz 18, P.Y. Nechaeva 137, F. Nechansky 148, L. Nedic 126, T.J. Neep 120,
A. Negri (5)73a,73b, M. Negrini (5)23b, C. Nellist (5)114, C. Nelson (5)104, K. Nelson (5)106,
S. Nemecek 131, M. Nessi 36,h, M.S. Neubauer 161, F. Neuhaus 1010, J. Neundorf 48,
R. Newhouse 163, P.R. Newman 20, C.W. Ng 129, Y.W.Y. Ng 48, B. Ngair 35e,
H.D.N. Nguyen <sup>108</sup>, R.B. Nickerson <sup>126</sup>, R. Nicolaidou <sup>135</sup>, J. Nielsen <sup>136</sup>, M. Niemeyer <sup>55</sup>,
N. Nikiforou 636, V. Nikolaenko 637,a, I. Nikolic-Audit 6127, K. Nikolopoulos 620, P. Nilsson 629,
I. Ninca 648, H.R. Nindhito 56, G. Ninio 5151, A. Nisati 575a, N. Nishu 2, R. Nisius 110,
J-E. Nitschke<sup>50</sup>, E.K. Nkadimeng<sup>33g</sup>, S.J. Noacco Rosende<sup>90</sup>, T. Nobe<sup>153</sup>, D.L. Noel<sup>32</sup>,
Y. Noguchi <sup>687</sup>, T. Nommensen <sup>6147</sup>, M.A. Nomura <sup>29</sup>, M.B. Norfolk <sup>6139</sup>, R.R.B. Norisam <sup>696</sup>,
B.J. Norman 634, J. Novak 693, T. Novak 648, L. Novotny 6132, R. Novotny 6112, L. Nozka 6122,
K. Ntekas <sup>159</sup>, N.M.J. Nunes De Moura Junior <sup>82b</sup>, E. Nurse<sup>96</sup>, J. Ocariz <sup>127</sup>, A. Ochi <sup>84</sup>,
I. Ochoa^{\bullet 130a}, S. Oerdek^{\bullet 160}, J.T. Offermann^{\bullet 39}, A. Ogrodnik^{\bullet 85a}, A. Oh^{\bullet 101},
C.C. Ohm 6144, H. Oide 83, R. Oishi 153, M.L. Ojeda 48, Y. Okazaki 87, M.W. O'Keefe92,
Y. Okumura <sup>153</sup>, L.F. Oleiro Seabra <sup>130a</sup>, S.A. Olivares Pino <sup>137d</sup>, D. Oliveira Damazio <sup>29</sup>,
D. Oliveira Goncalves <sup>682a</sup>, J.L. Oliver <sup>5159</sup>, M.J.R. Olsson <sup>5159</sup>, A. Olszewski <sup>86</sup>,
J. Olszowska <sup>6</sup>86,*, Ö.O. Öncel <sup>54</sup>, D.C. O'Neil <sup>142</sup>, A.P. O'Neill <sup>19</sup>, A. Onofre <sup>130a,130e</sup>,
P.U.E. Onyisi <sup>11</sup>, M.J. Oreglia <sup>39</sup>, G.E. Orellana <sup>90</sup>, D. Orestano <sup>77a,77b</sup>, N. Orlando <sup>13</sup>,
R.S. Orr <sup>155</sup>, V. O'Shea <sup>59</sup>, R. Ospanov <sup>62a</sup>, G. Otero y Garzon <sup>30</sup>, H. Otono <sup>89</sup>,
P.S. Ott 6<sup>3a</sup>, G.J. Ottino 1<sup>7a</sup>, M. Ouchrif 3<sup>3d</sup>, J. Ouellette 2<sup>2</sup>, F. Ould-Saada 1<sup>2</sup>5,
M. Owen <sup>59</sup>, R.E. Owen <sup>134</sup>, K.Y. Oyulmaz <sup>21a</sup>, V.E. Ozcan <sup>21a</sup>, N. Ozturk <sup>8</sup>,
S. Ozturk <sup>1</sup>0<sup>21d</sup>, H.A. Pacey <sup>3</sup>2, K. Pachal <sup>5</sup>1, A. Pacheco Pages <sup>1</sup>3, C. Padilla Aranda <sup>1</sup>3,
G. Padovano <sup>6</sup>75a,75b, S. Pagan Griso <sup>6</sup>17a, G. Palacino <sup>68</sup>, A. Palazzo <sup>670a,70b</sup>, S. Palestini <sup>636</sup>,
J. Pan 171, T. Pan 64a, D.K. Panchal 11, C.E. Pandini 114, J.G. Panduro Vazquez 195,
H. Pang <sup>14b</sup>, P. Pani <sup>48</sup>, G. Panizzo <sup>69a,69c</sup>, L. Paolozzi <sup>56</sup>, C. Papadatos <sup>108</sup>, S. Parajuli <sup>44</sup>,
A. Paramonov 6, C. Paraskevopoulos 10, D. Paredes Hernandez 6, T.H. Park 15,
M.A. Parker <sup>©32</sup>, F. Parodi <sup>©57b,57a</sup>, E.W. Parrish <sup>©115</sup>, V.A. Parrish <sup>©52</sup>, J.A. Parsons <sup>©41</sup>,
U. Parzefall <sup>654</sup>, B. Pascual Dias <sup>6108</sup>, L. Pascual Dominguez <sup>6151</sup>, F. Pasquali <sup>6114</sup>,
E. Pasqualucci <sup>675a</sup>, S. Passaggio <sup>57b</sup>, F. Pastore <sup>695</sup>, P. Pasuwan <sup>647a,47b</sup>, P. Patel <sup>86</sup>,
U.M. Patel<sup>51</sup>, J.R. Pater<sup>51</sup>, T. Pauly<sup>53</sup>, J. Pearkes<sup>5143</sup>, M. Pedersen<sup>5125</sup>, R. Pedro<sup>5130a</sup>,
S.V. Peleganchuk <sup>37</sup>, O. Penc <sup>36</sup>, E.A. Pender H. Peng <sup>62a</sup>, K.E. Penski <sup>109</sup>, M. Penzin <sup>37</sup>,
B.S. Peralva 682d,82d, A.P. Pereira Peixoto 660, L. Pereira Sanchez 647a,47b, D.V. Perepelitsa 629,af,
E. Perez Codina <sup>156a</sup>, M. Perganti <sup>10</sup>, L. Perini <sup>10</sup>, H. Pernegger <sup>16</sup>, S. Perrella <sup>16</sup>,
A. Perrevoort <sup>113</sup>, O. Perrin <sup>40</sup>, K. Peters <sup>48</sup>, R.F.Y. Peters <sup>101</sup>, B.A. Petersen <sup>36</sup>,
T.C. Petersen 642, E. Petit 10102, V. Petousis 132, C. Petridou 152,e, A. Petrukhin 141,
M. Pettee 17a, N.E. Pettersson 36, A. Petukhov 37, K. Petukhova 133, A. Peyaud 135,
R. Pezoa 137f, L. Pezzotti 36, G. Pezzullo 171, T.M. Pham 169, T. Pham 1515,
P.W. Phillips 134, M.W. Phipps 161, G. Piacquadio 145, E. Pianori 17a, F. Piazza 71a,71b,
R. Piegaia 630, D. Pietreanu 27b, A.D. Pilkington 101, M. Pinamonti 69a,69c, J.L. Pinfold 2,
```

```
B.C. Pinheiro Pereira 130a, C. Pitman Donaldson, D.A. Pizzi 34, L. Pizzimento 76a,76b,
A. Pizzini <sup>114</sup>, M.-A. Pleier <sup>29</sup>, V. Plesanovs <sup>4</sup>, V. Pleskot <sup>133</sup>, E. Plotnikova <sup>8</sup>, G. Poddar <sup>4</sup>,
R. Poettgen 198, L. Poggioli 127, D. Pohl 24, I. Pokharel 55, S. Polacek 133, G. Polesello 573a,
A. Poley 142,156a, R. Polifka 132, A. Polini 23b, C.S. Pollard 166, Z.B. Pollock 119,
V. Polychronakos <sup>©</sup> <sup>29</sup>, E. Pompa Pacchi <sup>75a,75b</sup>, D. Ponomarenko <sup>©</sup> <sup>113</sup>, L. Pontecorvo <sup>©</sup> <sup>36</sup>,
S. Popa <sup>©</sup> <sup>27a</sup>, G.A. Popeneciu <sup>©</sup> <sup>27d</sup>, D.M. Portillo Quintero <sup>©</sup> <sup>156a</sup>, S. Pospisil <sup>©</sup> <sup>132</sup>,
P. Postolache <sup>©27c</sup>, K. Potamianos <sup>©126</sup>, P.P. Potepa <sup>©85a</sup>, I.N. Potrap <sup>©38</sup>, C.J. Potter <sup>©32</sup>,
H. Potti<sup>1</sup>, T. Poulsen<sup>48</sup>, J. Poveda<sup>1</sup>, M.E. Pozo Astigarraga<sup>36</sup>, A. Prades Ibanez<sup>162</sup>,
M.M. Prapa 646, D. Price 1011, M. Primavera 707a, M.A. Principe Martin 99, R. Privara 122,
M.L. Proffitt <sup>138</sup>, N. Proklova <sup>128</sup>, K. Prokofiev <sup>64c</sup>, G. Proto <sup>76a,76b</sup>, S. Protopopescu <sup>29</sup>,
J. Proudfoot <sup>6</sup>, M. Przybycien <sup>85a</sup>, W.W. Przygoda <sup>85b</sup>, J.E. Puddefoot <sup>139</sup>, D. Pudzha <sup>37</sup>,
D. Pyatiizbyantseva 637, J. Qian 6106, D. Qichen 6101, Y. Qin 6101, S. Qiu 617a, T. Qiu 52,
A. Quadt <sup>655</sup>, M. Queitsch-Maitland <sup>6101</sup>, G. Quetant <sup>656</sup>, G. Rabanal Bolanos <sup>61</sup>,
D. Rafanoharana <sup>6</sup> <sup>54</sup>, F. Ragusa <sup>6</sup> <sup>71a,71b</sup>, J.L. Rainbolt <sup>6</sup> <sup>39</sup>, J.A. Raine <sup>6</sup> <sup>56</sup>, S. Rajagopalan <sup>6</sup> <sup>29</sup>,
E. Ramakoti <sup>637</sup>, K. Ran <sup>48,14d</sup>, N.P. Rapheeha <sup>63a</sup>, V. Raskina <sup>6127</sup>, D.F. Rassloff <sup>63a</sup>,
S. Rave^{\bullet 100}, B. Ravina^{\bullet 55}, I. Ravinovich^{\bullet 168}, M. Raymond^{\bullet 36}, A.L. Read^{\bullet 125},
N.P. Readioff <sup>139</sup>, D.M. Rebuzzi <sup>73a,73b</sup>, G. Redlinger <sup>29</sup>, K. Reeves <sup>45</sup>, J.A. Reidelsturz <sup>170</sup>,
D. Reikher <sup>151</sup>, A. Rej <sup>141</sup>, C. Rembser <sup>36</sup>, A. Renardi <sup>48</sup>, M. Renda <sup>27b</sup>, M.B. Rendel <sup>110</sup>,
F. Renner <sup>648</sup>, A.G. Rennie <sup>59</sup>, S. Resconi <sup>671a</sup>, M. Ressegotti <sup>57b,57a</sup>, E.D. Resseguie <sup>617a</sup>,
S. Rettie <sup>136</sup>, J.G. Reyes Rivera <sup>107</sup>, B. Reynolds <sup>119</sup>, E. Reynolds <sup>17a</sup>, M. Rezaei Estabragh <sup>170</sup>,
O.L. Rezanova \mathbb{D}^{37}, P. Reznicek \mathbb{D}^{133}, N. Ribaric \mathbb{D}^{91}, E. Ricci \mathbb{D}^{78a,78b}, R. Richter \mathbb{D}^{110},
S. Richter (10<sup>47a,47b</sup>, E. Richter-Was (10<sup>85b</sup>, M. Ridel (10<sup>127</sup>, S. Ridouani (10<sup>35d</sup>, P. Rieck (10<sup>117</sup>),
P. Riedler <sup>136</sup>, M. Rijssenbeek <sup>145</sup>, A. Rimoldi <sup>73a,73b</sup>, M. Rimoldi <sup>48</sup>, L. Rinaldi <sup>23b,23a</sup>,
T.T. Rinn ^{\circ}<sup>29</sup>, M.P. Rinnagel ^{\circ}<sup>109</sup>, G. Ripellino ^{\circ}<sup>160</sup>, I. Riu ^{\circ}<sup>13</sup>, P. Rivadeneira ^{\circ}<sup>48</sup>,
J.C. Rivera Vergara 6164, F. Rizatdinova 6121, E. Rizvi 694, C. Rizzi 656, B.A. Roberts 6166,
B.R. Roberts <sup>17a</sup>, S.H. Robertson <sup>104,w</sup>, M. Robin <sup>48</sup>, D. Robinson <sup>32</sup>,
C.M. Robles Gajardo<sup>137</sup>f, M. Robles Manzano<sup>100</sup>, A. Robson<sup>59</sup>, A. Rocchi<sup>76a,76b</sup>,
C. Roda ^{\circ}7^{4a,74b}, S. Rodriguez Bosca ^{\circ}6^{3a}, Y. Rodriguez Garcia ^{\circ}2^{2a}, A. Rodriguez Rodriguez ^{\circ}4,
A.M. Rodríguez Vera <sup>156b</sup>, S. Roe<sup>36</sup>, J.T. Roemer <sup>159</sup>, A.R. Roepe-Gier <sup>136</sup>, J. Roggel <sup>170</sup>,
O. Røhne<sup>125</sup>, R.A. Rojas<sup>103</sup>, B. Roland<sup>154</sup>, C.P.A. Roland<sup>168</sup>, J. Roloff<sup>199</sup>,
A. Romaniouk^{\odot 37}, E. Romano^{\odot 73a,73b}, M. Romano^{\odot 23b}, A.C. Romero Hernandez^{\odot 161},
N. Rompotis • 92, L. Roos • 127, S. Rosati • 75a, B.J. Rosser • 39, E. Rossi • 4, E. Rossi • 72a,72b,
L.P. Rossi 57b, L. Rossini 48, R. Rosten 119, M. Rotaru 27b, B. Rottler 54, C. Rougier 102,ai,
D. Rousseau 66, D. Rousso 32, G. Rovelli 73a,73b, A. Roy 161, S. Roy-Garand 155,
A. Rozanov <sup>102</sup>, Y. Rozen <sup>150</sup>, X. Ruan <sup>33g</sup>, A. Rubio Jimenez <sup>162</sup>, A.J. Ruby <sup>92</sup>,
V.H. Ruelas Rivera <sup>18</sup>, T.A. Ruggeri <sup>1</sup>, F. Rühr <sup>15</sup>, A. Ruiz-Martinez <sup>16</sup>, A. Rummler <sup>16</sup>,
Z. Rurikova <sup>6</sup> <sup>54</sup>, N.A. Rusakovich <sup>6</sup> <sup>38</sup>, H.L. Russell <sup>6</sup> <sup>164</sup>, J.P. Rutherfoord <sup>6</sup> <sup>7</sup>, K. Rybacki <sup>91</sup>,
M. Rybar <sup>133</sup>, E.B. Rye <sup>125</sup>, A. Ryzhov <sup>37</sup>, J.A. Sabater Iglesias <sup>56</sup>, P. Sabatini <sup>162</sup>,
L. Sabetta <sup>675a,75b</sup>, H.F-W. Sadrozinski <sup>6136</sup>, F. Safai Tehrani <sup>675a</sup>, B. Safarzadeh Samani <sup>6146</sup>,
M. Safdari 143, S. Saha 144, M. Sahinsoy 110, M. Saimpert 135, M. Saito 153, T. Saito 153,
D. Salamani <sup>36</sup>, A. Salnikov <sup>143</sup>, J. Salt <sup>162</sup>, A. Salvador Salas <sup>13</sup>, D. Salvatore <sup>436</sup>, A.
F. Salvatore <sup>146</sup>, A. Salzburger <sup>36</sup>, D. Sammel <sup>54</sup>, D. Sampsonidis <sup>152</sup>,
```

D. Sampsonidou 62d,62c, J. Sánchez 162, A. Sanchez Pineda 4, V. Sanchez Sebastian 162,

H. Sandaker ¹²⁵, C.O. Sander ⁴⁸, J.A. Sandesara ¹⁰³, M. Sandhoff ¹⁷⁰, C. Sandoval ^{22b}, D.P.C. Sankey ¹³⁴, T. Sano ⁸⁷, A. Sansoni ⁵³, L. Santi ^{675a,75b}, C. Santoni ⁴⁰, H. Santos 130a, 130b, S.N. Santpur 17a, A. Santra 168, K.A. Saoucha 139, J.G. Saraiva 130a, 130d, J. Sardain ¹⁰⁷, O. Sasaki ¹⁰⁸³, K. Sato ¹⁵⁷, C. Sauer ^{63b}, F. Sauerburger ¹⁵⁴, E. Sauvan ¹⁵⁴, P. Savard ^{155,ac}, R. Sawada ¹⁵³, C. Sawyer ¹³⁴, L. Sawyer ⁹⁷, I. Sayago Galvan ¹⁶², C. Sbarra ^{©23b}, A. Sbrizzi ^{©23b,23a}, T. Scanlon ^{©96}, J. Schaarschmidt ^{©138}, P. Schacht ^{©110}, D. Schaefer ³⁹, U. Schäfer ¹⁰⁰, A.C. Schaffer ^{66,44}, D. Schaile ¹⁰⁹, R.D. Schamberger ¹⁴⁵, E. Schanet ¹⁰ C. Scharf ¹⁸, M.M. Schefer ¹⁹, V.A. Schegelsky ¹³, D. Scheirich ¹³, F. Schenck ¹⁸, M. Schernau ¹⁵⁹, C. Scheulen ⁵⁵, C. Schiavi ^{57b,57a}, Z.M. Schillaci ²⁶, E.J. Schioppa ⁶70a,70b, M. Schioppa ⁶43b,43a, B. Schlag ⁶100, K.E. Schleicher ⁶54, S. Schlenker ⁶36, J. Schmeing ¹⁷⁰, M.A. Schmidt ¹⁷⁰, K. Schmieden ¹⁰⁰, C. Schmitt ¹⁰⁰, S. Schmitt ⁴⁸, L. Schoeffel ¹³⁵, A. Schoening ^{63b}, P.G. Scholer ⁵⁴, E. Schopf ¹²⁶, M. Schott ¹⁰⁰, J. Schovancova ⁶³⁶, S. Schramm ⁵⁶, F. Schroeder ⁶¹⁷⁰, H-C. Schultz-Coulon ^{63a}, M. Schumacher ⁵⁴, B.A. Schumm ⁵¹³⁶, Ph. Schune ⁵¹³⁵, H.R. Schwartz ⁵¹³⁶, A. Schwartzman ¹⁴³, T.A. Schwarz ¹⁰⁶, Ph. Schwemling ¹³⁵, R. Schwienhorst ¹⁰⁷, A. Sciandra ¹³⁶, G. Sciolla ²⁶, F. Scuri ^{74a}, F. Scutti ¹⁰⁵, C.D. Sebastiani ⁹², K. Sedlaczek ⁴⁹, P. Seema ¹⁸, S.C. Seidel ¹¹², A. Seiden ¹³⁶, B.D. Seidlitz ⁴¹, C. Seitz ⁴⁸, J.M. Seixas ^{82b}, G. Sekhniaidze 0^{72a} , S.J. Sekula 0^{44} , L. Selem 0^4 , N. Semprini-Cesari $0^{23b,23a}$, S. Sen 0^{51} , D. Sengupta 56, V. Senthilkumar 5162, L. Serin 566, L. Serkin 569a,69b, M. Sessa 77a,77b, H. Severini ¹²⁰, F. Sforza ^{57b,57a}, A. Sfyrla ⁵⁶, E. Shabalina ⁵⁵, R. Shaheen ¹⁴⁴, J.D. Shahinian ¹²⁸, D. Shaked Renous ¹⁶⁸, L.Y. Shan ^{14a}, M. Shapiro ^{17a}, A. Sharma ³⁶, A.S. Sharma ¹⁶³, P. Sharma ⁸⁰, S. Sharma ⁴⁸, P.B. Shatalov ³⁷, K. Shaw ¹⁴⁶, S.M. Shaw $^{\bullet 101}$, Q. Shen $^{\bullet 62c,5}$, P. Sherwood $^{\bullet 96}$, L. Shi $^{\bullet 96}$, C.O. Shimmin $^{\bullet 171}$, Y. Shimogama ¹⁶⁷, J.D. Shinner ⁹⁵, I.P.J. Shipsey ¹²⁶, S. Shirabe ⁶⁰, M. Shiyakova ³⁸, J. Shlomi ¹⁶⁸, M.J. Shochet ³⁹, J. Shojaii ¹⁰⁵, D.R. Shope ¹²⁵, S. Shrestha ^{119,ag}, E.M. Shrif 633g, M.J. Shroff 6164, P. Sicho 6131, A.M. Sickles 6161, E. Sideras Haddad 633g, A. Sidoti ^{23b}, F. Siegert ⁵⁰, Dj. Sijacki ¹⁵, R. Sikora ^{85a}, F. Sili ⁹⁰, J.M. Silva ²⁰, M.V. Silva Oliveira 636, S.B. Silverstein 647a, S. Simion 66, R. Simoniello 636, E.L. Simpson 59, H. Simpson ¹⁴⁶, L.R. Simpson ¹⁰⁶, N.D. Simpson ⁹⁸, S. Simsek ¹²¹, S. Sindhu ¹⁵⁵, P. Sinervo 155, S. Singh 142, S. Singh 155, S. Sinha 48, S. Sinha 33g, M. Sioli 23b,23a, I. Siral 636, S.Yu. Sivoklokov 637,*, J. Sjölin 647a,47b, A. Skaf 55, E. Skorda 98, P. Skubic 120, M. Slawinska 686, V. Smakhtin 8, B.H. Smart 134, J. Smiesko 36, S.Yu. Smirnov 37, Y. Smirnov ⁶³⁷, L.N. Smirnova ^{637,a}, O. Smirnova ⁶⁹⁸, A.C. Smith ⁶⁴¹, E.A. Smith ⁶³⁹, H.A. Smith ¹²⁶, J.L. Smith ⁹², R. Smith ¹⁴³, M. Smizanska ⁹¹, K. Smolek ¹³², A. Smykiewicz 686, A.A. Snesarev 637, H.L. Snoek 6114, S. Snyder 629, R. Sobie 6164, w, A. Soffer ¹⁵¹, C.A. Solans Sanchez ¹⁵⁶, E.Yu. Soldatov ¹⁵⁷, U. Soldevila ¹⁵¹, A.A. Solodkov ¹⁵⁷, S. Solomon ⁵⁴, A. Soloshenko ³⁸, K. Solovieva ⁵⁴, O.V. Solovyanov ⁴⁰, V. Solovyev ³⁷, P. Sommer • 36, A. Sonay • 13, W.Y. Song • 156b, J.M. Sonneveld • 114, A. Sopczak • 132, A.L. Sopio ⁶⁹⁶, F. Sopkova ^{628b}, V. Sothilingam ^{63a}, S. Sottocornola ⁶⁸, R. Soualah ^{6116c}, Z. Soumaimi ^{635e}, D. South ⁶⁴⁸, S. Spagnolo ^{670a,70b}, M. Spalla ⁶¹¹⁰, D. Sperlich ⁵⁴, G. Spigo ³⁶, M. Spina ¹⁴⁶, S. Spinali ⁹¹, D.P. Spiteri ⁵⁹, M. Spousta ¹³³, E.J. Staats ³⁴,

A. Stabile 071a,71b , R. Stamen 63a , M. Stamenkovic 0114 , A. Stampekis 020 , M. Standke 024 , E. Stanecka 086 , M.V. Stange 050 , B. Stanislaus 017a , M.M. Stanitzki 048 , M. Stankaityte 0126 ,

B. Stapf • 48, E.A. Starchenko • 37, G.H. Stark • 136, J. Stark • 102, ai, D.M. Starko 156b, P. Staroba¹³¹, P. Starovoitov^{63a}, S. Stärz¹⁰¹⁰⁴, R. Staszewski⁸⁶, G. Stavropoulos⁴⁶, J. Steentoft 6160, P. Steinberg 629, B. Stelzer 142,156a, H.J. Stelzer 129, O. Stelzer-Chilton 156a, H. Stenzel⁵⁸, T.J. Stevenson¹⁴⁶, G.A. Stewart³⁶, J.R. Stewart¹²¹, M.C. Stockton³⁶, G. Stoicea ^{©27b}, M. Stolarski ^{©130a}, S. Stonjek ^{©110}, A. Straessner ^{©50}, J. Strandberg ^{©144}, S. Strandberg ⁶4^{7a,47b}, M. Strauss ⁶1²⁰, T. Strebler ⁶1⁰², P. Strizenec ⁶2^{8b}, R. Ströhmer ⁶1⁶⁵, D.M. Strom 123, L.R. Strom 48, R. Stroynowski 44, A. Strubig 47a,47b, S.A. Stucci 29, B. Stugu 16, J. Stupak 120, N.A. Styles 48, D. Su 143, S. Su 62a, W. Su 62d, 138,62c, X. Su^{62a,66}, K. Sugizaki¹⁵³, V.V. Sulin³⁷, M.J. Sullivan⁹², D.M.S. Sultan^{78a,78b}, L. Sultanaliyeva ³⁷, S. Sultansoy ³, T. Sumida ⁸⁷, S. Sun ¹⁰⁶, S. Sun ¹⁶⁹, O. Sunneborn Gudnadottir ¹⁶⁰, M.R. Sutton ¹⁴⁶, M. Svatos ¹³¹, M. Swiatlowski ^{156a}, T. Swirski $^{\bullet 165}$, I. Sykora $^{\bullet 28a}$, M. Sykora $^{\bullet 133}$, T. Sykora $^{\bullet 133}$, D. Ta $^{\bullet 100}$, K. Tackmann $^{\bullet 48,u}$, A. Taffard ¹⁵⁹, R. Tafirout ^{156a}, J.S. Tafoya Vargas ⁶⁶, R.H.M. Taibah ¹²⁷, R. Takashima ⁸⁸, E.P. Takeva • 52, Y. Takubo • 83, M. Talby • 102, A.A. Talyshev • 37, K.C. Tam • 64b, N.M. Tamir¹⁵¹, A. Tanaka¹⁵³, J. Tanaka¹⁵³, R. Tanaka¹⁵⁶, M. Tanasini^{157b,57a}, J. Tang^{62c}, Z. Tao 163, S. Tapia Araya 137, S. Tapprogge 100, A. Tarek Abouelfadl Mohamed 1017, S. Tarem¹⁵⁰, K. Tariq^{62b}, G. Tarna^{102,27b}, G.F. Tartarelli^{671a}, P. Tas¹³³, M. Tasevsky 131, E. Tassi 143b,43a, A.C. Tate 1616, G. Tateno 153, Y. Tayalati 153e,v, G.N. Taylor ¹⁰⁵, W. Taylor ^{156b}, H. Teagle ⁹², A.S. Tee ¹⁶⁹, R. Teixeira De Lima ¹⁴³, P. Teixeira-Dias •95, J.J. Teoh •155, K. Terashi •153, J. Terron •99, S. Terzo •13, M. Testa •53, R.J. Teuscher ^{155,w}, A. Thaler ⁷⁹, O. Theiner ⁵⁶, N. Themistokleous ⁵², T. Theveneaux-Pelzer 1010, O. Thielmann 1710, D.W. Thomas 1520, E.A. Thompson $^{\bullet 17a}$, P.D. Thompson $^{\bullet 20}$, E. Thomson $^{\bullet 128}$, E.J. Thorpe $^{\bullet 94}$, Y. Tian $^{\bullet 55}$, V. Tikhomirov $0^{37,a}$, Yu.A. Tikhonov 0^{37} , S. Timoshenko³⁷, E.X.L. Ting 0^{1} , P. Tipton 0^{171} , S.H. Tlou \mathbb{D}^{33g} , A. Tnourji \mathbb{D}^{40} , K. Todome $\mathbb{D}^{23b,23a}$, S. Todorova-Nova \mathbb{D}^{133} , S. Todt⁵⁰, M. Togawa 683, J. Tojo 689, S. Tokár 628a, K. Tokushuku 683, O. Toldaiev 668, R. Tombs 632, M. Tomoto 683,111, L. Tompkins 6143,am, K.W. Topolnicki 685b, P. Tornambe 6103, E. Torrence ¹²³, H. Torres ⁵⁰, E. Torró Pastor ¹⁶², M. Toscani ³⁰, C. Tosciri ³⁹, M. Tost ¹¹, D.R. Tovey ¹³⁹, A. Traeet¹⁶, I.S. Trandafir ^{27b}, T. Trefzger ¹⁶⁵, A. Tricoli ²⁹, I.M. Trigger 156a, S. Trincaz-Duvoid 127, D.A. Trischuk 26, B. Trocmé 66, C. Troncon 71a, L. Truong $^{\bullet 33c}$, M. Trzebinski $^{\bullet 86}$, A. Trzupek $^{\bullet 86}$, F. Tsai $^{\bullet 145}$, M. Tsai $^{\bullet 106}$, A. Tsiamis $^{\bullet 152,e}$, P.V. Tsiareshka³⁷, S. Tsigaridas o^{156a}, A. Tsirigotis o^{152,s}, V. Tsiskaridze o¹⁴⁵, E.G. Tskhadadze^{149a}, M. Tsopoulou^{©152,e}, Y. Tsujikawa^{©87}, I.I. Tsukerman^{©37}, V. Tsulaia^{©17a}, S. Tsuno 683 , O. Tsur 150 , D. Tsybychev 6145 , Y. Tu 64b , A. Tudorache 627b , V. Tudorache 627b , A.N. Tuna ⁶³⁶, S. Turchikhin ⁶³⁸, I. Turk Cakir ^{63a}, R. Turra ^{671a}, T. Turtuvshin ^{638,x}, P.M. Tuts • 41, S. Tzamarias • 152,e, P. Tzanis • 10, E. Tzovara • 100, K. Uchida 153, F. Ukegawa • 157, P.A. Ulloa Poblete 137c, E.N. Umaka 29, G. Unal 36, M. Unal 11, A. Undrus 29, G. Unel 159, J. Urban ^{28b}, P. Urquijo ¹⁰⁵, G. Usai ⁸, R. Ushioda ¹⁵⁴, M. Usman ¹⁰⁸, Z. Uysal ^{21b}, L. Vacavant ¹⁰ V. Vacek ¹³², B. Vachon ¹⁰⁴, K.O.H. Vadla ¹²⁵, T. Vafeiadis ¹³⁶, A. Vaitkus ⁶⁹⁶, C. Valderanis ¹⁰⁹, E. Valdes Santurio ^{47a,47b}, M. Valente ^{156a}, S. Valentinetti $0^{23b,23a}$, A. Valero 0^{162} , A. Vallier $0^{102,ai}$, J.A. Valls Ferrer 0^{162} , D.R. Van Arneman ¹¹⁴, T.R. Van Daalen ¹³⁸, P. Van Gemmeren ⁶, M. Van Rijnbach ^{125,36},

S. Van Stroud ⁶ ⁹⁶, I. Van Vulpen ⁶ ¹¹⁴, M. Vanadia ⁶ ^{76a,76b}, W. Vandelli ⁶,

```
M. Vandenbroucke <sup>135</sup>, E.R. Vandewall <sup>121</sup>, D. Vannicola <sup>151</sup>, L. Vannoli <sup>57b,57a</sup>, R. Vari <sup>75a</sup>,
E.W. Varnes <sup>67</sup>, C. Varni <sup>617a</sup>, T. Varol <sup>6148</sup>, D. Varouchas <sup>66</sup>, L. Varriale <sup>6162</sup>,
K.E. Varvell 147, M.E. Vasile 276, L. Vaslin 4, G.A. Vasquez 164, F. Vazeille 4,
T. Vazquez Schroeder <sup>636</sup>, J. Veatch <sup>31</sup>, V. Vecchio <sup>6101</sup>, M.J. Veen <sup>6103</sup>, I. Veliscek <sup>6126</sup>,
L.M. Veloce <sup>155</sup>, F. Veloso <sup>130a,130c</sup>, S. Veneziano <sup>75a</sup>, A. Ventura <sup>70a,70b</sup>, A. Verbytskyi <sup>110</sup>,
M. Verducci \mathbb{D}^{74a,74b}, C. Vergis \mathbb{D}^{24}, M. Verissimo De Araujo \mathbb{D}^{82b}, W. Verkerke \mathbb{D}^{114},
J.C. Vermeulen <sup>114</sup>, C. Vernieri <sup>143</sup>, P.J. Verschuuren <sup>95</sup>, M. Vessella <sup>103</sup>, M.C. Vetterli <sup>142,ac</sup>,
A. Vgenopoulos <sup>152,e</sup>, N. Viaux Maira <sup>137f</sup>, T. Vickey <sup>139</sup>, O.E. Vickey Boeriu <sup>139</sup>,
G.H.A. Viehhauser <sup>126</sup>, L. Vigani <sup>63b</sup>, M. Villa <sup>23b,23a</sup>, M. Villaplana Perez <sup>162</sup>,
E.M. Villhauer<sup>52</sup>, E. Vilucchi <sup>653</sup>, M.G. Vincter <sup>634</sup>, G.S. Virdee <sup>620</sup>, A. Vishwakarma <sup>652</sup>,
C. Vittori 636, I. Vivarelli 6146, V. Vladimirov 66, E. Voevodina 6110, F. Vogel 1019, P. Vokac 6132,
J. Von Ahnen <sup>648</sup>, E. Von Toerne <sup>624</sup>, B. Vormwald <sup>636</sup>, V. Vorobel <sup>613</sup>, K. Vorobev <sup>637</sup>,
M. Vos 6 162, K. Voss 6 141, J.H. Vossebeld 6 92, M. Vozak 6 114, L. Vozdecky 6 94, N. Vranjes 6 15,
M. Vranjes Milosavljevic <sup>15</sup>, M. Vreeswijk <sup>114</sup>, R. Vuillermet <sup>36</sup>, O. Vujinovic <sup>100</sup>,
I. Vukotic <sup>139</sup>, S. Wada <sup>157</sup>, C. Wagner <sup>103</sup>, J.M. Wagner <sup>170</sup>, W. Wagner <sup>170</sup>, S. Wahdan <sup>170</sup>,
H. Wahlberg <sup>190</sup>, R. Wakasa <sup>157</sup>, M. Wakida <sup>111</sup>, J. Walder <sup>134</sup>, R. Walker <sup>109</sup>,
W. Walkowiak 141, A.M. Wang 161, A.Z. Wang 169, C. Wang 160, C. Wang 162c, H. Wang 17a,
J. Wang 64a, R.-J. Wang 100, R. Wang 61, R. Wang 66, S.M. Wang 148, S. Wang 62b,
T. Wang 62a, W.T. Wang 80, X. Wang 14c, X. Wang 161, X. Wang 62c, Y. Wang 62d,
Y. Wang 14c, Z. Wang 15c, Z. Wang 15c, Z. Wang 15c, Z. Wang 15c, A. Warburton 15c,
R.J. Ward <sup>620</sup>, N. Warrack <sup>59</sup>, A.T. Watson <sup>620</sup>, H. Watson <sup>59</sup>, M.F. Watson <sup>620</sup>,
G. Watts <sup>138</sup>, B.M. Waugh <sup>96</sup>, C. Weber <sup>29</sup>, H.A. Weber <sup>18</sup>, M.S. Weber <sup>19</sup>,
S.M. Weber 63a, C. Wei 62a, Y. Wei 6126, A.R. Weidberg 6126, E.J. Weik 6117, J. Weingarten 649,
M. Weirich 100, C. Weiser 54, C.J. Wells 48, T. Wenaus 29, B. Wendland 49, T. Wengler 36,
N.S. Wenke<sup>110</sup>, N. Wermes<sup>©24</sup>, M. Wessels<sup>©63a</sup>, K. Whalen<sup>©123</sup>, A.M. Wharton<sup>©91</sup>,
A.S. White 61, A. White 8, M.J. White 11, D. Whiteson 159, L. Wickremasinghe 124,
W. Wiedenmann <sup>169</sup>, C. Wiel <sup>50</sup>, M. Wielers <sup>134</sup>, C. Wiglesworth <sup>642</sup>, L.A.M. Wiik-Fuchs <sup>54</sup>,
D.J. Wilbern<sup>120</sup>, H.G. Wilkens<sup>136</sup>, D.M. Williams<sup>141</sup>, H.H. Williams<sup>128</sup>, S. Williams<sup>132</sup>,
S. Willocq 6103, B.J. Wilson 6101, P.J. Windischhofer 639, F. Winklmeier 6123, B.T. Winter 54,
J.K. Winter <sup>101</sup>, M. Wittgen <sup>143</sup>, M. Wobisch <sup>197</sup>, R. Wölker <sup>126</sup>, J. Wollrath <sup>159</sup>,
M.W. Wolter <sup>686</sup>, H. Wolters <sup>6130a,130c</sup>, V.W.S. Wong <sup>6163</sup>, A.F. Wongel <sup>648</sup>, S.D. Worm <sup>648</sup>,
B.K. Wosiek 686, K.W. Woźniak 86, K. Wraight 59, J. Wu 14a,14d, M. Wu 64a, M. Wu 113,
S.L. Wu<sup>169</sup>, X. Wu<sup>56</sup>, Y. Wu<sup>62a</sup>, Z. Wu<sup>5135,62a</sup>, J. Wuerzinger<sup>5110</sup>, T.R. Wyatt<sup>5101</sup>,
B.M. Wynne <sup>52</sup>, S. Xella <sup>42</sup>, L. Xia <sup>14c</sup>, M. Xia <sup>4b</sup>, J. Xiang <sup>64c</sup>, X. Xiao <sup>106</sup>, M. Xie <sup>62a</sup>,
X. Xie 62a, S. Xin 14a, 14d, J. Xiong 17a, I. Xiotidis 46, D. Xu 14a, H. Xu 24a, H. Xu 62a,
L. Xu<sup>6</sup><sup>62a</sup>, R. Xu<sup>128</sup>, T. Xu<sup>16</sup><sup>106</sup>, Y. Xu<sup>16</sup><sup>14b</sup>, Z. Xu<sup>16</sup><sup>62b</sup>, Z. Xu<sup>16</sup><sup>14a</sup>, B. Yabsley<sup>16</sup><sup>147</sup>,
S. Yacoob \mathbb{D}^{33a}, N. Yamaguchi \mathbb{D}^{89}, Y. Yamaguchi \mathbb{D}^{154}, H. Yamauchi \mathbb{D}^{157}, T. Yamazaki \mathbb{D}^{17a},
Y. Yamazaki 6<sup>84</sup>, J. Yan 6<sup>2c</sup>, S. Yan 6<sup>126</sup>, Z. Yan 6<sup>25</sup>, H.J. Yang 6<sup>2c</sup>, H.T. Yang 6<sup>62a</sup>,
S. Yang 62a, T. Yang 64c, X. Yang 62a, X. Yang 14a, Y. Yang 44, Y. Yang 4a,
Z. Yang 62a,106, W-M. Yao 17a, Y.C. Yap 48, H. Ye 14c, H. Ye 55, J. Ye 44, S. Ye 29,
X. Ye 6<sup>2a</sup>, Y. Yeh 9<sup>96</sup>, I. Yeletskikh 3<sup>8</sup>, B.K. Yeo 1<sup>7a</sup>, M.R. Yexley 9<sup>91</sup>, P. Yin 4<sup>1</sup>,
K. Yorita 6167, S. Younas 276, C.J.S. Young 54, C. Young 143, Y. Yu 62a, M. Yuan 166,
R. Yuan ^{\bullet 62b,k}, L. Yue ^{\bullet 96}, M. Zaazoua ^{\bullet 35e}, B. Zabinski ^{\bullet 86}, E. Zaid ^{52}, T. Zakareishvili ^{\bullet 149b},
```

- N. Zakharchuk ¹³⁴, S. Zambito ¹⁵⁶, J.A. Zamora Saa ^{137d,137b}, J. Zang ¹⁵³, D. Zanzi ¹⁵⁴,
- O. Zaplatilek ¹³², C. Zeitnitz ¹⁷⁰, H. Zeng ^{14a}, J.C. Zeng ¹⁶¹, D.T. Zenger Jr ²⁶,
- O. Zenin 637, T. Ženiš 628a, S. Zenz 694, S. Zerradi 635a, D. Zerwas 666, M. Zhai 614a,14d,
- B. Zhang ^{14c}, D.F. Zhang ¹³⁹, J. Zhang ^{62b}, J. Zhang ⁶, K. Zhang ^{14a,14d}, L. Zhang ^{14c},
- P. Zhang 14a, 14d, R. Zhang 169, S. Zhang 106, T. Zhang 153, X. Zhang 162c, X. Zhang 162b,
- Y. Zhang 62c,5, Z. Zhang 17a, Z. Zhang 66, H. Zhao 138, P. Zhao 51, T. Zhao 62b,
- Y. Zhao 6136, Z. Zhao 662a, A. Zhemchugov 138, X. Zheng 62a, Z. Zheng 143, D. Zhong 161,
- B. Zhou¹⁰⁶, C. Zhou¹⁰⁶, H. Zhou¹⁰⁷, N. Zhou^{1062c}, Y. Zhou⁷, C.G. Zhu^{1062b}, H.L. Zhu^{1062a},
- J. Zhu 106, Y. Zhu 166, Y. Zhu 166, Y. Zhu 166, X. Zhuang 1614, K. Zhukov 1637, V. Zhulanov 1637,
- N.I. Zimine ¹³⁸, J. Zinsser ^{63b}, M. Ziolkowski ¹⁴¹, L. Živković ¹⁵, A. Zoccoli ^{23b,23a},
- K. Zoch 656, T.G. Zorbas 6139, O. Zormpa 646, W. Zou 641, L. Zwalinski 536
 - ¹ Department of Physics, University of Adelaide, Adelaide; Australia
 - ² Department of Physics, University of Alberta, Edmonton AB; Canada
 - ³ (a) Department of Physics, Ankara University, Ankara; (b) Division of Physics, TOBB University of Economics and Technology, Ankara; Türkiye
 - ⁴ LAPP, Université Savoie Mont Blanc, CNRS/IN2P3, Annecy; France
 - ⁵ APC, Université Paris Cité, CNRS/IN2P3, Paris; France
 - ⁶ High Energy Physics Division, Argonne National Laboratory, Argonne IL; United States of America
 - ⁷ Department of Physics, University of Arizona, Tucson AZ; United States of America
 - ⁸ Department of Physics, University of Texas at Arlington, Arlington TX; United States of America
 - ⁹ Physics Department, National and Kapodistrian University of Athens, Athens; Greece
 - 10 Physics Department, National Technical University of Athens, Zografou; Greece
 - ¹¹ Department of Physics, University of Texas at Austin, Austin TX; United States of America
 - ¹² Institute of Physics, Azerbaijan Academy of Sciences, Baku; Azerbaijan
 - ¹³ Institut de Física d'Altes Energies (IFAE), Barcelona Institute of Science and Technology, Barcelona; Spain
 - ¹⁴ (a) Institute of High Energy Physics, Chinese Academy of Sciences, Beijing; (b) Physics Department, Tsinghua University, Beijing; (c) Department of Physics, Nanjing University, Nanjing; (d) University of Chinese Academy of Science (UCAS), Beijing; China
 - ¹⁵ Institute of Physics, University of Belgrade, Belgrade; Serbia
 - ¹⁶ Department for Physics and Technology, University of Bergen, Bergen; Norway
- ^{17 (a)} Physics Division, Lawrence Berkeley National Laboratory, Berkeley CA; ^(b) University of California, Berkeley CA; United States of America
- ¹⁸ Institut für Physik, Humboldt Universität zu Berlin, Berlin; Germany
- ¹⁹ Albert Einstein Center for Fundamental Physics and Laboratory for High Energy Physics, University of Bern, Bern; Switzerland
- ²⁰ School of Physics and Astronomy, University of Birmingham, Birmingham; United Kingdom
- ^{21 (a)} Department of Physics, Bogazici University, Istanbul; ^(b) Department of Physics Engineering, Gaziantep University, Gaziantep; ^(c) Department of Physics, Istanbul University, Istanbul; ^(d) Istinye University, Sariyer, Istanbul; Türkiye
- ^{22 (a)} Facultad de Ciencias y Centro de Investigaciónes, Universidad Antonio Nariño, Bogotá; (b) Departamento de Física, Universidad Nacional de Colombia, Bogotá; Colombia
- ^{23 (a)} Dipartimento di Fisica e Astronomia A. Righi, Università di Bologna, Bologna; ^(b) INFN Sezione di Bologna; Italy
- ²⁴ Physikalisches Institut, Universität Bonn, Bonn; Germany
- ²⁵ Department of Physics, Boston University, Boston MA; United States of America
- ²⁶ Department of Physics, Brandeis University, Waltham MA; United States of America
- ²⁷ (a) Transilvania University of Brasov, Brasov; (b) Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest; (c) Department of Physics, Alexandru Ioan Cuza University of Iasi,

- Iasi; (d) National Institute for Research and Development of Isotopic and Molecular Technologies, Physics Department, Cluj-Napoca; (e) University Politehnica Bucharest, Bucharest; (f) West University in Timisoara, Timisoara; (g) Faculty of Physics, University of Bucharest, Bucharest; Romania
- ^{28 (a)} Faculty of Mathematics, Physics and Informatics, Comenius University, Bratislava; (b) Department of Subnuclear Physics, Institute of Experimental Physics of the Slovak Academy of Sciences, Kosice; Slovak Republic
- ²⁹ Physics Department, Brookhaven National Laboratory, Upton NY; United States of America
- ³⁰ Universidad de Buenos Aires, Facultad de Ciencias Exactas y Naturales, Departamento de Física, y CONICET, Instituto de Física de Buenos Aires (IFIBA), Buenos Aires; Argentina
- 31 California State University, CA; United States of America
- ³² Cavendish Laboratory, University of Cambridge, Cambridge; United Kingdom
- ³³ (a) Department of Physics, University of Cape Town, Cape Town; (b) iThemba Labs, Western Cape; (c) Department of Mechanical Engineering Science, University of Johannesburg, Johannesburg; (d) National Institute of Physics, University of the Philippines Diliman (Philippines); (e) University of South Africa, Department of Physics, Pretoria; (f) University of Zululand, KwaDlangezwa; (g) School of Physics, University of the Witwatersrand, Johannesburg; South Africa
- ³⁴ Department of Physics, Carleton University, Ottawa ON; Canada
- ^{35 (a)} Faculté des Sciences Ain Chock, Réseau Universitaire de Physique des Hautes Energies Université Hassan II, Casablanca; ^(b) Faculté des Sciences, Université Ibn-Tofail, Kénitra; ^(c) Faculté des Sciences Semlalia, Université Cadi Ayyad, LPHEA-Marrakech; ^(d) LPMR, Faculté des Sciences, Université Mohamed Premier, Oujda; ^(e) Faculté des sciences, Université Mohammed V, Rabat; ^(f) Institute of Applied Physics, Mohammed VI Polytechnic University, Ben Guerir; Morocco
- 36 CERN, Geneva; Switzerland
- 37 Affiliated with an institute covered by a cooperation agreement with CERN
- 38 Affiliated with an international laboratory covered by a cooperation agreement with CERN
- ³⁹ Enrico Fermi Institute, University of Chicago, Chicago IL; United States of America
- 40 LPC, Universit\'e Clermont Auvergne, CNRS/IN2P3, Clermont-Ferrand; France
- ⁴¹ Nevis Laboratory, Columbia University, Irvington NY; United States of America
- ⁴² Niels Bohr Institute, University of Copenhagen, Copenhagen; Denmark
- ⁴³ (a) Dipartimento di Fisica, Università della Calabria, Rende; (b) INFN Gruppo Collegato di Cosenza, Laboratori Nazionali di Frascati; Italy
- ⁴⁴ Physics Department, Southern Methodist University, Dallas TX; United States of America
- ⁴⁵ Physics Department, University of Texas at Dallas, Richardson TX; United States of America
- ⁴⁶ National Centre for Scientific Research "Demokritos", Agia Paraskevi; Greece
- ⁴⁷ (a) Department of Physics, Stockholm University; (b) Oskar Klein Centre, Stockholm; Sweden
- ⁴⁸ Deutsches Elektronen-Synchrotron DESY, Hamburg and Zeuthen; Germany
- ⁴⁹ Fakultät Physik, Technische Universität Dortmund, Dortmund; Germany
- ⁵⁰ Institut für Kern- und Teilchenphysik, Technische Universität Dresden, Dresden; Germany
- ⁵¹ Department of Physics, Duke University, Durham NC; United States of America
- ⁵² SUPA School of Physics and Astronomy, University of Edinburgh, Edinburgh; United Kingdom
- ⁵³ INFN e Laboratori Nazionali di Frascati, Frascati; Italy
- ⁵⁴ Physikalisches Institut, Albert-Ludwigs-Universität Freiburg, Freiburg; Germany
- ⁵⁵ II. Physikalisches Institut, Georg-August-Universität Göttingen, Göttingen; Germany
- ⁵⁶ Département de Physique Nucléaire et Corpusculaire, Université de Genève, Genève; Switzerland
- ⁵⁷ (a) Dipartimento di Fisica, Università di Genova, Genova; (b) INFN Sezione di Genova; Italy
- ⁵⁸ II. Physikalisches Institut, Justus-Liebig-Universität Giessen, Giessen; Germany
- ⁵⁹ SUPA School of Physics and Astronomy, University of Glasgow, Glasgow; United Kingdom
- ⁶⁰ LPSC, Université Grenoble Alpes, CNRS/IN2P3, Grenoble INP, Grenoble; France
- ⁶¹ Laboratory for Particle Physics and Cosmology, Harvard University, Cambridge MA; United States of America
- ⁶² (a) Department of Modern Physics and State Key Laboratory of Particle Detection and Electronics, University of Science and Technology of China, Hefei; (b) Institute of Frontier and Interdisciplinary Science and Key Laboratory of Particle Physics and Particle Irradiation (MOE), Shandong University,

- Qingdao; ^(c) School of Physics and Astronomy, Shanghai Jiao Tong University, Key Laboratory for Particle Astrophysics and Cosmology (MOE), SKLPPC, Shanghai; ^(d) Tsung-Dao Lee Institute, Shanghai; China
- ⁶³ (a) Kirchhoff-Institut für Physik, Ruprecht-Karls-Universität Heidelberg, Heidelberg; (b) Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg; Germany
- ⁶⁴ (a) Department of Physics, Chinese University of Hong Kong, Shatin, N.T., Hong Kong; (b) Department of Physics, University of Hong Kong, Hong Kong; (c) Department of Physics and Institute for Advanced Study, Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong; China
- 65 Department of Physics, National Tsing Hua University, Hsinchu; Taiwan
- ⁶⁶ IJCLab, Université Paris-Saclay, CNRS/IN2P3, 91405, Orsay; France
- 67 Centro Nacional de Microelectrónica (IMB-CNM-CSIC), Barcelona; Spain
- ⁶⁸ Department of Physics, Indiana University, Bloomington IN; United States of America
- ⁶⁹ (a) INFN Gruppo Collegato di Udine, Sezione di Trieste, Udine; (b) ICTP, Trieste; (c) Dipartimento Politecnico di Ingegneria e Architettura, Università di Udine, Udine; Italy
- ⁷⁰ (a) INFN Sezione di Lecce; ^(b) Dipartimento di Matematica e Fisica, Università del Salento, Lecce; Italy
- ⁷¹ (a) INFN Sezione di Milano; ^(b) Dipartimento di Fisica, Università di Milano, Milano; Italy
- ⁷² (a) INFN Sezione di Napoli; (b) Dipartimento di Fisica, Università di Napoli, Napoli; Italy
- ⁷³ (a) INFN Sezione di Pavia; (b) Dipartimento di Fisica, Università di Pavia, Pavia; Italy
- ⁷⁴ (a) INFN Sezione di Pisa; ^(b) Dipartimento di Fisica E. Fermi, Università di Pisa, Pisa; Italy
- ⁷⁵ (a) INFN Sezione di Roma; ^(b) Dipartimento di Fisica, Sapienza Università di Roma, Roma; Italy
- ⁷⁶ (a) INFN Sezione di Roma Tor Vergata; (b) Dipartimento di Fisica, Università di Roma Tor Vergata, Roma: Italy
- ⁷⁷ (a) INFN Sezione di Roma Tre; (b) Dipartimento di Matematica e Fisica, Università Roma Tre, Roma; Italy
- ⁷⁸ (a) INFN-TIFPA; (b) Università degli Studi di Trento, Trento; Italy
- ⁷⁹ Universität Innsbruck, Department of Astro and Particle Physics, Innsbruck; Austria
- ⁸⁰ University of Iowa, Iowa City IA; United States of America
- ⁸¹ Department of Physics and Astronomy, Iowa State University, Ames IA; United States of America
- 82 (a) Departamento de Engenharia Elétrica, Universidade Federal de Juiz de Fora (UFJF), Juiz de Fora; Universidade Federal do Rio De Janeiro COPPE/EE/IF, Rio de Janeiro; Instituto de Física, Universidade de São Paulo, São Paulo; de Janeiro State University, Rio de Janeiro; Brazil
- 83 KEK, High Energy Accelerator Research Organization, Tsukuba; Japan
- ⁸⁴ Graduate School of Science, Kobe University, Kobe; Japan
- ⁸⁵ (a) AGH University of Science and Technology, Faculty of Physics and Applied Computer Science, Krakow; (b) Marian Smoluchowski Institute of Physics, Jagiellonian University, Krakow; Poland
- ⁸⁶ Institute of Nuclear Physics Polish Academy of Sciences, Krakow; Poland
- ⁸⁷ Faculty of Science, Kyoto University, Kyoto; Japan
- ⁸⁸ Kyoto University of Education, Kyoto; Japan
- ⁸⁹ Research Center for Advanced Particle Physics and Department of Physics, Kyushu University, Fukuoka ; Japan
- 90 Instituto de Física La Plata, Universidad Nacional de La Plata and CONICET, La Plata; Argentina
- 91 Physics Department, Lancaster University, Lancaster; United Kingdom
- ⁹² Oliver Lodge Laboratory, University of Liverpool, Liverpool; United Kingdom
- ⁹³ Department of Experimental Particle Physics, Jožef Stefan Institute and Department of Physics, University of Ljubljana, Ljubljana; Slovenia
- ⁹⁴ School of Physics and Astronomy, Queen Mary University of London, London; United Kingdom
- 95 Department of Physics, Royal Holloway University of London, Egham; United Kingdom
- ⁹⁶ Department of Physics and Astronomy, University College London, London; United Kingdom
- ⁹⁷ Louisiana Tech University, Ruston LA; United States of America
- ⁹⁸ Fysiska institutionen, Lunds universitet, Lund; Sweden
- 99 Departamento de Física Teorica C-15 and CIAFF, Universidad Autónoma de Madrid, Madrid; Spain
- ¹⁰⁰ Institut für Physik, Universität Mainz, Mainz; Germany

- ¹⁰¹ School of Physics and Astronomy, University of Manchester, Manchester; United Kingdom
- ¹⁰² CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille; France
- ¹⁰³ Department of Physics, University of Massachusetts, Amherst MA; United States of America
- ¹⁰⁴ Department of Physics, McGill University, Montreal QC; Canada
- 105 School of Physics, University of Melbourne, Victoria; Australia
- 106 Department of Physics, University of Michigan, Ann Arbor MI; United States of America
- 107 Department of Physics and Astronomy, Michigan State University, East Lansing MI; United States of America
- ¹⁰⁸ Group of Particle Physics, University of Montreal, Montreal QC; Canada
- ¹⁰⁹ Fakultät für Physik, Ludwig-Maximilians-Universität München, München; Germany
- $^{110}\ Max-Planck-Institut\ f\"{u}r\ Physik\ (Werner-Heisenberg-Institut),\ M\"{u}nchen;\ Germany$
- 111 Graduate School of Science and Kobayashi-Maskawa Institute, Nagoya University, Nagoya; Japan
- ¹¹² Department of Physics and Astronomy, University of New Mexico, Albuquerque NM; United States of America
- ¹¹³ Institute for Mathematics, Astrophysics and Particle Physics, Radboud University/Nikhef, Nijmegen; Netherlands
- ¹¹⁴ Nikhef National Institute for Subatomic Physics and University of Amsterdam, Amsterdam; Netherlands
- ¹¹⁵ Department of Physics, Northern Illinois University, DeKalb IL; United States of America
- ¹¹⁶ (a) New York University Abu Dhabi, Abu Dhabi; (b) United Arab Emirates University, Al Ain; (c) University of Sharjah, Sharjah; United Arab Emirates
- ¹¹⁷ Department of Physics, New York University, New York NY; United States of America
- ¹¹⁸ Ochanomizu University, Otsuka, Bunkyo-ku, Tokyo; Japan
- ¹¹⁹ Ohio State University, Columbus OH; United States of America
- ¹²⁰ Homer L. Dodge Department of Physics and Astronomy, University of Oklahoma, Norman OK; United States of America
- ¹²¹ Department of Physics, Oklahoma State University, Stillwater OK; United States of America
- ¹²² Palacký University, Joint Laboratory of Optics, Olomouc; Czech Republic
- ¹²³ Institute for Fundamental Science, University of Oregon, Eugene, OR; United States of America
- ¹²⁴ Graduate School of Science, Osaka University, Osaka; Japan
- ¹²⁵ Department of Physics, University of Oslo, Oslo; Norway
- ¹²⁶ Department of Physics, Oxford University, Oxford; United Kingdom
- ¹²⁷ LPNHE, Sorbonne Université, Université Paris Cité, CNRS/IN2P3, Paris; France
- ¹²⁸ Department of Physics, University of Pennsylvania, Philadelphia PA; United States of America
- ¹²⁹ Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh PA; United States of America
- ¹³⁰ (a) Laboratório de Instrumentação e Física Experimental de Partículas LIP, Lisboa; (b) Departamento de Física, Faculdade de Ciências, Universidade de Lisboa, Lisboa; (c) Departamento de Física, Universidade de Coimbra, Coimbra; (d) Centro de Física Nuclear da Universidade de Lisboa, Lisboa; (e) Departamento de Física, Universidade do Minho, Braga; (f) Departamento de Física Teórica y del Cosmos, Universidad de Granada, Granada (Spain); (g) Departamento de Física, Instituto Superior Técnico, Universidade de Lisboa, Lisboa; Portugal
- ¹³¹ Institute of Physics of the Czech Academy of Sciences, Prague; Czech Republic
- ¹³² Czech Technical University in Prague, Prague; Czech Republic
- ¹³³ Charles University, Faculty of Mathematics and Physics, Prague; Czech Republic
- ¹³⁴ Particle Physics Department, Rutherford Appleton Laboratory, Didcot; United Kingdom
- ¹³⁵ IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette; France
- ¹³⁶ Santa Cruz Institute for Particle Physics, University of California Santa Cruz, Santa Cruz CA; United States of America
- ¹³⁷ (a) Departamento de Física, Pontificia Universidad Católica de Chile, Santiago; (b) Millennium Institute for Subatomic physics at high energy frontier (SAPHIR), Santiago; (c) Instituto de Investigación Multidisciplinario en Ciencia y Tecnología, y Departamento de Física, Universidad de La Serena; (d) Universidad Andres Bello, Departament of Physics, Santiago; (e) Instituto de Alta Investigación,

- $\label{thm:controller} Universidad\ de\ Tarapac\'a,\ Arica; {}^{(f)}Departamento\ de\ F\'isica,\ Universidad\ T\'ecnica\ Federico\ Santa\ Mar\'ia,\ Valpara\'iso;\ Chile$
- ¹³⁸ Department of Physics, University of Washington, Seattle WA; United States of America
- ¹³⁹ Department of Physics and Astronomy, University of Sheffield, Sheffield; United Kingdom
- ¹⁴⁰ Department of Physics, Shinshu University, Nagano; Japan
- ¹⁴¹ Department Physik, Universität Siegen, Siegen; Germany
- ¹⁴² Department of Physics, Simon Fraser University, Burnaby BC; Canada
- ¹⁴³ SLAC National Accelerator Laboratory, Stanford CA; United States of America
- ¹⁴⁴ Department of Physics, Royal Institute of Technology, Stockholm; Sweden
- ¹⁴⁵ Departments of Physics and Astronomy, Stony Brook University, Stony Brook NY; United States of America
- ¹⁴⁶ Department of Physics and Astronomy, University of Sussex, Brighton; United Kingdom
- ¹⁴⁷ School of Physics, University of Sydney, Sydney; Australia
- ¹⁴⁸ Institute of Physics, Academia Sinica, Taipei; Taiwan
- ¹⁴⁹ (a) E. Andronikashvili Institute of Physics, Iv. Javakhishvili Tbilisi State University, Tbilisi; (b) High Energy Physics Institute, Tbilisi State University, Tbilisi; (c) University of Georgia, Tbilisi; Georgia
- ¹⁵⁰ Department of Physics, Technion, Israel Institute of Technology, Haifa; Israel
- ¹⁵¹ Raymond and Beverly Sackler School of Physics and Astronomy, Tel Aviv University, Tel Aviv; Israel
- ¹⁵² Department of Physics, Aristotle University of Thessaloniki, Thessaloniki; Greece
- ¹⁵³ International Center for Elementary Particle Physics and Department of Physics, University of Tokyo, Tokyo; Japan
- ¹⁵⁴ Department of Physics, Tokyo Institute of Technology, Tokyo; Japan
- ¹⁵⁵ Department of Physics, University of Toronto, Toronto ON; Canada
- ¹⁵⁶ (a) TRIUMF, Vancouver BC; (b) Department of Physics and Astronomy, York University, Toronto ON; Canada
- ¹⁵⁷ Division of Physics and Tomonaga Center for the History of the Universe, Faculty of Pure and Applied Sciences, University of Tsukuba, Tsukuba; Japan
- ¹⁵⁸ Department of Physics and Astronomy, Tufts University, Medford MA; United States of America
- 159 Department of Physics and Astronomy, University of California Irvine, Irvine CA; United States of America
- ¹⁶⁰ Department of Physics and Astronomy, University of Uppsala, Uppsala; Sweden
- ¹⁶¹ Department of Physics, University of Illinois, Urbana IL; United States of America
- ¹⁶² Instituto de Física Corpuscular (IFIC), Centro Mixto Universidad de Valencia CSIC, Valencia; Spain
- ¹⁶³ Department of Physics, University of British Columbia, Vancouver BC; Canada
- ¹⁶⁴ Department of Physics and Astronomy, University of Victoria, Victoria BC; Canada
- ¹⁶⁵ Fakultät für Physik und Astronomie, Julius-Maximilians-Universität Würzburg, Würzburg; Germany
- ¹⁶⁶ Department of Physics, University of Warwick, Coventry; United Kingdom
- 167 Waseda University, Tokyo; Japan
- ¹⁶⁸ Department of Particle Physics and Astrophysics, Weizmann Institute of Science, Rehovot; Israel
- ¹⁶⁹ Department of Physics, University of Wisconsin, Madison WI; United States of America
- ¹⁷⁰ Fakultät für Mathematik und Naturwissenschaften, Fachgruppe Physik, Bergische Universität Wuppertal, Wuppertal; Germany
- ¹⁷¹ Department of Physics, Yale University, New Haven CT; United States of America
 - a Also Affiliated with an institute covered by a cooperation agreement with CERN
- ^b Also at Borough of Manhattan Community College, City University of New York, New York NY; United States of America
- ^c Also at Bruno Kessler Foundation, Trento; Italy
- ^d Also at Center for High Energy Physics, Peking University; China
- ^e Also at Center for Interdisciplinary Research and Innovation (CIRI-AUTH), Thessaloniki; Greece
- ^f Also at Centro Studi e Ricerche Enrico Fermi; Italy
- ^g Also at CERN, Geneva; Switzerland

- h Also at Département de Physique Nucléaire et Corpusculaire, Université de Genève, Genève; Switzerland
- ⁱ Also at Departament de Fisica de la Universitat Autonoma de Barcelona, Barcelona; Spain
- ^j Also at Department of Financial and Management Engineering, University of the Aegean, Chios; Greece
- ^k Also at Department of Physics and Astronomy, Michigan State University, East Lansing MI; United States of America
- ^l Also at Department of Physics, Ben Gurion University of the Negev, Beer Sheva; Israel
- ^m Also at Department of Physics, California State University, East Bay; United States of America
- ⁿ Also at Department of Physics, California State University, Sacramento; United States of America
- $^o\ Also\ at\ Department\ of\ Physics,\ King's\ College\ London,\ London;\ United\ Kingdom$
- ^p Also at Department of Physics, University of Fribourg, Fribourg; Switzerland
- ^q Also at Department of Physics, University of Thessaly; Greece
- ^r Also at Department of Physics, Westmont College, Santa Barbara; United States of America
- ^s Also at Hellenic Open University, Patras; Greece
- ^t Also at Institucio Catalana de Recerca i Estudis Avancats, ICREA, Barcelona; Spain
- ^u Also at Institut für Experimentalphysik, Universität Hamburg, Hamburg; Germany
- v Also at Institute of Applied Physics, Mohammed VI Polytechnic University, Ben Guerir; Morocco
- ^w Also at Institute of Particle Physics (IPP); Canada
- ^x Also at Institute of Physics and Technology, Ulaanbaatar; Mongolia
- ^y Also at Institute of Physics, Azerbaijan Academy of Sciences, Baku; Azerbaijan
- ^z Also at Institute of Theoretical Physics, Ilia State University, Tbilisi; Georgia
- ^{aa} Also at Lawrence Livermore National Laboratory, Livermore; United States of America
- ^{ab} Also at The Collaborative Innovation Center of Quantum Matter (CICQM), Beijing; China
- ac Also at TRIUMF, Vancouver BC; Canada
- ^{ad} Also at Università di Napoli Parthenope, Napoli; Italy
- ae Also at University of Chinese Academy of Sciences (UCAS), Beijing; China
- af Also at University of Colorado Boulder, Department of Physics, Colorado; United States of America
- ^{ag} Also at Washington College, Maryland; United States of America
- ah Also at Physics Department, An-Najah National University, Nablus, Palestine
- ai Also at L2IT, Université de Toulouse, CNRS/IN2P3, UPS, Toulouse; France
- ^{aj} Also at National Institute of Physics, University of the Philippines Diliman (Philippines); Philippines
- ^{ak} Also at Technical University of Munich, Munich; Germany
- al Also at Yeditepe University, Physics Department, Istanbul; Türkiye
- ^{am} Also at Department of Physics, Stanford University, Stanford CA, United States of America
 - $^*\ Deceased$