


BMP readouts vs non_neural_ectoderm (expected high) and neural_plate (expected low), 75% epiboly - 6-somite: 88% of non_neural_ectoderm cells and 93% of neural_plate cells as expected
(high / low: smoothed readout score above / below the library average; readout genes removed from the programs)




Nodal readouts vs margin (expected high) and presumptive_neural (expected low), shield - bud: 43% of margin cells and 84% of presumptive_neural cells as expected
(high / low: smoothed readout score above / below the library average; readout genes removed from the programs)



Wnt readouts vs tailbud (expected high) and forebrain_eye (expected low), bud - 6-somite: 82% of tailbud cells and 99% of forebrain_eye cells as expected
(high / low: smoothed readout score above / below the library average; readout genes removed from the programs)



FGF readouts vs mesoderm (expected high) and non_neural_ectoderm (expected low), 75% epiboly - bud: 54% of mesoderm cells and 100% of non_neural_ectoderm cells as expected
(high / low: smoothed readout score above / below the library average; readout genes removed from the programs)



Notch readouts vs presomitic_mesoderm (expected high) and lateral_plate (expected low), bud - 6-somite: 65% of presomitic_mesoderm cells and 94% of lateral_plate cells as expected
(high / low: smoothed readout score above / below the library average; readout genes removed from the programs)

