


signal:BMP readouts vs non_neural_ectoderm (expected high) and neural_plate (expected low), high - 6-somite: 91% of non_neural_ectoderm cells and 81% of neural_plate cells as expected
(high / low: smoothed readout score above / below the library average; readout genes removed from the programs)




signal:Nodal readouts vs margin (expected high) and neural_plate (expected low), high - 6-somite: 40% of margin cells and 88% of neural_plate cells as expected
(high / low: smoothed readout score above / below the library average; readout genes removed from the programs)



signal:Wnt readouts vs tailbud (expected high) and forebrain_eye (expected low), high - 6-somite: 88% of tailbud cells and 97% of forebrain_eye cells as expected
(high / low: smoothed readout score above / below the library average; readout genes removed from the programs)



signal:FGF readouts vs mesoderm (expected high) and non_neural_ectoderm (expected low), high - 6-somite: 64% of mesoderm cells and 97% of non_neural_ectoderm cells as expected
(high / low: smoothed readout score above / below the library average; readout genes removed from the programs)



receiving:Notch readouts vs presomitic_mesoderm (expected high) and lateral_plate (expected low), dome - 6-somite: 42% of presomitic_mesoderm cells and 85% of lateral_plate cells as expected
(high / low: smoothed readout score above / below the library average; readout genes removed from the programs)

