Proxy coherence per library (* p < 0.05); grey = gene set not scorable/absent

Nodal readout [coherent]
Nodal secondary [not coherent]
Nodal sender [coherent]
Nodal receiver [not coherent]
BMP readout [coherent]
BMP secondary [not coherent]
BMP sender [coherent]
BMP receiver [not coherent]
BMP antagonist [coherent]
Wnt readout [coherent]
Wnt secondary [not coherent]
Wnt sender [not scorable (<2 usable genes)]
Wnt receiver [not coherent]
FGF readout [coherent]
FGF secondary [coherent]
FGF sender [coherent]
FGF receiver [not coherent]
Hh readout [not scorable (<2 usable genes)]
Hh secondary [coherent]
Hh sender [not scorable (<2 usable genes)]
Hh receiver [not scorable (<2 usable genes)]
Notch readout [coherent]
Notch secondary [not scorable (<2 usable genes)]
Notch sender [not scorable (<2 usable genes)]
Notch receiver [not scorable (<2 usable genes)]
VEGF readout [not scorable (<2 usable genes)]
VEGF secondary [not scorable (<2 usable genes)]
VEGF sender [not scorable (<2 usable genes)]
VEGF receiver [not scorable (<2 usable genes)]

high DS5
high DS5b
oblong DS5
oblong DS5b
dome DS5
30% DS5
30% DS5b
shield DS5
50% DS2
50% DS3
50% DS4
50% DS4b

-log10 empirical p (coherence vs matched random sets)
3.0
2.5
2.0
1.5
1.0
0.5
0.0