

# An exploratory analysis of patient episode data

Henry Wilde

Cardiff University School of Mathematics

April 30, 2018

# Outline

## Motivation

### Getting to know the data

Structure and  
origin

Summative  
analysis

Measuring  
variation

### Slice analysis

General  
methodology

Diabetic patient  
analysis

### Moving forward

## 1 Motivation

## 2 Getting to know the data

Structure and origin

Summative analysis

Measuring variation

## 3 Slice analysis

General methodology

Diabetic patient analysis

## 4 Moving forward

# Motivation

## Motivation

### Getting to know the data

Structure and  
origin

Summative  
analysis

Measuring  
variation

### Slice analysis

General  
methodology

Diabetic patient  
analysis

### Moving forward

- Observe and understand cost variation
- Identify important slices in the data
- Develop methods for examining slices of the data
- Analyse their impact on costs and comorbidity

# Structure and origin

# Structure and origin

All data is provided by the Cwm Taf University Health Board.

## Motivation

### Getting to know the data

#### Structure and origin

Summative  
analysis

Measuring  
variation

### Slice analysis

General  
methodology  
Diabetic patient  
analysis

### Moving forward

# Structure and origin

## Motivation

## Getting to know the data

### Structure and origin

Summative  
analysis

Measuring  
variation

## Slice analysis

General  
methodology

Diabetic patient  
analysis

## Moving forward

All data is provided by the Cwm Taf University Health Board.

We have:

- 2,447,299 patient episodes
- 1,946,545 patient spells
- 865,421 individual patients

# Structure and origin

All data is provided by the Cwm Taf University Health Board.

We have:

- 2,447,299 patient episodes
- 1,946,545 patient spells
- 865,421 individual patients

Each row is made up of roughly 250 attributes, including:

# Structure and origin

All data is provided by the Cwm Taf University Health Board.

We have:

- 2,447,299 patient episodes
- 1,946,545 patient spells
- 865,421 individual patients

Each row is made up of roughly 250 attributes, including:

- personal identifiers and demographic information



# Structure and origin

## Motivation

## Getting to know the data

### Structure and origin

### Summative analysis

### Measuring variation

## Slice analysis

### General methodology Diabetic patient analysis

## Moving forward

All data is provided by the Cwm Taf University Health Board.

We have:

- 2,447,299 patient episodes
- 1,946,545 patient spells
- 865,421 individual patients

Each row is made up of roughly 250 attributes, including:

- personal identifiers and demographic information
- condition and treatment indicators (HRG, OPCS4, ICD10)

# Structure and origin

## Motivation

## Getting to know the data

### Structure and origin

### Summative analysis

### Measuring variation

## Slice analysis

### General methodology Diabetic patient analysis

## Moving forward

All data is provided by the Cwm Taf University Health Board.

We have:

- 2,447,299 patient episodes
- 1,946,545 patient spells
- 865,421 individual patients

Each row is made up of roughly 250 attributes, including:

- personal identifiers and demographic information
- condition and treatment indicators (HRG, OPCS4, ICD10)
- other clinical references

# Structure and origin

## Motivation

## Getting to know the data

### Structure and origin

### Summative analysis

### Measuring variation

## Slice analysis

### General methodology Diabetic patient analysis

## Moving forward

All data is provided by the Cwm Taf University Health Board.

We have:

- 2,447,299 patient episodes
- 1,946,545 patient spells
- 865,421 individual patients

Each row is made up of roughly 250 attributes, including:

- personal identifiers and demographic information
- condition and treatment indicators (HRG, OPCS4, ICD10)
- other clinical references
- cost components

# Summative analysis

# Summative analysis

- Our data is skewed towards low-cost, short-stay episodes

# Summative analysis

- Our data is skewed towards low-cost, short-stay episodes
- This extends to the spell level with largely one or two-time visits from patients

# Summative analysis

## Motivation

### Getting to know the data

Structure and  
origin

**Summative  
analysis**

Measuring  
variation

### Slice analysis



General  
methodology

Diabetic patient  
analysis

### Moving forward



- Our data is skewed towards low-cost, short-stay episodes
- This extends to the spell level with largely one or two-time visits from patients
- Long and heavy tails are present in our costs and lengths of stay

# Number of spells





# Length of stay (spell-wise)



# Net cost

## Motivation

### Getting to know the data

Structure and  
origin


### Summative analysis

Measuring  
variation

### Slice analysis

General  
methodology  
Diabetic patient  
analysis

### Moving forward



# Summative analysis

Other areas of interest to us are:

- Demographic variables
- Interactions between variables

# Demographic analysis

As it stands, demographic information is not well-recorded in the data.

# Demographic analysis

As it stands, demographic information is not well-recorded in the data.



- Gender is strictly binary and not recorded for all patients or episodes

# Demographic analysis

As it stands, demographic information is not well-recorded in the data.

- Gender is strictly binary and not recorded for all patients or episodes
- Limited geographic information is encoded in the GP practice code of the patient

# Demographic analysis



# Correlation

Motivation

Getting to know the data

Structure and origin

Summative analysis

Measuring variation


Slice analysis

General methodology

Diabetic patient analysis

Moving forward

Correlation coefficients for spell-level cost components and other clinical variables





# Measuring variation

Variance is not scale invariant and did lead to misconceptions.

# Measuring variation

Variance is not scale invariant and did lead to misconceptions.

## Definition

Let  $\mu, \sigma^2$  denote the population mean and population variance of some population respectively. Then we define the *coefficient of variation*, denoted by  $C_v$ , to be:

$$C_v = \frac{\sigma}{\mu}$$

# Measuring variation

Variance is not scale invariant and did lead to misconceptions.


## Definition

Let  $\mu, \sigma^2$  denote the population mean and population variance of some population respectively. Then we define the *coefficient of variation*, denoted by  $C_v$ , to be:


$$C_v = \frac{\sigma}{\mu}$$

The coefficient of variation is scale invariant, and allows us to see the relative variation in each of our cost components.

# Measuring variation



# Cost component contribution



# General methods for slice analysis

## Motivation

### Getting to know the data

Structure and  
origin

Summative  
analysis

Measuring  
variation

### Slice analysis

General  
methodology

Diabetic patient  
analysis

### Moving forward

Given some slice of the data, we want to:

- Examine cost variations and general surface-level statistics
- Determine components and variable relationships of interest
- Consider the relative 'cost' of the patients in this slice
- Contrast this against its complement and the general dataset



# Diabetic patient analysis

This is a known area of interest to the health board.

Diabetic patients make up 10.8% of all the episodes in the dataset, and roughly 8.7% of the unique patients in the dataset.



Here we consider patients to be 'diabetic' if they have diabetes flagged as either a primary or secondary condition in their episode.

# Summative analysis








# Summative analysis



# Summative analysis



# Demographic analysis



# Cost variation

Motivation


Getting to  
know the data

- Structure and origin
- Summative analysis
- Measuring variation


Slice analysis

- General methodology
- Diabetic patient analysis

Moving  
forward



# Cost component contribution



# Correlation

Motivation  
Getting to know the data

Structure and origin  
Summative analysis  
Measuring variation

Slice analysis  
General methodology  
Diabetic patient analysis

Moving forward



# Correlation (differences)

Motivation

Getting to know the data


Structure and origin  
Summative analysis  
Measuring variation

Slice analysis

General methodology  
Diabetic patient analysis

Moving forward

Difference in correlation coefficients for diabetic patients and the general population



# Diabetic patient analysis

We will focus our definition of system 'cost' on three measures:



# Diabetic patient analysis

We will focus our definition of system 'cost' on three measures:

- Proportion of total daily admissions
- Average length of stay given admission date
- Proportion of net costs spent given admission date

# Diabetic patient analysis

We will focus our definition of system 'cost' on three measures:


- Proportion of total daily admissions
- Average length of stay given admission date
- Proportion of net costs spent given admission date

These are indicators of resources used and resources necessary.


This grouping by admission date will lead to a degree of misrepresentation in our plots.

Allows us to investigate patterns developing over time.


# Resource consumption



# Resource consumption



# Resource consumption



# Conclusions

- Relative resource consumption by diabetic patients is consistent
- Cost components are less variant than - and are comparable in their distribution to - non-diabetic patients

## Motivation

### Getting to know the data

Structure and  
origin

Summative  
analysis

Measuring  
variation

### Slice analysis

General  
methodology

Diabetic patient  
analysis

## Moving forward

# Moving forward

# Moving forward

- Resource consumption metric



# Moving forward

- Resource consumption metric
- Perform clustering analysis to find inherent slices in the data

# Moving forward

- Resource consumption metric
- Perform clustering analysis to find inherent slices in the data
- Incorporate external data
  - Decode GP practice codes for GeoPandas
  - Socio-economic analysis based on deprivation and geography
  - Temperature-based analysis

# Moving forward

- Resource consumption metric
- Perform clustering analysis to find inherent slices in the data
- Incorporate external data
  - Decode GP practice codes for GeoPandas
  - Socio-economic analysis based on deprivation and geography
  - Temperature-based analysis
- Severity and comorbidity analysis
  - Average severity of secondary conditions given some primary condition
  - Using the comorbidity index as a class label in some predictive analysis