

NOTES TITLE

Author Name

June 3, 2026

Abstract

These notes develop a small piece of theoretical machinery and work through its consequences in detail. The exposition is kept self-contained, with definitions, statements, short proofs, and worked examples interleaved with prose. The closing sections collect a few exercises for practice.

Contents

1	Setting	3
2	Definitions and Statements	3
2.1	Time Evolution	3
3	Examples	3
4	Exercises	4
5	Closing	5

1 Setting

These notes develop a small piece of theoretical machinery to illustrate the structure of the template. The exposition follows the typical pattern of mathematical and physics notes: definitions, statements of results, short proofs, and worked examples interleaved with prose.

Citations are rendered in brackets, e.g. [1], and multiple sources combine and compress to ranges automatically [1, 2].

2 Definitions and Statements

Definition 2.1. Let $\mathcal{H}$ be a complex separable Hilbert space. A *state* is a unit vector $\psi \in \mathcal{H}$, considered up to a phase factor $e^{i\theta}$.

Definition 2.2. An *observable* is a self-adjoint linear operator $\hat{A}$ acting on $\mathcal{H}$.

The expectation value of an observable in a given state is defined by

$$\langle A \rangle_\psi = \langle \psi | \hat{A} | \psi \rangle. \quad (1)$$

Theorem 2.3. *Expectation values of self-adjoint operators are real.*

Proof. By definition, $\overline{\langle A \rangle_\psi} = \overline{\langle \psi | \hat{A} | \psi \rangle} = \langle \psi | \hat{A}^\dagger | \psi \rangle = \langle \psi | \hat{A} | \psi \rangle = \langle A \rangle_\psi$, where we used $\hat{A}^\dagger = \hat{A}$. $\square$

Remark 2.4. The reality of expectation values is what makes self-adjoint operators the natural mathematical model of observable quantities in quantum mechanics.

2.1 Time Evolution

Time evolution in quantum mechanics is governed by the Schrödinger equation:

$$i\hbar \frac{\partial \psi}{\partial t} = \hat{H} \psi, \quad (2)$$

where $\hat{H}$ is the Hamiltonian. For time-independent $\hat{H}$, equation (2) integrates to

$$\psi(t) = e^{-i\hat{H}t/\hbar} \psi(0). \quad (3)$$

2.1.1 Stationary states A *stationary state* is an eigenstate of $\hat{H}$; its time dependence is a pure phase, so every expectation value is constant in time. Run-in headings like this one mark the lowest structural level and are kept out of the table of contents.

3 Examples

Figure 1 shows a typical structure that recurs throughout the rest of these notes. A few fundamental constants are collected in Table 1 for reference.




Figure 1: Schematic energy level diagram.

Table 1: Fundamental physical constants.

Quantity	Symbol	Value
Speed of light	c	2.998×10^8 m/s
Planck's constant	$\hbar$	1.055×10^{-34} Js
Elementary charge	e	1.602×10^{-19} C

4 Exercises

Notes can carry problems alongside the exposition. The `exercise` environment is numbered per section and is visually separated from the surrounding text by `\separator`. Subproblems use the `subproblems` environment, which labels parts a), b), c) without affecting ordinary `enumerate` lists.

Exercise 4.1

Show that the eigenvalues of a self-adjoint operator $\hat{A}$ are real, and that eigenvectors belonging to distinct eigenvalues are orthogonal.

Exercise 4.2

Consider a particle in a one-dimensional infinite square well of width L .

- Write down the normalized energy eigenfunctions.
- Compute the energy levels E_n .

The remaining parts build on the spectrum found above; use the result from part (b) when sketching the densities.

- c) Sketch the probability density for $n = 1, 2, 3$.
- d) Compute the expectation value $\langle x \rangle$ in the ground state.

5 Closing

These notes are a template; the content above exists only to demonstrate typography. Replace it with your own material as the notes develop.

References

- [1] Firstname Lastname. Example article title. *Journal Name*, 12(3):45–67, 2025.
- [2] Firstname Lastname. *Example Book Title*. Publisher Name, City, 2025.