

# Homework 10

April 22, 2025

## Exercise 2.2.10

Let  $X$  be the quotient space of  $S^2$  under the identifications  $x \sim -x$  for  $x$  in the equator  $S^1$ . Compute the homology groups  $H_i(X)$ . Do the same for  $S^3$  with antipodal points of the equatorial  $S^2 \subset S^3$  identified.

## Proof.

For the first  $X$ , because we have 1 0-cell, 1 1-cell, and 2 2-cells, we get the following chain complex:

$$0 \xrightarrow{f} \mathbb{Z} \oplus \mathbb{Z} \xrightarrow{g} \mathbb{Z} \xrightarrow{h} \mathbb{Z} \xrightarrow{\phi} 0$$

So that our homology groups are:

$$H_0(X) \cong \ker(\phi) / \operatorname{im}(h)$$

$$H_1(X) \cong \ker(h) / \operatorname{im}(g)$$

$$H_2(X) \cong \ker(g) / \operatorname{im}(f)$$

with all other homology groups for  $n > 2$  zero.

Clearly,  $\ker(\phi) \cong \mathbb{Z}$ , and because  $h$  is a map from 1-cells to 0-cells,  $h$  is the zero map. This is also evident because the boundary of the 1-cell is zero, because the ends are identified. Thus  $\operatorname{im}(h) \cong 0$ , and  $\ker(h) \cong \mathbb{Z}$ .

For the  $g$  map, it's easiest to first think about what the map is doing. Because the equator has  $x \sim -x$ , the 2-cell "wraps around" the equator twice, for both the north and south 2-cells. Thus both generators of  $\mathbb{Z} \oplus \mathbb{Z}$ ,  $(1, 0)$  and  $(0, 1)$ , map to 2. Thus the image of  $g$  is the map generated by 2, so  $\operatorname{im}(g) \cong 2\mathbb{Z}$ . When considering the kernel, the elements are of the form  $a(1, 0) + b(0, 1) = 0$ , so that  $b = -a$ . Thus the kernel is wholly generated by  $(-1, 1)$ , so  $\ker(g) \cong \mathbb{Z}$ . Finally, we can clearly see that  $\operatorname{im}(f) \cong 0$ .

Thus we are left with:

$$H_0(X) \cong \ker(\phi) / \operatorname{im}(h) \cong \mathbb{Z}/0 \cong \mathbb{Z}$$

$$H_1(X) \cong \ker(h) / \operatorname{im}(g) \cong \mathbb{Z} / 2\mathbb{Z}$$

$$H_2(X) \cong \ker(g) / \operatorname{im}(f) \cong \mathbb{Z}/0 \cong \mathbb{Z}$$

Now letting  $X$  be the second space, we have a similar CW Complex structure: 1 0-cell, 1 1-cell, 1 2-cell, 2 3-cells. Thus we have the following chain complex:

$$0 \xrightarrow{f} \mathbb{Z} \oplus \mathbb{Z} \xrightarrow{g} \mathbb{Z} \xrightarrow{h} \mathbb{Z} \xrightarrow{\phi} \mathbb{Z} \xrightarrow{\psi} 0$$

So that our homology groups are:

$$H_0(X) \cong \ker(\psi) / \operatorname{im}(\phi)$$

$$H_1(X) \cong \ker(\phi) / \operatorname{im}(h)$$

$$H_2(X) \cong \ker(h) / \operatorname{im}(g)$$

$$H_3(X) \cong \ker(g) / \operatorname{im}(f)$$

with all other homology groups for  $n > 2$  zero.

We can use a similar logic as the last time to find everything but the map  $h$  (note that although we have increased dimension, the 3-cell still "wraps around" the  $S^2$  equator twice, similar to before).

So we have

$$H_0(X) \cong \ker(\psi) / \operatorname{im}(\phi) \cong \mathbb{Z}/0 \cong \mathbb{Z}$$

$$H_1(X) \cong \ker(\phi) / \operatorname{im}(h) \cong \mathbb{Z}/\operatorname{im}(h)$$

$$H_2(X) \cong \ker(h) / \operatorname{im}(g) \cong \ker(h)/2\mathbb{Z}$$

$$H_3(X) \cong \ker(g) / \operatorname{im}(f) \cong \mathbb{Z}/0 \cong \mathbb{Z}$$

For the map  $h$ , as it maps the 2-cell to the 1-cell, ■