

Columbia HW Problems

Hui Sun

August 5, 2025

Contents

1	HW2	3
2	HW3	4

Chapter 1

HW2

Problem 1.1 (1). Determine whether the following statements are true or false. Justify your answers.

- (a) Any subring of a field is an integral domain.
- (b) The ring $\mathbb{Z}/49\mathbb{Z}$ is an integral domain.
- (c) The direct product $F_1 \times F_2$ of two fields is a field.
- (d) An element ab of a ring R is invertible if and only if both a and b are invertible.
- (e) The ring $\mathbb{Z} \times \mathbb{Z}$ has exactly four idempotents.

Hint: First find all idempotents in the ring \mathbb{Z} . An idempotent is an element e such that $e^2 = e$.

Proof. (a) True (b) False (c) False, consider $(1, 0) \cdot (0, 1)$ (d) False Consider $5 + 5 = 1$ in $\mathbb{Z}/6\mathbb{Z}$ (e) True. \square

Problem 1.2 (6). 6. An element x of a ring R is called *nilpotent* if $x^n = 0$ for some $n > 0$. Note that $0 \in R$ is always nilpotent. (Remark: A nonzero nilpotent element is a zero divisor, while a zero divisor does not have to be nilpotent.)

- (a) (5 points) Show that 0 is the only nilpotent element of an integral domain R .
- (b) (5 points) Find all nilpotent elements in the following rings:

$$\mathbb{Z}, \quad \mathbb{Q}, \quad \mathbb{Z}/9\mathbb{Z}, \quad \mathbb{Z}/12\mathbb{Z}, \quad \mathbb{Q}[x].$$

- (c) (optional, 10 points) Show that if x, y are nilpotent then $x+y$ is nilpotent (assume that $x^n = 0$, $y^m = 0$, use that the ring is commutative and apply the binomial theorem from lecture 1 to some large power of $x + y$).

Proposition 1.1. If $a \in \mathbb{Z}/n\mathbb{Z}$ is nilpotent, then a is nilpotent in $\mathbb{Z}/d\mathbb{Z}$ for all divisors d of n . Moreover, units are not nilpotent.

(b) For $\mathbb{Z}/9\mathbb{Z}$: $\{0, 3, 6\}$ are nilpotents. For $\mathbb{Z}/12\mathbb{Z}$: $\{0, 6\}$ are nilpotents. (c) try $(x + y)^{m+n}$. \square

Proposition 1.2. An integral domain must have characteristic 0 or equal to some prime p ; however, there exists char prime ring that is not an integral domain: $\mathbb{Z}/p\mathbb{Z} \times \mathbb{Z}/p\mathbb{Z}$.

Chapter 2

HW3

Problem 2.1 (Q7). Let $e \in R$ be an idempotent (i.e., $e^2 = e$) in a commutative ring R .

- (a) Check that Re and $R(1 - e)$ are ideals of R and that their intersection $Re \cap R(1 - e) = 0$. Show that any element $a \in R$ has a unique presentation as a sum of an element in Re and an element in $R(1 - e)$.
- (b) Prove that Re is a ring, with identity e and addition and multiplication inherited from R . Likewise for $R(1 - e)$. (Since $1 - e$ is also an idempotent, you don't need to repeat your arguments twice.)
- (c) Show that the map $\phi : Re \times R(1 - e) \rightarrow R$ defined by $\phi(a, b) = a + b$ is an isomorphism of rings.

Proof. (a) Let $a = re = r'(1 - e)$, then

$$a = re = r' - r'e = r'e - r'e = 0$$

Clearly we have $a = ae + a(1 - e)$, and one can show this presentation is unique.

(b) True.

(c) It is surjective and injective both by part (a).

Proposition 2.1. Let $e \in R$ be a nontrivial idempotent, then viewing $Re, R(1 - e)$ as rings with identities $e, (1 - e)$, we can decompose R into a product of two rings:

$$R \cong Re \cong R(1 - e)$$

□