

Calc III Sections

Fall 2025

Hui Sun

November 11, 2025

Calc III-Week 12 (11/10-14)

next week, please focus on line integrals, the fundamental theorem of line integrals and its applications.

Definition 1 (path integral). Let $c : [a, b] \rightarrow \mathbb{R}^3$ be a path of C^1 and $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ is such that $f \circ c$ is continuous on $[a, b]$. The **path integral** of $f(x, y, z)$ along the path c is given by

$$\begin{aligned}\int_c f ds &= \int_a^b f(c(t)) \|c'(t)\| dt \\ &= \int_a^b f(x(t), y(t), z(t)) \|c'(t)\| dt\end{aligned}$$

Definition 2 (line integral). Let F be a vector field on \mathbb{R}^3 that is continuous on the C^1 path $c : [a, b] \rightarrow \mathbb{R}^3$, where $c(t) = (x(t), y(t), z(t))$. We define $\int_c F \cdot ds$, the **line integral** of F along c by the following

$$\begin{aligned}\int_c F \cdot ds &= \int_a^b F(c(t)) \cdot c'(t) dt \\ &= \int_a^b \left(F_1 \frac{dx}{dt} + F_2 \frac{dy}{dt} + F_3 \frac{dz}{dt} \right) dt \\ &:= \int_c F_1 dx + F_2 dy + F_3 dz\end{aligned}$$

the expression $F_1 dx + F_2 dy + F_3 dz$ is called the **differential form**.

Example 1 (work done along a path). The force field F on a particle moving along a path $c : [a, b] \rightarrow \mathbb{R}^3$ is given by

$$\text{work done by } F = \int_a^b F(c(t)) \cdot c'(t) dt$$

Definition 3 (reparametrization). Let $h : I \rightarrow I_1$ be a C^1 real-valued bijective function. Let $c : I_1 \rightarrow \mathbb{R}^3$ be a piecewise C^1 path. Then we call the composition

$$p = c \circ h : I \rightarrow \mathbb{R}^3$$

a **reparametrization** of c .

Example 2. Let $c : [0, 1] \rightarrow \mathbb{R}^3$ be a C^1 path, then consider $h : [0, 1] \rightarrow [0, 1]$, where $h(t) = 1 - t$. Then the path

$$c_{\text{op}} = c \circ h(t) = c(1 - t)$$

is the same path in the opposite direction.

Proposition 1 (reparametrization for path integrals). Let c be a C^1 path and c' be any reparametrization of c , and let f be a continuous function on the image of c , then

$$\int_c f(x, y, z) ds = \int_{c'} f(x, y, z) ds$$

Proposition 2 (reparametrization for line integrals). Let F be a vector field continuous on the C^1 path $c : [a, b] \rightarrow \mathbb{R}^3$, and let $c' : [a', b'] \rightarrow \mathbb{R}^3$ be a reparametrization of c . If c' is orientation-preserving, then

$$\int_{c'} F \cdot ds = \int_c F \cdot ds$$

If c' is orientation-reversing, then

$$\int_{c'} F \cdot ds = - \int_c F \cdot ds$$

Proposition 3 (fundamental theorem of line integrals). Suppose $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ is of C^1 and that $c : [a, b] \rightarrow \mathbb{R}^3$ is piecewise C^1 . Then

$$\int_c \nabla f \cdot ds = f(c(b)) - f(c(a))$$

Problem 1. Let $c : [a, b] \rightarrow \mathbb{R}^3$ be a C^1 path, find a reparametrization $\tilde{c} = c \circ h$, such that $\tilde{c} : [0, 1] \rightarrow \mathbb{R}^3$.

Problem 2. Suppose that $\nabla f(x, y, z) = (2x^2, y, -z)$, and that $f(0, 1, -1) = 2$. What is the value of $f(2, 1, 1)$?

Problem 3. Suppose path c has length l , and $\|F\| \leq M$. Prove that

$$\left| \int_c F \cdot ds \right| \leq Ml$$

(Hint: Cauchy-Schwarz.)

Problem 4 (Group work). Given vector field $F(x, y) = (1, 1)$, and the following heart is given by the parametrization:

$$c(t) = ((1 - \sin t) \cos t, (1 - \sin t) \sin t) \quad \text{where } 0 \leq t \leq 2\pi$$

Group 1: what is the work done by going up the top of the heart? ($0 \leq t \leq \pi$).

Group 2: what is the work done by going up the bottom of the heart? ($\pi \leq t \leq 2\pi$).

