

False Discovery Rates, A New Deal

Matthew Stephens

2014/2/24

Before we get started: Getting Organized

- Over ~10 years of working with graduate students + postdocs, I've noticed something.

Before we get started: Getting Organized

- Over ~10 years of working with graduate students + postdocs, I've noticed something.
- Organized researchers get more done (and better!).

Before we get started: Getting Organized

- Over ~10 years of working with graduate students + postdocs, I've noticed something.
- Organized researchers get more done (and better!).
- Many of them are more organized than I am!

Before we get started: Getting Organized

- Over ~10 years of working with graduate students + postdocs, I've noticed something.
- Organized researchers get more done (and better!).
- Many of them are more organized than I am!
- Thought: I should get organized; I should help others get organized.

So what can you do?

- Buy a notebook; bring it to meetings; make notes!

So what can you do?

- Buy a notebook; bring it to meetings; make notes!
- Come to meetings with a written agenda.

So what can you do?

- Buy a notebook; bring it to meetings; make notes!
- Come to meetings with a written agenda.
- While doing research, record what you did and what the outcome was.

So what can you do?

- Buy a notebook; bring it to meetings; make notes!
- Come to meetings with a written agenda.
- While doing research, record what you did and what the outcome was.
- Use version control git and internet repositories (bitbucket, github) to organize notes, code, etc.

So what can you do?

- Buy a notebook; bring it to meetings; make notes!
- Come to meetings with a written agenda.
- While doing research, record what you did and what the outcome was.
- Use version control git and internet repositories (bitbucket, github) to organize notes, code, etc.
- Use knitr to help make your research reproducible.

What are these repository things?

- A repository: a central place in which an aggregation of data is kept and maintained in an organized way (searcharticle.com)

What are these repository things?

- A repository: a central place in which an aggregation of data is kept and maintained in an organized way (searcharticle.com)
- Great for sharing material across multiple people (eg student and advisor!)

What are these repository things?

- A repository: a central place in which an aggregation of data is kept and maintained in an organized way (searcharticle.com)
- Great for sharing material across multiple people (eg student and advisor!)
- An amateur example: <http://github.com/stephens999/ash>

What is knitr?

- An R package

What is knitr?

- An R package
- A tool for literate programming

What is knitr?

- An R package
- A tool for literate programming
- Text, and R code are interleaved

What is knitr?

- An R package
- A tool for literate programming
- Text, and R code are interleaved
- When you compile the document, the code is run, and output inserted into the text.

What is knitr?

- An R package
- A tool for literate programming
- Text, and R code are interleaved
- When you compile the document, the code is run, and output inserted into the text.
- Great for writing reports, and keeping a track of what you did and what the result was!

What is knitr?

- An R package
- A tool for literate programming
- Text, and R code are interleaved
- When you compile the document, the code is run, and output inserted into the text.
- Great for writing reports, and keeping a track of what you did and what the result was!
- This talk was written with knitr (with RStudio)!

What is Reproducible Research?

- Principle: when publishing results of computational procedures, we should publish the code that produced the results.

What is Reproducible Research?

- Principle: when publishing results of computational procedures, we should publish the code that produced the results.
- “publishing figures or results without the complete software environment could be compared to a mathematician publishing an announcement of a mathematical theorem without giving the proof” (Buckheit and Donohoe)

What is Reproducible Research?

- Principle: when publishing results of computational procedures, we should publish the code that produced the results.
- “publishing figures or results without the complete software environment could be compared to a mathematician publishing an announcement of a mathematical theorem without giving the proof” (Buckheit and Donohoe)
- “an article about a computational result is advertising, not scholarship. The actual scholarship is the full software environment, code and data, that produced the result.” [Claerbout]

Why is reproducibility important?

- Not only because people are forgetful, error-prone, or dishonest!

Why is reproducibility important?

- Not only because people are forgetful, error-prone, or dishonest!
- Reproducing work is also the first step to extending it.

Why is reproducibility important?

- Not only because people are forgetful, error-prone, or dishonest!
- Reproducing work is also the first step to extending it.
- Helps communications among researchers (eg student + advisor).

Why is reproducibility important?

- Not only because people are forgetful, error-prone, or dishonest!
- Reproducing work is also the first step to extending it.
- Helps communications among researchers (eg student + advisor).
- If you do not publish code implementing your methods, your methods will likely go unused.

More on git, github, knitr, reproducibility

- Google “The git book”, to get started on git.

More on git, github, knitr, reproducibility

- Google “The git book”, to get started on git.
- Google “Karl Broman github tutorial” for statistics-oriented intro to github.

More on git, github, knitr, reproducibility

- Google “The git book”, to get started on git.
- Google “Karl Broman github tutorial” for statistics-oriented intro to github.
- Google “donohoe buckheit” for “Wavelab and reproducible research”

The Canonical Genomics Experiment

- Measure lots of things, with error

The Canonical Genomics Experiment

- Measure lots of things, with error
- Get estimates of effects β_j ($\hat{\beta}_j$) and their standard errors s_j

The Canonical Genomics Experiment

- Measure lots of things, with error
- Get estimates of effects β_j ($\hat{\beta}_j$) and their standard errors s_j
- Turn these into Z-scores, $z_j = \hat{\beta}_j/s_j$

The Canonical Genomics Experiment

- Measure lots of things, with error
- Get estimates of effects β_j ($\hat{\beta}_j$) and their standard errors s_j
- Turn these into Z-scores, $z_j = \hat{\beta}_j/s_j$
- Turn these into p values, p_j

The Canonical Genomics Experiment

- Measure lots of things, with error
- Get estimates of effects β_j ($\hat{\beta}_j$) and their standard errors s_j
- Turn these into Z-scores, $z_j = \hat{\beta}_j/s_j$
- Turn these into p values, p_j
- Apply `qvalue` to identify findings “significant” at a given False Discovery Rate.

The Canonical Genomics Experiment

- Measure lots of things, with error
- Get estimates of effects β_j ($\hat{\beta}_j$) and their standard errors s_j
- Turn these into Z-scores, $z_j = \hat{\beta}_j/s_j$
- Turn these into p values, p_j
- Apply `qvalue` to identify findings “significant” at a given False Discovery Rate.
- ...?

FDR, local fdr, and q values

Although precise definitions vary depending on whether one takes a Bayesian or Frequentist approach to the problem, roughly

- The FDR at a threshold P is

$$\text{FDR}(P) = \Pr(\beta_j = 0 | p_j < P).$$

FDR, local fdr, and q values

Although precise definitions vary depending on whether one takes a Bayesian or Frequentist approach to the problem, roughly

- The FDR at a threshold P is

$$\text{FDR}(P) = \Pr(\beta_j = 0 | p_j < P).$$

- The q value for observation j is $q_j = \text{FDR}(p_j)$.

FDR, local fdr, and q values

Although precise definitions vary depending on whether one takes a Bayesian or Frequentist approach to the problem, roughly

- The FDR at a threshold P is

$$\text{FDR}(P) = \Pr(\beta_j = 0 | p_j < P).$$

- The q value for observation j is $q_j = \text{FDR}(p_j)$.
- The local false discovery rate, fdr , at threshold P is

$$\text{fdr}(P) = \Pr(\beta_j = 0 | p_j = P).$$

FDR, local fdr, and q values

Although precise definitions vary depending on whether one takes a Bayesian or Frequentist approach to the problem, roughly

- The FDR at a threshold P is


$$\text{FDR}(P) = \Pr(\beta_j = 0 | p_j < P).$$

- The q value for observation j is $q_j = \text{FDR}(p_j)$.
- The local false discovery rate, fdr , at threshold P is

$$\text{fdr}(P) = \Pr(\beta_j = 0 | p_j = P).$$


- The fdr is more relevant, but slightly harder to estimate than FDR because it involves density estimation rather than tail-area estimation.

Example: FDR estimation




Example: FDR estimation

Distribution of p values




Data from Hedenfalk et al. comparing BRCA1 vs BRCA2 expression


Example: FDR estimation




Example: fdr estimation



Example: fdr estimation



Example: fdr estimation



FDR problem 1: different measurement precision

- If some effects are measured very imprecisely, those tests “lack power” and simply add noise

FDR problem 1: different measurement precision

- If some effects are measured very imprecisely, those tests “lack power” and simply add noise
- In particular, such tests increase the estimated number of nulls, and increase the FDR for other tests

FDR problem 1: different measurement precision


- If some effects are measured very imprecisely, those tests “lack power” and simply add noise
- In particular, such tests increase the estimated number of nulls, and increase the FDR for other tests
- It would seem preferable to simply ignore the tests with very low precision. Summarizing each test by a p value (or Z score) loses the information about precision.

Example: Mouse Heart Data


##	gene	lv1	lv2	rv1	rv2	genelength
## 1	Itm2a	2236	2174	9484	10883	1626
## 2	Sergef	97	90	341	408	1449
## 3	Fam109a	383	314	1864	2384	2331
## 4	Dhx9	2688	2631	18501	20879	4585
## 5	Ssu72	762	674	2806	3435	1446
## 8	Eif2b2	736	762	3081	3601	1565

- Data on 150 mouse hearts, dissected into left and right ventricle (courtesy Scott Schmemo, Marcelo Nobrega)


Example: Mouse Heart Data




Mouse Data: Counts vary considerably across genes




Lower count genes, less power



Higher count genes, more power



FDR problem 1: low count genes add noise, increase q values



FDR problem 1: Summary

- Analyzing p values or Z scores doesn't fully account for measurement precision.

Problem 2: The Zero Assumption (ZA)

- The standard q value approach assumes that all the p values near 1 are null.

Problem 2: The Zero Assumption (ZA)

- The standard q value approach assumes that all the p values near 1 are null.
- Analogously, one can assume that all Z scores near 0 are null. Efron refers to this as the “Zero Assumption”.

Problem 2: The Zero Assumption (ZA)

- The standard q value approach assumes that all the p values near 1 are null.
- Analogously, one can assume that all Z scores near 0 are null. Efron refers to this as the “Zero Assumption”.
- The ZA allows us to estimate the null proportion, π_0 , using the density of p values near 1 (or Z scores near 0).


Problem 2: The ZA

- The ZA seems initially natural.


Problem 2: The ZA

- The ZA seems initially natural.
- However, it turns out to imply unrealistic assumptions about the distribution of non-zero effects.


Implied distribution of p values under H_1




Implied distribution of Z scores under alternative (fdrtool)



Implied distribution of Z scores under alternative (locfdr)



Implied distribution of Z scores under alternative (mixfdr)



Problems: Summary

- By summarizing each observation by a Z score or p value, standard fdr tools ignore precision of different measurements

Problems: Summary

- By summarizing each observation by a Z score or p value, standard fdr tools ignore precision of different measurements
- Standard tools make the ZA, which implies actual effects have a (probably unrealistic) bimodal distribution. [and tends to overestimate π_0 , losing power]

Problems: Summary

- By summarizing each observation by a Z score or p value, standard fdr tools ignore precision of different measurements
- Standard tools make the ZA, which implies actual effects have a (probably unrealistic) bimodal distribution. [and tends to overestimate π_0 , losing power]
- Also standard tools focus only on zero vs non-zero effects. (eg what if we would like to identify genes that have at least a 2-fold change?)

FDR via Empirical Bayes

- Following previous work (e.g. Newton, Efron, Muralidharan) we take an empirical Bayes approach to FDR.

FDR via Empirical Bayes

- Following previous work (e.g. Newton, Efron, Muralidharan) we take an empirical Bayes approach to FDR.
- Eg Efron assumes that the Z scores come from a mixture of null, and alternative:

$$Z_j \sim f_Z(.) = \pi_0 N(., 0, 1) + (1 - \pi_0) f_1(.)$$

where f_1 is to be estimated from the data.

FDR via Empirical Bayes

- Following previous work (e.g. Newton, Efron, Muralidharan) we take an empirical Bayes approach to FDR.
- Eg Efron assumes that the Z scores come from a mixture of null, and alternative:

$$Z_j \sim f_Z(.) = \pi_0 N(., 0, 1) + (1 - \pi_0) f_1(.)$$

where f_1 is to be estimated from the data.

- Various semi-parametric approaches taken to estimating f_1 . For example, Efron uses Poisson regression; Muralidharan uses mixture of normal distributions.

FDR via Empirical Bayes

- Following previous work (e.g. Newton, Efron, Muralidharan) we take an empirical Bayes approach to FDR.
- Eg Efron assumes that the Z scores come from a mixture of null, and alternative:

$$Z_j \sim f_Z(.) = \pi_0 N(., 0, 1) + (1 - \pi_0) f_1(.)$$

where f_1 is to be estimated from the data.

- Various semi-parametric approaches taken to estimating f_1 . For example, Efron uses Poisson regression; Muralidharan uses mixture of normal distributions.
- $\text{fdr}(Z) \approx \pi_0 N(Z; 0, 1) / f_Z(Z)$

FDR: The New Deal

- Instead of modelling Z scores, model the effects β ,

$$\beta_j \sim \pi_0 \delta_0(.) + (1 - \pi_0)g(.)$$

FDR: The New Deal

- Instead of modelling Z scores, model the effects β ,

$$\beta_j \sim \pi_0 \delta_0(.) + (1 - \pi_0)g(.)$$

- Constrain g to be unimodal about 0; estimate g from data.

FDR: The New Deal

- Instead of modelling Z scores, model the effects β ,

$$\beta_j \sim \pi_0 \delta_0(.) + (1 - \pi_0)g(.)$$

- Constrain g to be unimodal about 0; estimate g from data.
- *Incorporate precision* of each observation $\hat{\beta}$ into the likelihood. Specifically, approximate likelihood for β_j by a normal:

$$L(\beta_j) \propto \exp(-0.5(\beta_j - \hat{\beta}_j)^2/s_j^2).$$

[From $\hat{\beta}_j \sim N(\beta_j, s_j)$]

FDR: The New Deal

- Instead of modelling Z scores, model the effects β ,

$$\beta_j \sim \pi_0 \delta_0(.) + (1 - \pi_0)g(.)$$

- Constrain g to be unimodal about 0; estimate g from data.
- *Incorporate precision* of each observation $\hat{\beta}$ into the likelihood. Specifically, approximate likelihood for β_j by a normal:

$$L(\beta_j) \propto \exp(-0.5(\beta_j - \hat{\beta}_j)^2/s_j^2).$$

[From $\hat{\beta}_j \sim N(\beta_j, s_j)$]

- fdr given by

$$p(\beta_j = 0|\hat{\beta}_j) = \pi_0 p(\hat{\beta}_j|\beta_j = 0)/p(\hat{\beta}_j)$$

FDR - A New Deal

- A convenient way to model g is by a mixture of 0-centered normal distributions:

$$g(\beta; \pi) = \sum_{k=1}^K \pi_k N(\beta; 0, \sigma_k^2)$$

FDR - A New Deal

- A convenient way to model g is by a mixture of 0-centered normal distributions:

$$g(\beta; \pi) = \sum_{k=1}^K \pi_k N(\beta; 0, \sigma_k^2)$$

- Estimating g comes down to estimating π . Joint estimation of π_0, π easy by maximum likelihood (EM algorithm) or variational Bayes.

FDR - A New Deal

- A convenient way to model g is by a mixture of 0-centered normal distributions:

$$g(\beta; \pi) = \sum_{k=1}^K \pi_k N(\beta; 0, \sigma_k^2)$$

- Estimating g comes down to estimating π . Joint estimation of π_0, π easy by maximum likelihood (EM algorithm) or variational Bayes.
- By allowing K large, and σ_k to span a dense grid of values, we get a fairly flexible unimodal symmetric distribution.

FDR - A New Deal

- A convenient way to model g is by a mixture of 0-centered normal distributions:

$$g(\beta; \pi) = \sum_{k=1}^K \pi_k N(\beta; 0, \sigma_k^2)$$

- Estimating g comes down to estimating π . Joint estimation of π_0, π easy by maximum likelihood (EM algorithm) or variational Bayes.
- By allowing K large, and σ_k to span a dense grid of values, we get a fairly flexible unimodal symmetric distribution.
- Can approximate, arbitrarily closely, any scale mixture of normals. Includes almost all priors used for sparse regression problems (spike-and-slab, double exponential/Laplace/Bayesian Lasso, horseshoe).

FDR - A New Deal

- Alternatively, a mixture of uniforms, with 0 as one end-point of the range, provides still more flexibility, and in particular allows for asymmetry.

FDR - A New Deal

- Alternatively, a mixture of uniforms, with 0 as one end-point of the range, provides still more flexibility, and in particular allows for asymmetry.
- If allow a very large number of uniforms this provides the non-parametric mle for g ; cf Grenander 1953; Campy + Thomas.

Illustration: g a mixture of 0-centered normals




Illustration: g a mixture of 0-centered normals




Illustration: g a mixture of 0-anchored uniforms





Illustration: g a mixture of 0-anchored uniforms



Issue: identifiability of π_0

- For estimating False Discoveries, we are asking whether $\beta_j = 0$.

Issue: identifiability of π_0

- For estimating False Discoveries, we are asking whether $\beta_j = 0$.
- However, the data cannot distinguish between $\beta_j = 0$ and β_j “very small”

Issue: identifiability of π_0

- For estimating False Discoveries, we are asking whether $\beta_j = 0$.
- However, the data cannot distinguish between $\beta_j = 0$ and β_j “very small”
- As a result π_0 is formally unidentifiable. Eg data can never rule out $\pi_0 = 0$.

Issue: identifiability of π_0

- The Zero assumption (ZA) solves the identifiability problem by assuming that there *are* no β_j near zero!

Issue: identifiability of π_0

- The Zero assumption (ZA) solves the identifiability problem by assuming that there *are* no β_j near zero!
- The ZA makes π_0 identifiable.

Issue: identifiability of π_0

- The Zero assumption (ZA) solves the identifiability problem by assuming that there *are* no β_j near zero!
- The ZA makes π_0 identifiable.
- Another view is that the estimate of π_0 under ZA will systematically tend to overestimate π_0 , and so is “conservative”.

Issue: identifiability of π_0

- The Zero assumption (ZA) solves the identifiability problem by assuming that there *are* no β_j near zero!
- The ZA makes π_0 identifiable.
- Another view is that the estimate of π_0 under ZA will systematically tend to overestimate π_0 , and so is “conservative”.
- That is it provides an “upper bound” on π_0

Identifiability of π_0 : Solution 1

- We replaced the ZA with the unimodal assumption on g .

Identifiability of π_0 : Solution 1

- We replaced the ZA with the unimodal assumption on g .
- This does not make π_0 identifiable, but it does effectively provide an upper bound on π_0 .


Identifiability of π_0 : Solution 1

- We replaced the ZA with the unimodal assumption on g .
- This does not make π_0 identifiable, but it does effectively provide an upper bound on π_0 .
- Indeed, we saw that when we estimated π_0 under the ZA the data then contradicted the unimodal assumption on g . Thus the upper bound is more conservative than under ZA.

Identifiability of π_0 : Solution 1


- We replaced the ZA with the unimodal assumption on g .
- This does not make π_0 identifiable, but it does effectively provide an upper bound on π_0 .
- Indeed, we saw that when we estimated π_0 under the ZA the data then contradicted the unimodal assumption on g . Thus the upper bound is more conservative than under ZA.
- In practice, implement upper bound by using penalized likelihood that encourages π_0 to be as big as possible.

Illustration: Simulated Example




Example: BRCA data

Recall Problem: distribution of alternative Z values multimodal



Problem Fixed: distribution of alternative Z values unimodal



BRCA1: Compare π_0 estimates

```
round(c(hh.fdrtool$param[3], hh.locfdr$fp0[1, 3], hh.mixfdr$pi0[1, 2])
```

```
## [1] 0.64 0.74 0.80 0.21
```

BRCA1: Compare number significant at $\text{fdr} < 0.05$

```
c(sum(hh.fdrtool$lfd < 0.05), sum(hh.locfdr$fdr < 0.05), sum(
  0.05), sum(hh.ashz$ZeroProb < 0.05))
```

```
## [1] 154 171 162 341
```

Identifiability of π_0 and the False Sign Rate

- Identifiability of π_0 is primarily an issue if we insist on asking question is $\beta_j = 0$?

Identifiability of π_0 and the False Sign Rate

- Identifiability of π_0 is primarily an issue if we insist on asking question is $\beta_j = 0$?
- How about we change focus: assume *none* of the β_j are zero (“one group approach”), and ask for which β_j are we confident about the sign (Gelman et al, 2012).

Identifiability of π_0 and the False Sign Rate

- Identifiability of π_0 is primarily an issue if we insist on asking question is $\beta_j = 0$?
- How about we change focus: assume *none* of the β_j are zero (“one group approach”), and ask for which β_j are we confident about the sign (Gelman et al, 2012).
- Positive and negative effects are often treated differently in practice anyway.


Identifiability of π_0 and the False Sign Rate

- Identifiability of π_0 is primarily an issue if we insist on asking question is $\beta_j = 0$?
- How about we change focus: assume *none* of the β_j are zero (“one group approach”), and ask for which β_j are we confident about the sign (Gelman et al, 2012).
- Positive and negative effects are often treated differently in practice anyway.
- That is we replace fdr with False Sign Rate (fsr), the probability that if we say an effect is positive (negative), it is not.


Identifiability of π_0 and the False Sign Rate

- Identifiability of π_0 is primarily an issue if we insist on asking question is $\beta_j = 0$?
- How about we change focus: assume *none* of the β_j are zero (“one group approach”), and ask for which β_j are we confident about the sign (Gelman et al, 2012).
- Positive and negative effects are often treated differently in practice anyway.
- That is we replace fdr with False Sign Rate (fsr), the probability that if we say an effect is positive (negative), it is not.
- Example: suppose we estimate that $\Pr(\beta_j < 0) = 0.975$ and $\Pr(\beta_j > 0) = 0.025$. Then we report β_j as a “(negative) discovery“, and estimate its fsr as 0.025.

The fdr is more robust than fdr



The fsr is more robust than fdr



Estimation and Shrinkage

- Besides allowing one to estimate fdr and fsr , this approach also provides a full posterior distribution for each β_j .

Estimation and Shrinkage

- Besides allowing one to estimate fdr and fsr , this approach also provides a full posterior distribution for each β_j .
- So for example we can easily compute fdrs for discoveries other than “non-zero” (eg compute $\Pr(\beta_j > 2|\hat{\beta}_j)$).

Estimation and Shrinkage

- Besides allowing one to estimate fdr and fsr , this approach also provides a full posterior distribution for each β_j .
- So for example we can easily compute fdrs for discoveries other than “non-zero” (eg compute $\Pr(\beta_j > 2|\hat{\beta}_j)$).
- And use it to obtain point estimates and credible intervals for each β_j , taking account of information from all the other β_j .

Estimation and Shrinkage

- Besides allowing one to estimate fdr and fsr , this approach also provides a full posterior distribution for each β_j .
- So for example we can easily compute fdrs for discoveries other than “non-zero” (eg compute $\Pr(\beta_j > 2|\hat{\beta}_j)$).
- And use it to obtain point estimates and credible intervals for each β_j , taking account of information from all the other β_j .
- Because $f(\beta)$ is unimodal, the point estimates will tend to be “shrunk” towards the overall mean (0).


Estimation and Shrinkage

- Besides allowing one to estimate fdr and fsr , this approach also provides a full posterior distribution for each β_j .
- So for example we can easily compute fdrs for discoveries other than “non-zero” (eg compute $\Pr(\beta_j > 2|\hat{\beta}_j)$).
- And use it to obtain point estimates and credible intervals for each β_j , taking account of information from all the other β_j .
- Because $f(\beta)$ is unimodal, the point estimates will tend to be “shrunk” towards the overall mean (0).
- Because $f(\beta)$ is estimated from the data, the amount of shrinkage is adaptive to the data. And because of the role of s_j , the amount of shrinkage adapts to the information on each gene.


Estimation and Shrinkage

- Besides allowing one to estimate fdr and fsr , this approach also provides a full posterior distribution for each β_j .
- So for example we can easily compute fdrs for discoveries other than “non-zero” (eg compute $\Pr(\beta_j > 2|\hat{\beta}_j)$).
- And use it to obtain point estimates and credible intervals for each β_j , taking account of information from all the other β_j .
- Because $f(\beta)$ is unimodal, the point estimates will tend to be “shrunk” towards the overall mean (0).
- Because $f(\beta)$ is estimated from the data, the amount of shrinkage is adaptive to the data. And because of the role of s_j , the amount of shrinkage adapts to the information on each gene.
- So we call the approach “Adaptive Shrinkage” (ASH).


Example: ASH applied to mouse data




Example: ASH applied to mouse data



Example: ASH applied to mouse data



Example: ASH applied to mouse data



Summary

- ASH provides a generic approach to shrinkage estimation, as well as false discovery (sign) rates.

Summary

- ASH provides a generic approach to shrinkage estimation, as well as false discovery (sign) rates.
- But by using two numbers ($\hat{\beta}$, s) instead of one (p values or z scores) precision of different measurements can be better accounted for.

Summary

- ASH provides a generic approach to shrinkage estimation, as well as false discovery (sign) rates.
- But by using two numbers ($\hat{\beta}$, s) instead of one (p values or z scores) precision of different measurements can be better accounted for.
- Unimodal assumption for effects reduces conservatism

Summary

- ASH provides a generic approach to shrinkage estimation, as well as false discovery (sign) rates.
- But by using two numbers ($\hat{\beta}$, s) instead of one (p values or z scores) precision of different measurements can be better accounted for.
- Unimodal assumption for effects reduces conservatism
- False Sign Rate is more robust to assumptions, and perhaps therefore preferable, than False Discovery Rate.

Other Applications

- Widely applicable: requiring only an estimated effect size and standard error for each object.

Other Applications

- Widely applicable: requiring only an estimated effect size and standard error for each object.
- Currently applying it to wavelet shrinkage applications.

Guarantees?

- “I think you have some nice ideas. How will you convince people to use them?” (C Morris)

Guarantees?

- “I think you have some nice ideas. How will you convince people to use them?” (C Morris)
- Theory anyone?

Next steps?

- Incorporate t likelihood as well as normal.

Next steps?

- Incorporate t likelihood as well as normal.
- Incorporate shrinkage of variances and not just means.

Next steps?

- Incorporate t likelihood as well as normal.
- Incorporate shrinkage of variances and not just means.
- Extend to allow $g(\cdot; \pi)$ to depend on covariates X .

Next steps?

- Incorporate t likelihood as well as normal.
- Incorporate shrinkage of variances and not just means.
- Extend to allow $g(\cdot; \pi)$ to depend on covariates X .
- Extend to allow for correlations in the measured $\hat{\beta}_j$.

Thanks

- to the several postdoctoral researchers and students who have worked with me on related topics.

Thanks

- to the several postdoctoral researchers and students who have worked with me on related topics.
- Including Scott Powers, Mengyin Lu, Tian Sen, Wei Wang, Zhengrong Xing.

Reproducible research

- This document is produced with **knitr**, **Rstudio** and **Pandoc**.

Reproducible research

- This document is produced with **knitr**, **Rstudio** and **Pandoc**.
- For more details see my `stephens999/ash` repository at <http://www.github.com/stephens999/ash>

Reproducible research

- This document is produced with **knitr**, **Rstudio** and **Pandoc**.
- For more details see my `stephens999/ash` repository at <http://www.github.com/stephens999/ash>
- Website: <http://stephenslab.uchicago.edu>

Pandoc Command used

```
pandoc -s -S -i --template=my.beamer -t beamer -V  
theme:CambridgeUS -V colortheme:beaver slides.md -o  
slides.pdf
```

(alternative to produce html slides; but figures would need reworking)

```
pandoc -s -S -i -t dzslides --mathjax slides.md -o  
slides.html
```

Here is my session info:

```
print(sessionInfo(), locale = FALSE)  
  
## R version 3.0.2 (2013-09-25)  
## Platform: x86_64-apple-darwin10.8.0 (64-bit)  
##  
## attached base packages:  
## [1] splines      parallel     stats        graphics     grDevices    utils  
## [8] methods     base
```

Some odd things in the data




Figure : plot of chunk unnamed-chunk-40

Error: incorrect number of dimensions

A technicality

- Suppose you estimate $\Pr(\beta_j < 0) = 0.98$, $\Pr(\beta_j > 0) = 0.01$,
 $\Pr(\beta_j = 0) = 0.01$.

A technicality


- Suppose you estimate $\Pr(\beta_j < 0) = 0.98$, $\Pr(\beta_j > 0) = 0.01$, $\Pr(\beta_j = 0) = 0.01$.
- Should you declare an fdr of 0.01 or 0.02?

A technicality


- Suppose you estimate $\Pr(\beta_j < 0) = 0.98$, $\Pr(\beta_j > 0) = 0.01$, $\Pr(\beta_j = 0) = 0.01$.
- Should you declare an fdr of 0.01 or 0.02?
- Maybe fsr makes more sense anyway?

Shrinkage is adaptive to information

Need to fix counts.associate to use fdr method in ash




Shrinkage is adaptive to information




Shrinkage is adaptive to information

##	gene	lv1	lv2	rv1	rv2	pval	zdat.ash\$lfr
## 19422	Mgat5b	7	10	320	452	0.03795	0
## 20432	Sec63	1042	1034	5496	6649	0.04908	0

Recall FDR problem 1: q values increased by low count genes



ASH q values more robust to inclusion of low count genes



Compare fitted $f(\beta)$, both estimating π_0 and fixing $\pi_0 = 0$.