


21. Sheu S-J, Ger L-P, Ho W-L. Late increased risk of retinal detachment after cataract extraction. *Am J Ophthalmol*. 2010;149(1):113–119.e1.
22. Tielsch JM, Legro MW, Cassard SD, et al. Risk factors for retinal detachment after cataract surgery: a population-based case-control study. *Ophthalmology*. 1996;103(10):1537–1545.
23. Jindachomthong KK, Cabral H, Subramanian ML, et al. Incidence and risk factors for delayed retinal tears after an acute, symptomatic posterior vitreous detachment. *Ophthalmol Retina*. 2023;7(4):318–324.
24. Ferrara M, Mehta A, Qureshi H, et al. Phenotype and outcomes of phakic versus pseudophakic primary rhegmatogenous retinal detachments: cataract or cataract surgery related? *Am J Ophthalmol*. 2021;222: 318–327.
25. Joseph DP, Ryan EH, Ryan CM, et al. Primary retinal detachment outcomes study: pseudophakic retinal detachment outcomes: primary retinal detachment outcomes study report number 3. *Ophthalmology*. 2020;127(11): 1507–1514.

Pictures & Perspectives



A Child with Posterior Ectopic Cilia

An otherwise healthy 9-year-old boy presented with irritation and tearing of the left eye for 6 months. There was no family history of a similar disorder, and the patient had no medical history of any ocular surgery or trauma. Slit-lamp examination showed a single black hair located in the palpebral conjunctiva of the upper lid (A). The lesion, including the hair and a tarsal plate tissue, was completely removed. Histopathologic examination showed posterior ectopic cilia (B and C). (Magnified version of Figure A-C is available online at www.aaojournal.org).

JIAQI GUAN, MD
 HUIJING YE, MD, PhD
 HUASHENG YANG, MD, PhD

State Key Laboratory of Ophthalmology, Zhongshan Ophthalmic Center, Sun Yat-sen University, Guangdong Provincial Key Laboratory of Ophthalmology and Visual Science, Guangzhou, China
