


22. Francisco BM, Salvador M, Amparo N. Oxidative stress in myopia. *Oxid Med Cell Longev*. 2015;2015: 750637.
23. Salzano AD, Khanal S, Cheung NL, et al. Repeated low-level red-light therapy: the next wave in myopia management? *Optom Vis Sci*. 2023;100(12):812–822.
24. Jonas JB, Spaide RF, Ostrin LA, et al. IMI-nonpathological human ocular tissue changes with axial myopia. *Invest Ophthalmol Vis Sci*. 2023;64(6):5.
25. Liu H, Yang Y, Guo J, et al. Retinal damage after repeated low-level red-light laser exposure. *JAMA Ophthalmol*. 2023;141(7):693–695.

Pictures & Perspectives



An Iris Cyst as a Presenting Feature of Axenfeld-Rieger Malformation

A 3-month-old girl presented with a whitish opacity in her right eye (OD). Examination revealed an abnormal iris morphology OD. She had a posterior embryotoxon (PE; **A**, black arrow), along with peripheral iris atrophy and holes (**A**, white arrow). On ultrasound biomicroscopy, multiple loculated cysts could be observed OD (**B**). Her left eye had a grossly normal iris morphology (**C**), although on gonioscopy a well-defined PE (**D**, black arrows), with high iris processes were observed (**D**, white arrowhead). The child had a known mutation (p.Gly34ThrfsTer8) that is associated with Axenfeld-Rieger malformation. (Magnified version of Figure **A-D** is available online at www.aaojournal.org).

ARNAV PANIGRAHI, MD¹

SHIKHA GUPTA, MD¹

VINEY GUPTA, MD¹

¹Dr Rajendra Prasad Centre for Ophthalmic Sciences, All India Institute of Medical Sciences, New Delhi, India