


Ophthalmic Images

Vascular Conjunctival Lesion in a Patient With a Recent Diagnosis of HIV Infection

Sofia De Arrigunaga, MD, MPH; Jason Fan, MD, PhD; Carol L. Karp, MD

A Slitlamp photography, conjunctival tumor



B Optical coherence tomography, subepithelial mass




Figure. A, Elevated vascular conjunctival tumor (white arrowheads) with hemorrhage (blue arrowheads). B, High-resolution optical coherence tomography reveals a subepithelial hyperreflective, highly cellular (arrowheads) mass with multiple vascular channels (asterisks). Inset, *Human herpesvirus 8* demonstrates positive nuclear staining within tumor cells (original magnification $\times 200$).

A young patient, diagnosed with HIV infection 5 months before presentation and taking highly active antiretroviral therapy (HAART), presented with a left eye lesion. At diagnosis, the CD4⁺ count was 69 cells/mm³. Examination revealed a highly vascular, supero-nasal heterogeneous mass with associated hemorrhage (Figure, A). High-resolution optical coherence tomography showed a subepithelial, hyperreflective, hypercellular tissue (Figure, B). The lesion was ex-

cised using the no-touch technique and cryotherapy. Histopathology revealed spindle-shaped cells positive for *Human herpesvirus 8* stain (Figure, B, inset). Conjunctival Kaposi sarcoma (KS) lesions are usually red, dark brown, or violaceous; flat or elevated; and may be associated with subconjunctival hemorrhage.^{1,2} Localized KS lesions can be treated with surgical excision.¹ In patients who are positive for HIV infection, HAART can be used as sole treatment³ or in combination with interferon injections⁴ or chemotherapeutic agents.⁵ Ocular KS lesions have become uncommon amid the widespread use of HAART.⁶

ARTICLE INFORMATION

Author Affiliations: Department of Ophthalmology, Bascom Palmer Eye Institute, University of Miami Miller School of Medicine, Miami, Florida.

Corresponding Author: Carol L. Karp, MD, Department of Ophthalmology, Bascom Palmer Eye Institute, University of Miami Miller School of Medicine, 900 NW 17th St, Miami, FL 33136 (ckarp@med.miami.edu).

Conflict of Interest Disclosures: Dr Karp reported receiving grants from the National Institutes of Health (P30EY014801) and Research to Prevent Blindness; institutional grants from Dr Ronald and Alicia Lepke grant, the Lee and Claire Hager grant, the Grant and Diana Stanton-Thornbrough grant, the Robert Baer Family grant, the Emilyn Page and Mark Feldberg grant, the Robert Farr Family grant, the Calvin and Flavia Oak Foundation, the Jose Ferreira de Melo grant, the Mr and Mrs Irwin Friedman grant, the Roberto and Antonia Menendez Family grant, the Stephen Takach grant, the Richard and Kathy Lesser grant, the Ragheb

Family grant, the Honorable A. Jay Cristol grant, the Michele and Ted Kaplan grant, the Christian Kathke grant, the Carol Soffer grant, and the Richard Azar Family grant; advisory board fees or stock options from Glaukos and Interfeen Biologics; and having a patent pending for PCT/US2022/029842. No other disclosures were reported.

REFERENCES

1. Venkateswaran N, Ramos JC, Cohen AK, et al. Spotlight on ocular Kaposi sarcoma: an update on the presentation, diagnosis, and management options. *Expert Rev Ophthalmol*. 2021;16(6):477-489. doi:10.1080/17469899.2021.1962294
2. Ulloa-Padilla JP, Ghassibi MP, Dubovy SR, Kerr DA. Clinicopathologic correlation of Kaposi sarcoma involving the ocular adnexa: immunophenotyping of diagnostic and therapeutic targets. *Ophthalmic Plast Reconstr Surg*. 2020;36(2):185-190. doi:10.1097/IOP.0000000000001506
3. Leder HA, Galor A, Peters GB, Kedhar SR, Dunn JP, Thorne JE. Resolution of conjunctival Kaposi

sarcoma after institution of highly active antiretroviral therapy alone. *Br J Ophthalmol*. 2008;92(1):151-151. doi:10.1136/bjo.2007.116186

4. Shields JA, Mashayekhi A, Kligman BE, et al. Vascular tumors of the conjunctiva in 140 cases. *Ophthalmology*. 2011;118(9):1747-1753. doi:10.1016/j.opht.2011.04.034

5. Eduardo-Sánchez YW, Fernández-Agrafiojo D. Sarcoma de Kaposi conjuntival gigante por el virus de inmunodeficiencia humana: remisión total tras terapia antirretroviral y quimioterapia sistémica. *Arch Soc Esp Oftalmol (Engl Ed)*. 2018;93(6):310-312. doi:10.1016/j.oftal.2017.07.010

6. Arora R, Sandhu N, Dokania P, Subramanian A. Ocular Manifestations in Patients of HIV(Human Immunodeficiency Virus) Infection on Combined Anti-Retroviral Therapy (CART). *Ocul Immunol Inflamm*. 2022;30(6):1399-1407. doi:10.1080/09273948.2021.1881562