
- 34. Kempen JH, Altaweel MM, Drye LT, et al. Benefits of systemic anti-inflammatory therapy versus fluocinolone acetonide intraocular implant for intermediate uveitis, posterior uveitis, and panuveitis: fifty-four-month results of the Multicenter Uveitis Steroid Treatment (MUST) Trial and Follow-up Study. *Ophthalmology*. 2015;122:1967—1975.
- 35. Tomkins-Netzer O, Lightman S, Drye L, et al. Outcome of treatment of uveitic macular edema: the Multicenter Uveitis Steroid Treatment Trial 2-year results. *Ophthalmology*. 2015;122:2351–2359.
- **36.** Pavesio C, Zierhut M, Bairi K, et al. Evaluation of an intravitreal fluocinolone acetonide implant versus standard systemic therapy in noninfectious posterior uveitis. *Ophthalmology*. 2010;117:567–575.
- 37. Jaffe GJ, Pavesio CE, Study Investigators. Effect of a fluocinolone acetonide insert on recurrence rates in noninfectious intermediate, posterior, or panuveitis: three-year results. *Ophthalmology*. 2020;127:1395—1404.
- 38. Kempen JH, Altaweel MM, Holbrook JT, et al. The Multicenter Uveitis Steroid Treatment Trial: rationale, design, and baseline characteristics. *Am J Ophthalmol*. 2010;149: 550–561.
- 39. Thorne JE, Sugar EA, Holbrook JT, et al. Periocular triamcinolone vs. intravitreal triamcinolone vs. intravitreal dexamethasone implant for the treatment of uveitic macular edema: the PeriOcular vs. INTravitreal corticosteroids for uveitic macular edema (POINT) trial. *Ophthalmology*. 2019;126:283–295.

Pictures & Perspectives



Dendritic Corneal Stromal Pigmentation: A Spillover of Nevus of Ota?

A 6-year-old girl presented (**A**) with congenital unilateral slate-grey hyperpigmentation of the eyelids, cheek, sclera, and (**B**) she had a brown, macular periumbilical dermal nevus. She had an intracorneal, brown, carpet-like lesion within the inferonasal corneal stroma, sparing the pupillary area. Her vision and intraocular pressure were within normal limits. Dendrite-like pigmentation fanned out intrastromally (**C**) within posterior stroma as on anterior-segment OCT (**D**), adjacent to scleral pigmentation, possibly representing scleral extension of Nevus of Ota. (Magnified version of Figure **A-D** is available online at www.aaojournal.org).

SOMYA KUMARI, MD ABHISHEK SINGH, BOPTOM SHIKHA GUPTA, MD

Dr. Rajendra Prasad Centre for Ophthalmic Sciences, All India Institute of Medical Sciences, New Delhi, India