


- cohort of 6336 patients and results of a survey. *Ann Surg.* 2004;240(2):205.
36. Strasberg SM, Linehan DC, Hawkins WG. The accordion severity grading system of surgical complications. *Ann Surg.* 2009;250(2):177–186.
37. Strong VE, Selby LV, Sovel M, et al. Development and assessment of Memorial Sloan Kettering Cancer Center's surgical secondary events grading system. *Ann Surg Oncol.* 2015;22:1061–1067.
38. Brown EG. Using MedDRA: implications for risk management. *Drug Saf.* 2004;27(8):591–602.
39. Sii S, Barton K, Pasquale LR, et al. Reporting harm in glaucoma surgical trials: systematic review and a consensus-derived new classification system. *Am J Ophthalmol.* 2018;194:153–162.
40. Munk MR, Kashani AH, Tadayoni R, et al. Recommendations for OCT angiography reporting in retinal vascular disease: a Delphi approach by international experts. *Ophthalmol Retina.* 2022;6(9):753–761.
41. Kodjikian L, Baillif S, Couturier A, et al. Recommendations for the management of diabetic macular oedema with intravitreal dexamethasone implant: a national Delphi consensus study. *Eur J Ophthalmol.* 2022;32(5):2845–2856.
42. Cruz-Pimentel M, Wu L. Complement inhibitors for advanced dry age-related macular degeneration (geographic atrophy): some light at the end of the tunnel? *J Clin Med.* 2023;12(15):5131.
43. Ecken P, Gnatzy T, von der Gracht HA. Desirability bias in foresight: consequences for decision quality based on Delphi results. *Technol Forecast Soc Change.* 2011;78(9):1654–1670.

## Pictures & Perspectives



### Dilated Conjunctival Vasculature in a Patient Up-Titrating Treprostinil Therapy

A 61-year-old man with a history of heritable pulmonary hypertension (+BNPR2) was hospitalized for up-titration of treprostinil. The patient developed acute, bilateral “red eye” without visual symptoms, visual acuity of 20/20, normal intraocular pressures, and symmetrically reactive pupils. Examination demonstrated bilateral, dilated circumferential conjunctival vasculature surrounding the limbus with temporal extension (A) and relative nasal sparing (B). Computed tomography angiography was reassuring against carotid-cavernous fistula. Dilated conjunctival vasculature has been reported secondary to epoprostenol; however, this is the first reported case secondary to treprostinil therapy. He remained asymptomatic with preserved afferent function on follow-up. (Magnified version of Figure A–B is available online at [www.aaojournal.org](http://www.aaojournal.org)).

JOHN WILSON, MD<sup>1</sup>

ELI PRATTE, MD<sup>1</sup>

NATHANIEL BLECHER, MD<sup>1</sup>

<sup>1</sup>Department of Ophthalmology, Washington University School of Medicine, St. Louis, Missouri