


- GAN. In: Shen D, Liu T, Peters TM, et al, eds. *Medical Image Computing and Computer Assisted Intervention—MICCAI 2019*. Springer International Publishing; 2019:623–631.
28. Chiu SJ, Allingham MJ, Mettu PS, et al. Kernel regression based segmentation of optical coherence tomography images with diabetic macular edema. *Biomed Opt Express*. 2015;6(4):1172–1194.
 29. Kim J, Kim M, Kang H, Lee K. U-GAT-IT: Unsupervised generative attentional networks with adaptive layer-instance normalization for image-to-image translation. *International Conference on Learning Representations*. 1; 2019. <http://arxiv.org/abs/1907.10830>. Accessed April 1, 2021.
 30. Ronneberger O, Fischer P, Brox T. U-Net: convolutional Networks for biomedical image segmentation. In: Navab N, Frangi AF, Wells W, Maie A. *Medical Image Computing and Computer-Assisted Intervention—MICCAI*. Springer International Publishing; 2015:234–241.
 31. Ma X, Huang H, Wang Y, et al. Normalized loss functions for deep learning with noisy labels. In: Iii HD, Singh A, eds. *Proceedings of the 37th International Conference on Machine Learning*. Vol 119. *Proceedings of Machine Learning Research*. Proceedings of Machine Learning Research; 2020: 6543–6553.
 32. Miller AR, Roisman L, Zhang Q, et al. Comparison between spectral-domain and swept-source optical coherence tomography angiographic imaging of choroidal neovascularization. *Invest Ophthalmol Vis Sci*. 2017;58(3):1499–1505.
 33. Yasaka K, Abe O. Deep learning and artificial intelligence in radiology: current applications and future directions. *PLoS Med*. 2018;15(11):e1002707.
 34. Pazos M, Dyrda AA, Biarnés M, et al. Diagnostic accuracy of Spectralis SD OCT automated macular layers segmentation to discriminate normal from early glaucomatous eyes. *Ophthalmology*. 2017;124(8):1218–1228.
 35. Mehta N, Lee CS, Mendonça LSM, et al. Model-to-data approach for deep learning in optical coherence tomography intraretinal fluid segmentation. *JAMA Ophthalmol*. 2020;138(10):1017–1024.
 36. Liefers B, Colijn JM, González-Gonzalo C, et al. A deep learning model for segmentation of geographic atrophy to study its long-term natural history. *Ophthalmology*. 2020;127(8):1086–1096.
 37. Van Brummen A, Owen JP, Spaide T, et al. PeriorbitAI: artificial intelligence automation of eyelid and periorbital measurements. *Am J Ophthalmol*. 2021;230: 285–296.

Pictures & Perspectives



Squamous Cell Papilloma of the Conjunctiva

A 74-year-old man presented to the ophthalmology department with an 8-month history of a tumor in his left eye. Examination showed a fleshy tumor with a cauliflower-like appearance involving the conjunctiva and cornea (Fig A). An excisional biopsy was performed, and pathological analysis showed squamous cell papilloma of the conjunctiva. Histology slide showed papilloma of the conjunctiva (Fig B). Conjunctival papilloma is an acquired benign epithelial tumor of the conjunctiva, and may manifest as a sessile or pedunculated conjunctival mass. Conjunctival papilloma often requires surgical excision. Other interventions include perilesional cryotherapy, intralesional treatment, or multiple choices of topical treatment modalities for managing conjunctival papilloma, and recurrences are not uncommon (Magnified version of Fig A-B is available online at www.aaojournal.org).

LIPING WANG, MD¹

GUOPING WANG, MD²

¹Ineye Hospital of Chengdu University of Traditional Chinese Medicine, Chengdu, Sichuan, China; ²Eye School of Chengdu University of Traditional Chinese Medicine, Chengdu, Sichuan, China