


- intraocular lenses. Results of a multicenter questionnaire. *J Cataract Refract Surg*. 2001;27:245–249.
5. Vingolo EM, Grenga P, Iacobelli L, Grenga R. Visual acuity and contrast sensitivity: AcrySof ReSTOR apodized diffractive versus AcrySof SA60AT monofocal intraocular lenses. *J Cataract Refract Surg*. 2007;33:1244–1247.
  6. United States Food and Drug Administration. Patient-reported outcome measures: use in medical product development to support labeling claims. 2009. Available at: <https://www.fda.gov/regulatory-information/search-fda-guidance-documents/patient-reported-outcome-measures-use-medical-product-development-support-labeling-claims>. Accessed February 2, 2023.
  7. Cronbach LJ. Coefficient alpha and the internal structure of tests. *Psychometrika*. 1951;16:297–334.
  8. SAS Institute. SAS®9.4 TS level 1M5. 2010. <http://www.sas.com/>. Accessed February 1, 2023.
  9. Schallhorn SC, Hettinger KA, Pelouskova M, et al. Effect of residual astigmatism on uncorrected visual acuity and patient satisfaction in pseudophakic patients. *J Cataract Refract Surg*. 2021;47:991–998.
  10. McAlinden C, Pesudovs K, Moore JE. The development of an instrument to measure quality of vision: the Quality of Vision (QOV) questionnaire. *Invest Ophthalmol Vis Sci*. 2010;51:5537–5545.
  11. Lasch K, Marcus JC, Seo C, et al. Development and validation of a visual symptom-specific patient-reported outcomes instrument for adults with cataract intraocular lens implants. *Am J Ophthalmol*. 2022;237:91–103.

## Pictures & Perspectives



### Rosai-Dorfman Disease

An 18-year-old man presented with a 3-month history of a rapidly growing, subconjunctival, salmon-colored lesion in the left medial palpebral aperture (A). Excisional biopsy removed a well-defined, gelatinous mass (B). Histopathology revealed lymphocytes, plasma cells and histiocytes with large, round nuclei, voluminous cytoplasm, and occasional emperipolesis (engulfment of lymphocytes) (C, arrow). Histiocytes demonstrated positive S100 immunostaining (D), confirming the diagnosis of extra-nodal Rosai-Dorfman disease. Rosai-Dorfman disease is a rare, non-Langerhans histiocytosis of unknown etiology predominantly affecting children and young adults. Unifocal disease can be treated surgically, although immunosuppressive agents are used for more extensive disease. The clinical course is typically benign (Magnified version of Figure A–D is available online at [www.aaojournal.org](http://www.aaojournal.org)).

BENJAMIN ZHOU, BS<sup>1</sup>

DAVID I. SUSTER, MD<sup>2</sup>

PAUL D. LANGER, MD, FACS<sup>1</sup>

<sup>1</sup>Institute of Ophthalmology and Visual Science, Rutgers New Jersey Medical School, Newark, New Jersey; <sup>2</sup>Department of Pathology, Rutgers New Jersey Medical School, Newark, New Jersey