

Clothesline Phenomenon


An 81-year-old man with a history of repaired rhegmatogenous retinal detachment in his left eye 45 years prior presented with a nasal scotoma after blunt trauma. A localized superotemporal rhegmatogenous retinal detachment occurred at the site of intrusion of the Arruga's suprapiramid ring suture (A), known as "Clothesline phenomenon." Postoperative appearance after 23-G pars plana vitrectomy (B) and laser to the area around the intruded suture. (Magnified version of Figure A-B is available online at www.aaojournal.org).

JOSE GRAU VERGER, MD¹

PERE GARCIA BRU, MD¹

HELENA BROSÀ MORROS, MD²

¹Department of Ophthalmology, Bellvitge University Hospital, Hospitalet de Llobregat, Spain; ²Department of Ophthalmology, Vall Hebron University Hospital, Barcelona, Spain



Spontaneous Rupture of Posttraumatic Iris Cyst

A 34-year-old man with a 1-year history of repaired temporal paralimbal scleral perforation presented with a well-circumscribed pearl-type inferior iris cyst and residual eyelash (A, arrow) in anterior chamber. Swept-source anterior-segment OCT demonstrated a hyperreflective dome-shaped mass obscuring inferior angle (B). On follow-up, it ruptured spontaneously (C). Ultrasound biomicroscopy revealed homogeneously echogenic ruptured cyst limited to iris, confirming its origin (D). Surgical excision with iridotomy and the removal of an eyelash was done. Traumatic cysts may present with a varying latent period and rupture spontaneously. Imaging plays a vital role to rule out underlying tumors (Magnified version of Figure A-D is available online at www.aaojournal.org).

AAFREEN BARI, MD

RITWEEZ SAHU, MBBS

RAJESH SINHA, MD

Cornea, Cataract & Refractive Surgery Services, Dr Rajendra Prasad Centre for Ophthalmic Sciences, All India Institute of Medical Sciences, New Delhi, India
