


40. Randleman JB, Wolfe JD, Woodward M, et al. The resident surgeon phacoemulsification learning curve. *Arch Ophthalmol.* 2007;125:1215–1219.
41. Heier JS, Khanani AM, Quezada Ruiz C, et al. Efficacy, durability, and safety of intravitreal faricimab up to every 16 weeks for neovascular age-related macular degeneration (TENAYA and LUCERNE): two randomised, double-masked, phase 3, non-inferiority trials. *Lancet.* 2022;399:729–740.
42. Regeneron Pharmaceuticals Inc. *Aflibercept 8 mg late-breaking data presented at Retina Society in Diabetic Macular Edema and Wet Age-Related Macular Degeneration.* Tarrytown, NY: Regeneron Pharmaceuticals, Inc.; 2022.

## Pictures & Perspectives

---



### Infrequent Bilateral Orbital Tumors

A 44-year-old woman presented with a 1-year history of progressive, painless bilateral proptosis. Magnetic resonance imaging revealed well-circumscribed, enhancing, central masses (A–B). Right mass resection was performed via an inferior fornix incision. Left mass resection was performed via a lateral orbitotomy. Histopathology showed the right-sided mass to be rich in vascular sinuses, lined with flat endothelial cells separated by fibrous intervals (characteristic of cavernous hemangioma, C). Histopathology of the left-sided mass demonstrated variably sized, exuberantly proliferous nerve fascicles with myelinated axons, consistent with a schwannoma (D). Follow-up over 1 year has shown no recurrence (Magnified version of Figure A–D is available online at [www.aaojournal.org](http://www.aaojournal.org)).

RUIMIAO LI, MMED

MINGYU REN, MMED

LIMIN LIU, MMED

Department of Orbital Disease and Ocular Tumor, Hebei Eye Hospital, Xingtai, Hebei, China

---