


## Ophthalmic Images


# Laser in Situ Keratomileusis Flap Free-Floating on Interface Edema in Acute Corneal Hydrops

Evan J. Warner, MD

A Slitlamp photography



B Anterior segment OCT



**Figure.** A, Slitlamp photograph showing severe stromal edema and a honeycomb pattern of diffuse epithelial bullae. B, Anterior segment optical coherence tomography (OCT) demonstrates stromal edema, a lamella of fluid in the laser in situ keratomileusis flap interface, and intact flap with diffuse epithelial fluid cysts. (Photograph courtesy of Dr Warner).

**A 62-year-old patient** with a remote history of laser in situ keratomileusis (LASIK) awoke with cloudy vision, tearing, photophobia, and sharp pain in the left eye. She reported requiring a rigid gas permeable contact for 11 years due to postrefractive ectasia. Slitlamp examination was consistent with acute hydrops (Figure, A). Central pachymetry measured over 1400  $\mu\text{m}$  (fellow eye, 588  $\mu\text{m}$ ). Optical coherence tomography identified fluid channels throughout the stroma and an unstable LASIK flap resting on fluid in the flap interface (Figure, B). She was treated con-

servatively with topical moxifloxacin, prednisolone, and timolol, and the edema resolved by 11 weeks, leaving a dense scar. Six weeks later, she developed redness, pain, and photophobia and was found to have a small, branching posterior stromal infiltrate concerning for fungal keratitis. Penetrating keratoplasty was performed, with pathology identifying *Aspergillus* at 80% stromal depth. The postoperative course was uncomplicated, and final visual acuity was correctable to 20/15. It was suspected that use of topical corticosteroid in conjunction with channels of edema communicating with tenuous surface bullae led to a deep fungal keratitis.

## ARTICLE INFORMATION

**Author Affiliation:** Department of Ophthalmology and Visual Sciences, University of Wisconsin-Madison School of Medicine and Public Health, Madison.

**Corresponding Author:** Evan J. Warner, MD, Department of Ophthalmology and Visual Sciences, University of Wisconsin-Madison School of

Medicine and Public Health, 2870 University Ave, Ste 206, Madison, WI 53705 (ejwarner@wisc.edu).

**Conflict of Interest Disclosures:** Dr Warner reported receiving grants from Research to Prevent Blindness Inc. No other disclosures were reported.

**Funding/Support:** This work was supported in part by an unrestricted grant from Research to Prevent Blindness Inc to the University of

Wisconsin-Madison Department of Ophthalmology and Visual Sciences.

**Role of the Funder/Sponsor:** The funder had no role in the decision to submit this case report for publication.

**Additional Contributions:** We thank the patient for granting permission to publish this information.