

Ophthalmic Images

Double Optic Disc Pit With Retinal Pigmentation in a Woman With High Myopia

Zhiyan Xu, MD; Zhikun Yang, MD




Figure. A, Color fundus montage photograph of the left eye showing an enlarged optic disc with 2 optic disc pits (ODPs, black arrowheads) surrounded by areas of retinal pigmentation and depigmentation, where some large choroidal vessels are visible. B, Swept-source optical coherence tomography of the optic disc with a scanned green line through the ODPs. This oblique line scan revealed ODPs as 2 distinct excavations of the optic disc (yellow arrowheads) with discontinuance of the ellipsoid zone and disorganization of the outer retinal layers in the adjacent retina (right image).

A 37-year-old woman with high myopia presented for a routine ocular checkup visit. Her refractive error was -10.00 diopters (D) OD, -7.75 D OS, and best-corrected visual acuity was 20/20 OU. Her axial length (IOL Master [Zeiss]) was 27.27 mm in the right eye and 26.01 mm in the left eye. Intraocular pressure and anterior segment examination were normal. Fundus examination revealed an enlarged optic disc with 2 optic disc pits (ODPs) in the left eye, located at the nasal and inferotemporal rim (Figure, A, black arrowheads). Two areas of retinal pigmentation adjacent to

the pits were noted. B-scan swept-source optical coherence tomography imaging revealed 2 distinct excavations of the optic disc (Figure, B, yellow arrowheads). Visual field test of the left eye showed an enlarged blind spot and a localized temporal defect, correlating with the pigmented areas nasal to the disc. Patients with ODPs are usually asymptomatic unless they develop maculopathy.^{1,2} This patient's fundus examination and visual field test remained unchanged over a 1.5-year period of follow-up.

ARTICLE INFORMATION

Author Affiliations: Department of Ophthalmology, Peking Union Medical College Hospital, Beijing, China (Xu, Yang); Key Laboratory of Ocular Fundus Diseases, Chinese Academy of Medical Sciences, Beijing, China (Xu, Yang).

Corresponding Author: Zhikun Yang, MD, Department of Ophthalmology, Peking Union Medical College Hospital, No. 1 Shuaifuyuan Wangfujing, Dongcheng District, Beijing 100730, China (yangzhikun@pumch.cn).

Conflict of Interest Disclosures: None reported.

Additional Contributions: We thank the patient for granting permission to publish this information.

REFERENCES

- Boese EA, Huang D, Tehrani S. Unilateral double optic nerve head pits. *Ophthalmology*. 2018;125(3):458. doi:10.1016/j.ophtha.2017.12.009
- Georgalas I, Ladas I, Georgopoulos G, Petrou P. Optic disc pit: a review. *Graefes Arch Clin Exp*

Ophthalmol. 2011;249(8):1113-1122. doi:10.1007/s00417-011-1698-5