

1. Congressional Budget Office. Prescription drugs: spending, use, and prices. 2022. Accessed December 6, 2023. [https://www.cbo.gov/publication/57772#\\_idTextAnchor004](https://www.cbo.gov/publication/57772#_idTextAnchor004)
2. Cubanski J, Neuman T. A small number of drugs account for a large share of Medicare Part D spending. KFF. July 12, 2023. Accessed December 6, 2023. <https://www.kff.org/medicare/issue-brief/a-small-number-of-drugs-account-for-a-large-share-of-medicare-part-d-spending/>
3. DrugBank Online. Home page. Accessed April 6, 2023. <https://go.drugbank.com/>
4. Centers for Medicare & Medicaid Services. 2022 Medicare Part D spending by drug. Accessed April 6, 2023. <https://data.cms.gov/summary-statistics-on-use-and-payments/medicare-medicaid-spending-by-drug/medicare-part-d-spending-by-drug/data>
5. Chen EM, Kombo N, Teng CC, Mruthyunjaya P, Nwanyanwu K, Parikh R. Ophthalmic medication expenditures and out-of-pocket spending: an analysis of United States prescriptions from 2007 through 2016. *Ophthalmology*. 2020; 127(10):1292-1302. doi:10.1016/j.ophtha.2020.04.037
6. Berkowitz ST, Groth S, Sternberg P Jr, Patel S. Drug savings for Medicare Part D beneficiaries using a direct-to-consumer model. *Ophthalmology*. 2024;131(4):509-510. doi:10.1016/j.ophtha.2023.12.004

## OBSERVATION

### Treatment of Vernal Keratoconjunctivitis Using Upadacitinib


Vernal keratoconjunctivitis (VKC) is a severe and chronic ocular allergic disease that predominantly affects children. Clinically, VKC is characterized by the formation of giant papillae on the upper palpebral conjunctiva and corneal damage caused by potent type 2 and eosinophilic inflammation, resulting in vision loss. As severe cases are often refractory to topical treatment using steroids or immunosuppressants, the development of novel therapeutic strategies is imperative.<sup>1-3</sup> Janus kinase (JAK), an intracellular tyrosine kinase, regulates cytokine signaling pathways. Currently, JAK inhibitors are used to treat atopic dermatitis (AD).<sup>4</sup> This report presents a case of improvement of VKC associated with initiation of an oral JAK inhibitor upadacitinib.

**Report of a Case** | A 13-year-old girl presented with exacerbation of symptoms including photophobia, itchy eyes, and pain in her right eye. Slitlamp examination revealed giant papillae on the upper tarsal conjunctiva and superficial

punctate keratitis in the right eye (**Figure 1**). Her best-corrected visual acuity was 20/63 OD and 20/16 OS. She had previously been diagnosed with VKC associated with AD at 9 years old and for 4 years was treated with eye drops containing tacrolimus, cyclosporine, steroids, and antiallergic agents. Despite topical treatment with eye drops, the giant papillae on the tarsal conjunctiva continued to worsen and resolve seasonally. Corneal lesions such as a shield ulcer and superficial punctate keratitis in her right eye persisted for 3 months. AD was observed on her skin, including the face, and was treated with tacrolimus ointments, steroid ointments, and a moisturizer. As the patient's AD was severe with persistent skin rash refractory to treatment with topical medications, oral upadacitinib (15 mg/d) was administered for AD. Scores for the Eczema Area and Severity Index and the Investigator's Global Assessment before and 2 weeks after upadacitinib treatment improved from 28.0 to 1.2 and 3 to 2, respectively. The pain and itching in the right eye improved within a month after upadacitinib administration. Additionally, the giant papillae gradually flattened, and corneal lesions improved after upadacitinib administration. After 3 months of oral upadacitinib administration, the giant papillae and punctate epithelial keratitis resolved completely (**Figure 2A and B**), and vision improved to 20/16 OD. After 11 months of treatment, upadacitinib dosage was increased to 30 mg/d owing to the exacerbation of AD. After more than 14 months of upadacitinib treatment, no recurrence of giant papillae or corneal damage was observed with topical treatment of only antiallergic eye drops (**Figure 2C and D**), and visual acuity was maintained at 20/16 OD and 20/20 OS.


**Discussion** | Keratoconjunctivitis has been reported to improve during treatment of AD by upadacitinib. Hayama et al<sup>5</sup> reported improvement of dupilumab-associated conjunctivitis after switching to upadacitinib in a patient with AD. Ghigloni et al<sup>6</sup> were the first to show the improvement of atopic keratoconjunctivitis during treatment with upadacitinib. In this case involving VKC resistant to topical treatment, the giant papillae of conjunctiva and corneal lesions resolved within 3

**Figure 1. Slitlamp Images for Upper Tarsal Conjunctiva and Cornea With Fluorescein Staining Before Upadacitinib Administration**



Giant papillae on the upper tarsal conjunctiva (arrowheads) (A) and superficial punctate keratitis (B) were apparent before upadacitinib administration.

**Figure 2. Slitlamp Images for Upper Tarsal Conjunctiva and Cornea With Fluorescein Staining After Upadacitinib Administration**



Giant papillae flattened (arrowheads) (A) and punctate epithelial keratitis (B) disappeared completely after 3 months of oral administration of upadacitinib. C, No recurrence of giant papillae or corneal damage was observed for the next 14 months.

months following initiation of oral upadacitinib, without recurrence for the next 14 months. To the best of our knowledge, this report is the first to describe this effect in association with initiation of upadacitinib in patients with VKC associated with AD. Important cytokines for ocular allergic inflammation including type 2 cytokines interleukin 4 and interleukin 13 and epithelium-derived cytokine thymic stromal lymphopoietin signal through the JAK1 pathway.<sup>4</sup> As upadacitinib is a selective JAK1 inhibitor, this case suggests that cytokine signaling through the JAK1 pathway may play critical roles in VKC pathogenesis. Although this single-case report cannot prove a cause-and-effect relationship and further studies and clinical trials would be needed to determine its efficacy, upadacitinib is a potential novel therapeutic agent against VKC, especially in cases with AD.

**Kazuhiro Kito, MD, PhD**

**Ken Fukuda, MD, PhD**

**Tatsuma Kishimoto, MD, PhD**

**Kenji Yamashiro, MD, PhD**

**Akiko Yagami, MD, PhD**

**Author Affiliations:** Ueji Eye Clinic, Okazaki City, Aichi, Japan (Kito); Department of Ophthalmology and Visual Science Kochi Medical School, Kochi University, Nankoku City, Kochi, Japan (Fukuda, Kishimoto, Yamashiro); Department of Allergology, Fujita Health University School of Medicine, Nagoya, Aichi, Japan (Yagami).

**Corresponding Author:** Ken Fukuda, MD, PhD, Department of Ophthalmology and Visual Science, Kochi Medical School, Kochi University, Kohasu, Oko-cho, Nankoku City, Kochi 783-8505, Japan ([k.fukuda@kochi-u.ac.jp](mailto:k.fukuda@kochi-u.ac.jp)).

**Published Online:** May 16, 2024. doi:10.1001/jamaophthalmol.2024.1439

**Conflict of Interest Disclosures:** Dr Fukuda reported lecture fees from Santen Pharmaceutical, Senju Pharmaceutical Co, Sanofi, Sandoz K.K., Taiho Pharmaceutical, and AbbVie. Dr Kishimoto reported personal fees from AbbVie Inc and Otsuka Pharmaceutical. No other disclosures were reported.

**Additional Contributions:** We thank the patient's parent for granting permission to publish this information.

1. Ali A, Bielory L, Dotchin S, Hamel P, Strube YNJ, Koo EB. Management of vernal keratoconjunctivitis: navigating a changing treatment landscape. *Surv Ophthalmol*. 2023;50(3):6257(23)00138-8. doi:10.1016/j.survophthal.2023.10.008
2. Cheung AY, Choi DS, Ahmad S, et al; American Academy of Ophthalmology Preferred Practice Pattern Cornea/External Disease Panel. Conjunctivitis preferred practice pattern. *Ophthalmology*. 2024;131(4):134-P204. doi:10.1016/j.ophtha.2023.12.037
3. Bielory L, Delgado L, Katelaris CH, Leonardi A, Rosario N, Vichyanoud P. ICON: diagnosis and management of allergic conjunctivitis. *Ann Allergy Asthma Immunol*. 2020;124(2):118-134. doi:10.1016/j.ana.2019.11.014
4. Nakashima C, Yanagihara S, Otsuka A. Innovation in the treatment of atopic dermatitis: emerging topical and oral Janus kinase inhibitors. *Allergol Int*. 2022; 71(1):40-46. doi:10.1016/j.alit.2021.10.004
5. Hayama K, Fujita H. Improvement of dupilumab-associated conjunctivitis after switching to upadacitinib in a patient with atopic dermatitis. *Dermatol Ther*. 2022;35(7):e15575. doi:10.1016/j.dth.15575
6. Ghigloni DG, Cozzi L, Pigazzi C, et al. Improvement of atopic keratoconjunctivitis during treatment with upadacitinib for atopic dermatitis. *Aust J Dermatol*. 2023;2023(10):1103.