


30. Xie J, Arnold A-L, Keeffe J, et al. Prevalence of self-reported diabetes and diabetic retinopathy in indigenous Australians: the National Indigenous Eye Health Survey. *Clin Exp Ophthalmol*. 2011;39:487–493.
31. Moynihan V, Turner A. Coordination of diabetic retinopathy screening in the Kimberley region of Western Australia. *Aust J Rural Health*. 2017;25:110–115.
32. Brazionis L, Jenkins A, Keech A, et al. Diabetic retinopathy in a remote Indigenous primary healthcare population: a Central Australian diabetic retinopathy screening study in the Telehealth Eye and Associated Medical Services Network project. *Diabet Med*. 2018;35:630–639.
33. O'Halloran RA, Turner AW. Evaluating the impact of optical coherence tomography in diabetic retinopathy screening for an Aboriginal population. *Clin Exp Ophthalmol*. 2018;46:116–121.
34. Quinn N, Yang F, Ryan C, et al. Screening for diabetic retinopathy and reduced vision among Indigenous Australians in Top End primary care health services: a TEAMSnet sub-study. *Intern Med J*. 2021;51:1897–1905.
35. Tapp RJ, Shaw JE, Harper CA, et al. The prevalence of and factors associated with diabetic retinopathy in the Australian population. *Diabetes Care*. 2003;26:1731–1737.
36. Cugati S, Kifley A, Mitchell P, et al. Temporal trends in the age-specific prevalence of diabetes and diabetic retinopathy in older persons: population-based survey findings. *Diabetes Res Clin Pract*. 2006;74:301–308.
37. Wong J, Molyneaux L, Constantino M, et al. Timing is everything: age of onset influences long-term retinopathy risk in type 2 diabetes, independent of traditional risk factors. *Diabetes Care*. 2008;31:1985–1990.
38. Copeland S, Muir J, Turner A. Understanding Indigenous patient attendance: a qualitative study. *Aust J Rural Health*. 2017;25:268–274.
39. Chia MA, Turner AW. Benefits of integrating telemedicine and artificial intelligence into outreach eye care: stepwise approach and future directions. *Front Med*. 2022;9:835804.

Pictures & Perspectives



Ultrawide-field Swept Source-OCT Angiography of Retinitis Pigmentosa

A 42-year-old man diagnosed with retinitis pigmentosa (RP) underwent widefield Optos imaging and swept-source OCT angiography (SS-OCTA) (24 × 20 mm SS-OCTA; BM-400K BMizar, TowardPi Medical Technology) examination. **A**, Fundus image revealed classic features of RP, including intraretinal pigmentation, waxy pallor of the optic disc, and attenuated retinal blood vessels. **B**, The retinal flow image displayed perfusion limited to the posterior pole with vascular remodeling at the borders. **C**, The retinal structure image showed intraretinal pigmentation outside the arcade (red arrows) and the empty shells of nonperfused “ghost” vessels (yellow arrows) (Magnified version of Fig A-C is available online at www.aaojournal.org).

FANG ZHENG, MD
JINGLIANG HE, MD
XIAOYUN FANG, MD

Eye Center, The Second Affiliated Hospital Zhejiang University School of Medicine, Zhejiang University, Hangzhou, China