


## Ophthalmic Images

## Morning Glory Syndrome in a Newborn Aged 2 Weeks

Francesco Pozzo Giuffrida, MD; Marco Nassisi, MD, PhD; Silvia Osnaghi, MD



**Figure.** Red-free fundus image and magnetic resonance imaging (MRI) scan of a male newborn with morning glory syndrome. A, Red-free fundus photography shows the morning glory disc anomaly. B, Axial T2-weighted MRI image shows a funnel-shaped morphologic pattern of the left optic disc (blue arrowhead) and frontonasal encephalocele (yellow arrowheads). Asymmetric appearance of brain parenchyma is due to head tilting.

**A 2-week-old newborn** was referred to a pediatric ophthalmology department for divergent strabismus. The infant had a superior cleft lip, hypertelorism, and a morning glory disc anomaly in the left eye (Figure, A). Family history and prenatal karyotype test results were unremarkable. A magnetic resonance imaging scan of the brain revealed a funnel-shaped optic disc in the left eye with ipsilateral prechiasmatic optic nerve thickening and a skull base defect with sphenothmoidal encephalocele and ectopia of the gland

(Figure, B). A diagnosis of morning glory syndrome<sup>1</sup> was made because systemic findings were associated with the optic disc anomaly. The patient needed surgical intervention to correct the herniated tissue and the skull defect, which was postponed until the infant weighed over 7 kg. Morning glory disc anomaly is frequently associated with midline defects<sup>2</sup> and cerebrovascular anomalies.<sup>3,4</sup> Early interdisciplinary evaluation, including neuroimaging, would seem important to rule out potential brain and basal skull defects that can lead to serious surgical complications if not diagnosed in time.

## ARTICLE INFORMATION

**Author Affiliations:** Department of Clinical Sciences and Community Health, University of Milan, Milan, Italy (Giuffrida, Nassisi); Ophthalmological Unit, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy (Giuffrida, Nassisi, Osnaghi).

**Corresponding Author:** Marco Nassisi, MD, PhD, Ophthalmological Unit, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan 20122, Italy ([marco.nassisi@unimi.it](mailto:marco.nassisi@unimi.it)).

**Conflict of Interest Disclosures:** None reported.

**Additional Contributions:** We thank the patient's mother for granting permission to publish this

information and Laura Dell'Arti, MD, and Francesco Viola, MD (Ospedale Maggiore Policlinico), and Ilaria Cocchi, MD, Irene Degrassi, MD, and Chiara Doneda, MD (Vittore Buzzi Children's Hospital), all of whom contributed to the management of the clinical case. No one was financially compensated for their contribution.

## REFERENCES

- Kindler P. Morning glory syndrome: unusual congenital optic disk anomaly. *Am J Ophthalmol*. 1970;69(3):376-384. doi:10.1016/0002-9394(70)92269-5
- Bakri SJ, Siker D, Masaryk T, Luciano MG, Traboulsi EI. Ocular malformations, moyamoya

disease, and midline cranial defects: a distinct syndrome. *Am J Ophthalmol*. 1999;127(3):356-357. doi:10.1016/S0002-9394(98)00317-1

3. Eustis HS, Sanders MR, Zimmerman T. Morning glory syndrome in children—association with endocrine and central nervous system anomalies. *Arch Ophthalmol*. 1994;112(2):204-207. doi:10.1001/archophth.1994.01090140080027

4. Lenhart PD, Lambert SR, Newman NJ, et al. Intracranial vascular anomalies in patients with morning glory disk anomaly. *Am J Ophthalmol*. 2006;142(4):644-650. doi:10.1016/j.ajo.2006.05.040