


29. Mitchell P, Holz FG, Hykin P, et al. Efficacy and safety of intravitreal aflibercept using a treat-and-extend regimen for neovascular age-related macular degeneration: the ARIES study: a randomized clinical trial. *Retina*. 2021;41:1911–1920.
30. Ebnetter A, Michels S, Prunte C, et al. Two-year outcomes of intravitreal aflibercept in a Swiss routine treat and extend regimen for patients with neovascular age-related macular degeneration. *Sci Rep*. 2020;10:20256.
31. Singh RP, Yu JS, Guruprasad B, Agashivala N. Anti-VEGF treatment patterns in patients with wet age-related macular degeneration in clinical practice. *Ophthalmic Surg Lasers Imaging Retina*. 2021;52:263–272.

## Pictures & Perspectives



### Conjunctival Spindle Cell/Sclerosing Rhabdomyosarcoma

A 13-year-old Asian boy had a fast-growing solid mass on his right lower palpebral conjunctiva and medial canthus for 1 month (A). Magnetic resonance imaging revealed 2 masses in the medial and inferior orbit. Histopathology of the conjunctival tumor showed ovoid and spindled cells with hyperchromatic nuclei and abundant eosinophilic cytoplasm, arranged in a fascicular pattern within hyalinized stroma (arrowheads). Some exhibited a pseudovascular pattern (B, arrows). Immunohistochemical staining results were positive for desmin (C) and MyoD1 (D). Molecular analysis revealed an absence of the *PAX3/FOXO1* fusion gene. The boy was diagnosed with spindle cell/sclerosing rhabdomyosarcoma and received concurrent chemotherapy. (Magnified version of Figure A–D is available online at [www.aaojournal.org](http://www.aaojournal.org)).

YI-WEN SU, MD<sup>1</sup>

LEI-CHI WANG, MD, PhD<sup>2,3</sup>

NAI-WEN FAN, MD, PhD<sup>1,3</sup>

<sup>1</sup>Department of Ophthalmology, Taipei Veterans General Hospital, Taipei, Taiwan; <sup>2</sup>Department of Pathology and Laboratory Medicine, Taipei Veterans General Hospital, Taipei, Taiwan; <sup>3</sup>School of Medicine, National Yang Ming Chiao Tung University, Taipei, Taiwan