


Correspondence:

John C. Buchan, FRCOphth, MD, London School of Hygiene and Tropical Medicine, Keppel St., London WC1E 7HT, UK. E-mail: John.Buchan@lshtm.ac.uk.

References

1. Relhan N, Forster RK, Flynn HW. Endophthalmitis: then and now. *Am J Ophthalmol.* 2018;187:xx–xxvii.
2. Pershing S, Lum F, Hsu S, et al. Endophthalmitis after cataract surgery in the United States: a report from the Intelligent Research in Sight Registry, 2013–2017. *Ophthalmology.* 2020;127:151–158.
3. Yannuzzi NA, Si N, Relhan N, et al. Endophthalmitis after clear corneal cataract surgery: outcomes over two decades. *Am J Ophthalmol.* 2017;174:155–159.
4. Moussa G, Bhatt H, Reekie I, et al. Using the West Midlands CONCERT to characterise regional incidence of acute-onset post cataract surgery endophthalmitis. *Eye (Lond).* 2021;35:1730–1740.
5. Rahmani S, Elliott D. Postoperative endophthalmitis: a review of risk factors, prophylaxis, incidence, microbiology, treatment, and outcomes. *Semin Ophthalmol.* 2018;33:95–101.
6. Gower EW, Lindsley K, Tulenko SE, et al. Perioperative antibiotics for prevention of acute endophthalmitis after cataract surgery. *Cochrane Database Syst Rev.* 2017;2:CD006364.

Pictures & Perspectives



Hemangioma of Conjunctiva and Eyelid

A 56-year-old man presented with a dilated conjunctival vessel in the left eye and nonprogressive left upper eyelid swelling for 2 years without diminution of vision. Examination of the left eye showed a smooth, red, lobulated mass on the temporal side of the conjunctiva, with a dilated tortuous conjunctival vessel (B). The left upper eyelid showed a smooth, dark brown mass consisting of tortuous vessels (A). There was neither bruit nor pulsation. Ocular and systemic examination was otherwise within normal limits. A clinical diagnosis of conjunctival hemangioma of the left eye with left upper eyelid hemangioma was made (Magnified version of Figure A-B is available online at www.aaojournal.org).

GIRENDRA YADAV, MBBS¹

SHILPA KHANAL, MBBS, MD²

RAKSHITA YADAV, BDS¹

¹B. P. Koirala Institute of Health Sciences, Dharan, Nepal; ²Department of Ophthalmology, B. P. Koirala Institute of Health Sciences, Dharan, Nepal