


- clinical diagnosis and disease epidemiology. *Ophthalmol Retina*. 2022;6:650–656.
15. R. Version 4.2.1. R Foundation for Statistical Computing. Available at: <https://www.R-project.org/>; 2022. Accessed February 1, 2022.
 16. STARD 2015: An Updated List of Essential Items for Reporting Diagnostic Accuracy Studies. <https://www.equator-network.org/reporting-guidelines/stard/>. Accessed October 5, 2022.
 17. Shukla R, Murthy GVS, Gilbert C, et al. Operational guidelines for ROP in India: a summary. *Indian J Ophthalmol*. 2020;68(Suppl 1):S108–S114.
 18. Brown JM, Campbell JP, Beers A, et al. Automated diagnosis of plus disease in retinopathy of prematurity using deep convolutional neural networks. *JAMA Ophthalmol*. 2018;136:803.
 19. Taylor S, Brown JM, Gupta K, et al. Monitoring disease progression with a quantitative severity scale for retinopathy of prematurity using deep learning. *JAMA Ophthalmol*. 2019;137:1022.
 20. Sabri K, Ells AL, Lee EY, et al. Retinopathy of prematurity: a global perspective and recent developments. *Pediatrics*. 2022;150:e2021053924.
 21. Al-Khaled T, Valikodath NG, Patel SN, et al. Addressing the third epidemic of retinopathy of prematurity through telemedicine and technology: a systematic review. *J Pediatr Ophthalmol Strabismus*. 2021;58:261–269.
 22. Chee RI, Darwish D, Fernandez-Vega A, et al. Retinal telemedicine. *Curr Ophthalmol Rep*. 2018;6:36–45.
 23. Shah PK, Narendran V, Kalpana N. Aggressive posterior retinopathy of prematurity in large preterm babies in South India. *Arch Dis Child Fetal Neonatal Ed*. 2012;97:F371–F375.
 24. Salverda HH, Beelen DML, Cramer SJE, et al. Clinical outcomes of preterm infants while using automated controllers during standard care: comparison of cohorts with different automated titration strategies. *Arch Dis Child Fetal Neonatal Ed*. 2023;108:26–30.
 25. Srivatsa B, Hagan JL, Clark RH, Kupke KG. Oxygenation factors associated with retinopathy of prematurity in infants of extremely low birth weight. *J Pediatr*. 2022;247:46–52.e4.
 26. Gilbert C, Malik ANJ, Nahar N, et al. Epidemiology of ROP update — Africa is the new frontier. *Semin Perinatol*. 2019;43:317–322.
 27. Wang D, Duke R, Chan RP, Campbell JP. Retinopathy of prematurity in Africa: a systematic review. *Ophthalmic Epidemiol*. 2019;26:223–230.
 28. Quinn GE. Retinopathy of prematurity blindness worldwide: phenotypes in the third epidemic. *Eye Brain*. 2016;8:31–36.
 29. Coyner AS, Chen JS, Singh P, et al. Single-examination risk prediction of severe retinopathy of prematurity. *Pediatrics*. 2021;148:e2021051772.

Pictures & Perspectives



Vitrectomy for Retinoschisis in Papillorenal Syndrome Related to *PAX2* Mutation

A 36-year-old woman with a history of kidney transplantation presented with visual disturbance in the right eye. Optic disc excavation, absence of central retinal vessels, numerous cilioretinal vessels in both eyes, and macular thickening were noted in the right eye (A and B). A *PAX2* mutation was confirmed by genetic testing. OCT showed schisis of the outer plexiform and nerve fiber layers and foveal detachment with outer macular hole (C). The patient underwent vitrectomy with gas tamponade. The retina was reattached 1 year after surgery (D), and final visual acuity in the right eye improved from 20/200 to 20/63 (Magnified version of Figure A–D is available online at www.aaojournal.org).

HYUK JUN LEE, MD^{1,2}

JANG WON HEO, MD, PhD³

JAE HO JUNG, MD, PhD^{1,2}

¹Department of Ophthalmology, Seoul National University College of Medicine, Seoul, Korea; ²Department of Ophthalmology, Seoul National University Hospital, Seoul, Korea; ³The One Seoul Eye Clinic, Seoul, Korea