


- spectral-domain optical coherence tomography. *J Glaucoma*. 2015;24(5):e109–e115.
7. Gaylor DW, Sweeney HC. Design for optimal prediction in simple linear regression. *J Am Stat Assoc*. 1965;60:205–216.
 8. Crabb DP, Garway-Heath DF. Intervals between visual field tests when monitoring the glaucomatous patient: wait-and-see approach. *Invest Ophthalmol Vis Sci*. 2012;53:2770–2776.
 9. Wu Z, Medeiros FA. Impact of different visual field testing paradigms on sample size requirements for glaucoma clinical trials. *Sci. Rep*. 2018;8:4889.
 10. Leung CK, Cheung CYL, Weinreb RN, et al. Evaluation of retinal nerve fiber layer progression in glaucoma: a study on optical coherence tomography guided progression analysis. *Invest Ophthalmol Vis Sci*. 2010;51:217–222.
 11. Mwanza JC, Kim HY, Budenz DL, et al. Residual and dynamic range of retinal nerve fiber layer thickness in glaucoma: comparison of three OCT platforms. *Invest Ophthalmol Vis Sci*. 2015;56:6344–6351.
 12. Sung MS, Heo H, Park SW. Structure-function relationship in advanced glaucoma after reaching the RNFL floor. *J Glaucoma*. 2019;28:1006–1011.
 13. Hodapp E, Parrish RK, Anderson DR. *Clinical Decisions in Glaucoma*. Mosby; 1993.
 14. Stubeda H, Quach J, Gao J, et al. Comparing criteria for evaluating glaucomatous visual fields. *Am J Ophthalmol*. 2022;237:154–163.
 15. Corcoran Consulting Group. Medicare reimbursement for SCODI of the posterior segment (ZEISS). <https://www.corcoranccg.com/products/faqs/scodi-p-zeiss/>; 2019. Accessed August 25, 2022.
 16. Jammal AA, Thomson AC, Mariottoni EB, et al. Rates of glaucomatous structural and functional change from a large clinical population. *Am J Ophthalmol*. 2021;222:238–247.
 17. Shin JW, Sung KR, Song MK. Ganglion cell-inner plexiform layer and retinal nerve fiber layer changes in glaucoma suspects enable prediction of glaucoma development. *Am J Ophthalmol*. 2020;210:26–34.
 18. Chauhan BC, Garway-Heath DF, Goni FJ, et al. Practical recommendations for measuring visual field change in glaucoma. *Br J Ophthalmol*. 2008;92:569–573.

Pictures & Perspectives



Epithelial Demarcation Line after Crosslinking

A 14-year-old girl with bilateral keratoconus was evaluated in our outpatient clinic. Pentacam imaging revealed the thinnest pachymetry to be 380 μm . In view of the patient's age and thin pachymetry, a decision was made to proceed with collagen crosslinking (CXL) in the right eye. Accelerated CXL using customized epithelial debridement technique was performed, leaving an island of epithelium over the thinnest area of the cone. At 1 week, the epithelial island could be clearly differentiated from the new epithelium with a demarcation line in between (Fig A). Although there was no epithelial defect, the epithelial island had punctate staining with fluorescein (Fig B). (Magnified version of Fig A-B is available online at www.aaojournal.org).

RASHMI DESHMUKH, MS

VIVEK MAHENDRA SINGH, MS

Refractive and Cataract Services, LV Prasad Eye Institute, Hyderabad, India