



Histopathology and Ultrasound Biomicroscopy Correlation in Conjunctival Intraepithelial Neoplasia

An 83-year-old man with squamous cell carcinoma of the left conjunctiva presented for tumor recurrence. Slit-lamp examination showed an amelanotic lesion on the left bulbar conjunctiva extending from the 5 to 9 o'clock meridians involving fornix and caruncle (A). Prominent feeder and intrinsic vessels were visible (white arrows). Ultrasound biomicroscopy (UBM) revealed a solid, well-circumscribed lesion, measuring $1.4 \times 15 \times 9$ mm, without scleral invasion (B, C, longitudinal and transverse scan, respectively). Internally, multiple linear formations with high reflectivity (white arrows) were visible. Pathology confirmed a diagnosis of conjunctival intraepithelial neoplasia (CIN III). Large, dilated blood vessels (white arrow) were visible (D), consistent with the noted UBM findings. Therefore, UBM is a noninvasive aid in the diagnosis and management of CIN (Magnified version of Figure A-D is available online at www.aojournal.org).

JESSICA J. WANINGER, BS, BA¹

HAKAN DEMIRCI, MD²

BERNADETE AYRES, MD²

¹University of Michigan Medical School, Ann Arbor, Michigan; ²Department of Ophthalmology and Visual Sciences, Kellogg Eye Center, University of Michigan, Ann Arbor, Michigan

Footnotes and Disclosures

Supported by the Medical Scientist Training Program T32 (GM007863) and the University of Michigan Rackham Merit Fellowship (to J.W.W.) and Richard N. and Marilyn K. Witham professorship (to H.D.).