



### Lacrimal Gland Adenoid Cystic Carcinoma in a 7-year-old Boy

A 7-year-old boy presented with left orbital proptosis and globe dystopia (A); visual acuity and ophthalmic examination were normal. Orbital magnetic resonance imaging showed a lobular, well-defined lacrimal gland mass (B) without hypercellular features on an apparent diffusion coefficient map (C). Surgery was conducted with a primary diagnosis of lacrimal gland pleomorphic adenoma, revealing a lobular tumor with a gelatin-like texture and no capsule. A pathological examination revealed a tumor with cribriform (D, blue arrow) and canaliculalike (D, black arrow) characteristics, composed of inner ductal and outer myoepithelial cells. On immunostains, cytokeratin 7 highlighted all the cells (E, left), and calponin highlighted myoepithelial components (E, right). Fluorescent in situ hybridization results showed rearrangements of an *MYB* gene in 60% of the cells, confirming the rare diagnosis of lacrimal gland adenoid cystic carcinoma. Treatment included radiation therapy at the hospital's pediatric oncology clinic. (Magnified version of Figure A-E is available online at [www.aaojournal.org](http://www.aaojournal.org)).

TIRAN GOLANI, MD<sup>1,2</sup>  
GAHL GREENBERG, MD<sup>1,2</sup>  
GUY BEN SIMON, MD<sup>1,2</sup>

<sup>1</sup>Faculty of Medicine, Tel-Aviv University, Tel-Aviv, Israel; <sup>2</sup>Goldschleger Eye Institute, Sheba Medical Center, Tel-Hashomer, Israel