


24. Guidugli L, Mehine M, Tuupanen S, et al. Characterization of copy number variants (CNVs) identified by genetic testing of inherited retinal disorders. *Invest Ophthalmol Vis Sci.* 2019;60:2944.
25. Kong A, Gudbjartsson DF, Sainz J, et al. A high-resolution recombination map of the human genome. *Nat Genet.* 2002;31:241–247.
26. Kwon EM, Holt SK, Fu R, et al. Androgen metabolism and JAK/STAT pathway genes and prostate cancer risk. *Cancer Epidemiol.* 2012;36:347–353.
27. Kersten E, Geerlings MJ, Pauper M, et al. Genetic screening for macular dystrophies in patients clinically diagnosed with dry age-related macular degeneration. *Clin Genet.* 2018;94:569–574.
28. Lupski JR. Genomic disorders: structural features of the genome can lead to DNA rearrangements and human disease traits. *Trends Genet.* 1998;14:417–422.
29. Lee JA, Carvalho CM, Lupski JR. A DNA replication mechanism for generating nonrecurrent rearrangements associated with genomic disorders. *Cell.* 2007;131:1235–1247.
30. Zhang F, Khajavi M, Connolly AM, et al. The DNA replication FoSTeS/MMBIR mechanism can generate genomic, genic and exonic complex rearrangements in humans. *Nat Genet.* 2009;41:849–853.
31. Coe BP, Witherspoon K, Rosenfeld JA, et al. Refining analyses of copy number variation identifies specific genes associated with developmental delay. *Nat Genet.* 2014;46:1063–1071.
32. Perry GH, Yang F, Marques-Bonet T, et al. Copy number variation and evolution in humans and chimpanzees. *Genome Res.* 2008;18:1698–1710.

Pictures & Perspectives



Macular Star Associated with Fibrinous Central Serous Chorioretinopathy

A 44-year-old woman with systemic lupus erythematosus, receiving prednisolone, presented with a macular star and an oval spot surrounded by a ring-shaped subretinal fibrin (Fig A, arrowhead). The oval spot corresponded with the site of leak on fluorescein angiography (Fig B). OCT showed subretinal fluid, intraretinal hyperreflective foci, and a hyporeflective vacuole associated with the leakage site (Figs C, arrowhead, D). The patient was diagnosed with central serous chorioretinopathy. Severe fibrinous exudate not only accumulated in the subretinal space but also invaded the outer plexiform layer of the retina (yellow arrows), resulting in macular star formation (Magnified version of Fig A-D is available online at www.aaojournal.org).

CHIKAKO HARA, MD, PhD¹

TAKU WAKABAYASHI, MD, PhD^{1,2}

KOJI NISHIDA, MD, PhD¹

¹Department of Ophthalmology, Osaka University Graduate School of Medicine, Osaka, Japan; ²Wills Eye Hospital, Mid Atlantic Retina, Thomas Jefferson University, Philadelphia, Pennsylvania