


40. Gupta R, Sharma P, Menon V. A prospective clinical evaluation of augmented Anderson procedure for idiopathic infantile nystagmus. *J AAPOS*. 2006;10:312–317.
41. Graf M, Hausmann A, Lorenz B. High-dose Anderson operation for nystagmus-related anomalous head turn. *Graefes Arch Clin Exp Ophthalmol*. 2019;257:2033–2041.
42. Hertle RW, Felius J, Yang D, Kaufman M. Eye muscle surgery for infantile nystagmus syndrome in the first two years of life. *Clin Ophthalmol*. 2009;3:615–624.
43. ElKamshoushy A, Shawky D, ElMassry A, et al. Improved visual acuity and recognition time in nystagmus patients following four-muscle recession or Kestenbaum-Anderson procedures. *J AAPOS*. 2012;16:36–40.
44. Akbari MR, Akbari-Kamrani M, Mohseni A, et al. Effect of four horizontal rectus muscle tenotomy and replacement (TAR) alone and in combination with recessions for strabismus, on visual function and eye movements in patients with infantile nystagmus syndrome (INS) without abnormal head posture (AHP). *Binocul Vis Strabolog Q Simms Romano*. 2013;28:211–221.
45. Hertle RW, Curtis M, Boydston I, et al. Clinical and electrophysiological outcomes after eye muscle surgery in 81 adults with infantile nystagmus syndrome. *J Pediatr Ophthalmol Strabismus*. 2021;58:93–104.
46. Singh A, Ashar J, Sharma P, et al. A prospective evaluation of retroequatorial recession of horizontal rectus muscles and Hertle-Dell'Osso tenotomy procedure in patients with infantile nystagmus with no definite null position. *J AAPOS*. 2016;20:96–99.
47. Dell'Osso LF, Hertle RW, Jacobs JB. Clinical and ocular motor complications of extraocular muscle extirpation for infantile nystagmus syndrome. *J AAPOS*. 2018;22:110–114 e111.

Pictures & Perspectives



Concurrent Mucocoele and Silent Sinus Syndrome after Orbital Fracture Repair

A 59-year-old man sustained a left medial wall-floor fracture requiring repair (Fig A). Eight years later, the patient presented with left-sided proptosis and binocular diplopia. Visual acuity was normal with no afferent pupillary defect. Extraocular movements were limited in supraduction and abduction in the left eye. Orbital computed tomography demonstrated a large soft tissue density representing a mucocoele encasing the implant (Fig B). There was also opacification of the left maxillary sinus and bowing of the orbital floor consistent with silent sinus syndrome. Although rare, a mucocoele may result in silent sinus syndrome by obstruction of the osteomeatal complex leading to maxillary sinus atelectasis (Magnified version of Fig A-B is available online at www.aaojournal.org).

BENYAM KINDE, MD, PhD

MELEHA AHMAD, MD, MS

M. REZA VAGEFI, MD

Department of Ophthalmology, University of California, San Francisco, San Francisco, California

Footnotes and Disclosures

Funded in part by an institutional grant from Research to Prevent Blindness.