


41. de Groot S, Röttgering B, Gelderblom H, et al. Unraveling the resistance of IGF-pathway inhibition in Ewing sarcoma. *Cancers (Basel)*. 2020;12(12):3568.
42. Haluska P, Shaw HM, Batzel GN, et al. Phase I dose escalation study of the anti insulin-like growth factor-I receptor monoclonal antibody CP-751,871 in patients with refractory solid tumors. *Clin Cancer Res*. 2007;13(19):5834–5840.
43. Shah K, Charitou M. A novel case of hyperglycemic hyperosmolar state after the use of teprotumumab in a patient with thyroid eye disease. *AACE Clin Case Rep*. 2022;8(4):148–149.
44. Amariqwa L, Mohamed A, Kim SH, et al. Teprotumumab-related hyperglycemia. *J Clin Endocrinol Metab*. 2023;108(4):858–864.
45. Douglas RS, Wang Y, Dailey RA, et al. Teprotumumab in clinical practice: recommendations and considerations from the OPTIC Trial Investigators. *J Neuroophthalmol*. 2021;41:461–468.
46. Lee HBH, Mariash CN, Yoon MK, et al. Teprotumumab and hyperglycemia guidelines to monitor for hyperglycemia in teprotumumab. *Ophthalmic Plast Reconstr Surg*. 2021;37(4):393.
47. Kossler AL, Douglas R, Dosiou C. Teprotumumab and the evolving therapeutic landscape in thyroid eye disease. *J Clin Endocrinol Metab*. 2022;107(Suppl 1):S36–S46.

## Pictures & Perspectives



### Neglected Metastatic Neuroblastoma

A 5-year-old child was referred to our clinic with a history of neglected and untreated metastatic neuroblastoma. **A**, The child exhibited severe proptosis, severe chemosis, exposure keratopathy, corneal ulcers, limited extraocular movements, and decreased visual acuity in both eyes. **B**, Cerebrospinal fluid analysis showed a cluster of neuroblastoma cells. The child underwent transconjunctival orbital decompression surgery to alleviate proptosis and relieve pressure on the optic nerve. Additionally, intrathecal chemotherapy was administered to target central nervous system involvement. **C**, The child's response to treatment was remarkable, as the proptosis subsided, and extraocular movements returned to normal. Although corneal scarring remains (**D**), there has been an improvement in visual acuity. (Magnified version of Figure A–D is available online at [www.aaojournal.org](http://www.aaojournal.org)).

KOUROSH SHAHRAKI, MD  
DONNY W. SUH, MD, MBA

Department of Ophthalmology, Gavin Herbert Eye Institute, University of California Irvine, Irvine, California