

Ophthalmic Images

Live Larva in Ocular Toxocariasis

Sara Geada, MD; Cristina Fonseca, MD; Rui Proença, MD, PhD




Figure. Live larva in ocular toxocariasis. A, Ultra-widefield color fundus photograph with amplification cuts of consecutive 4-month interval photographs (bottom) demonstrating a coiling/uncoiling image compatible with a live nematode. B, Optical coherence tomography showing a hyperreflective image (arrowhead; transversal cut) in the subretinal space.

A 23-year-old man presented for follow-up of ocular toxocariasis (OT), diagnosed 5 years prior during an evaluation for vitritis and chorioretinal granuloma. Diagnosis was confirmed through serum and



CME at jamacmelookup.com

aqueous humor *Toxocara canis* antibodies, and treatment with albendazole and oral steroids was instituted. Treatment resulted in inflammation improvement; however, macular scarring led to permanent vision loss. Ultra-widefield color fundus photographs taken at consecutive 4-month intervals revealed a linear

image (Figure, A) compatible with a live larva. This was corroborated by optical coherence tomography as a hyperreflective element in the subretinal space of the detached retina (Figure, B). Diagnosis of OT is generally presumptive, with larva demonstration occurring only in rare cases.^{1,2} Treatment remains controversial; options include albendazole with adjunctive photocoagulation (impossible in this case due to detached retina) or vitrectomy.^{3,4} Considering the stable cicatricial findings and the poor visual prognosis, it was decided to withhold any course of treatment and closely monitor the patient for recurrence.

ARTICLE INFORMATION

Author Affiliations: Department of Ophthalmology, Centro Hospitalar e Universitário de Coimbra, Coimbra, Portugal (Geada, Fonseca, Proença); Clinical Academic Centre of Coimbra, Coimbra, Portugal (Geada, Fonseca, Proença).

Corresponding Author: Sara Geada, MD, Department of Ophthalmology, Centro Hospitalar e Universitário de Coimbra, Praceta Professor Mota Pinto, 3004-561 Coimbra, Portugal (sarageadabatista@gmail.com).

Conflict of Interest Disclosures: None reported.

Additional Contributions: We thank the patient for granting permission to publish this information.

REFERENCES

1. Fonseca C, Silva AM, Freire S, Proença R. Ocular toxocariasis: atypical clinical course. *BMJ Case Rep*. 2019;12(4):e228717. doi:10.1136/bcr-2018-228717
2. Hernanz I, Moll-Udina A, Llorenç BV, Adan CA. Ocular toxocariasis: beyond typical patterns through the new imaging technologies. *Ocul*

Immunol Inflamm. 2021;29(7-8):1252-1258. doi:10.1080/09273948.2020.1793370

3. Ahn SJ, Ryoo NK, Woo SJ. Ocular toxocariasis: clinical features, diagnosis, treatment, and prevention. *Asia Pac Allergy*. 2014;4(3):134-141. doi:10.5415/apallergy.2014.4.3.134

4. Despreaux R, Fardeau C, Touhami S, et al. Ocular toxocariasis: clinical features and long-term visual outcomes in adult patients. *Am J Ophthalmol*. 2016;166:162-168. doi:10.1016/j.ajo.2016.03.050