
- 38. Tomita G, Araie M, Kitazawa Y, Tsukahara S. A three-year prospective, randomized and open comparison between latanoprost and timolol in Japanese normal-tension glaucoma patients. *Eye.* 2004;18:984—989.
- **39.** Lee JW, Ho WL, Chan JC, Lai JS. Efficacy of selective laser trabeculoplasty for normal tension glaucoma: 1 year results. *BMC Ophthalmol*. 2015;15:1.
- Lee JWY, Shum JJW, Chan JCH, Lai JSM. Two-year clinical results after selective laser trabeculoplasty for normal tension glaucoma. *Medicine*. 2015;94:e984.
- 41. Lee JWY, Liu CCL, Chan JCH, Lai JSM. Predictors of success in selective laser trabeculoplasty for chinese open-angle glaucoma. *J Glaucoma*. 2014;23:321–325.
- Wallace DK, Melia M. Post hoc power calculations. *Ophthalmology*. 2008;115:2098. author reply 2098–2099.
- 43. Wright DM, Konstantakopoulou E, Montesano G, et al. Visual field outcomes from the multicenter, randomized controlled Laser in Glaucoma and Ocular Hypertension Trial (LiGHT). *Ophthalmology*. 2020;127:1313—1321.

## **Pictures & Perspectives**



## Ophthalmic Sequelae of Late-Stage Primary Hyperoxaluria Type 1

A 56-year-old woman with primary hyperoxaluria type 1 presented with count fingers vision and significantly elevated intraocular pressures bilaterally. Examination revealed bilateral angle closure due to neovascular peripheral anterior synechiae (A), advanced optic disc cupping, severe oxalate retinopathy, and attenuated, crystal-filled vessels (B). OCT of the macula showed crystal deposition in every retinal layer and hyperreflectivity and irregularity of the retinal pigment epithelium with overlying loss of the ellipsoid zone (C). Fluorescein angiography demonstrated widespread vascular leakage (D). Primary hyperoxaluria type 1 is an extremely rare inborn error of metabolism characterized by systemic deposition of oxalate. The authors report a unique case of oxalate-compromised vasculature leading to the development of neovascularization and subsequent angle-closure glaucoma. (Magnified version of Figure A-D is available online at www.aaojournal.org).

KIMBERLY NGUYEN, MD LYNN SUN, MD, PHD

Department of Ophthalmology, Casey Eye Institute, Oregon Health & Science University, Portland, Oregon