

EVERY n POINTS IN THE PLANE CONTAIN $(m_1 + o(1))n$ POINTS WITH NO UNIT DISTANCE

SHISHENG LI

ABSTRACT. Let $f(n)$ be the largest integer such that every set of n points in the plane contains $f(n)$ points no two at distance 1, and let m_1 be the supremum of the upper densities of Lebesgue measurable planar sets with no two points at distance 1. Larman and Rogers observed that $f(n) \geq m_1 n$. We prove that $f(n)/n \rightarrow m_1$; that is, $f(n) = (m_1 + o(1))n$. This is the first half of a conjecture of Dúcz and Varga, and with the bound $m_1 \leq 0.247$ of Ambrus, Csiszárík, Matolcsi, Varga and Zsámboki it gives $f(n) \leq (0.247 + o(1))n$. The proof builds a random independent subset of the algebraic plane, invariant under algebraic translations and rotations, whose probability of containing the origin is at least the worst finite independence ratio, and turns it into a measurable planar set of at least that upper density. Its only non-classical input is the Haar rigidity theorem for wild character laws from the OpenAI preprint *The Euclidean plane is not five-colorable*, which has a machine-checked Lean proof. The whole argument has been formalized in Lean 4 on top of it.

1. INTRODUCTION

For a finite set $P \subset \mathbb{R}^2$ let G_P be its unit-distance graph (vertices P , edges the pairs at Euclidean distance exactly 1) and $\alpha(P)$ its independence number. Erdős [7] asked to estimate

$$f(n) = \min\{\alpha(P) : P \subset \mathbb{R}^2, |P| = n\},$$

and in particular whether $f(n) \geq n/4$ (Problem #1070 in Bloom's database [4]). The Moser spindle gives $f(n) \leq \frac{2}{7}n + O(1)$. Matolcsi, Ruzsa, Varga and Zsámboki [9] proved $f(n) \leq (\frac{1}{4} + o(1))n$ and conjectured that this is sharp. Dúcz and Varga [6] disproved that conjecture by exhibiting a finite unit-distance graph with independence ratio below $1/4$, answering the second question of Erdős negatively; their result has been formalized in Lean by Benkő [3].

For the lower bound, let m_1 be the supremum of the upper densities of measurable sets $A \subset \mathbb{R}^2$ containing no two points at distance 1. Larman and Rogers [8] observed $f(n) \geq m_1 n$; Croft [5] constructed such a set of density $\delta_{\text{Croft}} = 0.22936\dots$, so $m_1 \geq \delta_{\text{Croft}}$; and Ambrus, Csiszárík, Matolcsi, Varga and Zsámboki [1] proved $m_1 \leq 0.2470$, settling Erdős's question whether $m_1 < 1/4$ (Problem #232 of [4]). Dúcz and Varga [6, §2.1] state: “*Conjecture 1. $f(n)/n = m_1(\mathbb{R}^2) + o(1)$ and in particular that $m_1(\mathbb{R}^2) = \delta_{\text{Croft}}$.*” We prove the first half, $f(n)/n = m_1 + o(1)$.

Theorem 1.1. $\lim_{n \rightarrow \infty} f(n)/n = m_1$. Moreover $f(n) \geq m_1 n$ for every $n \geq 1$.

Corollary 1.2. $0.22936 n \leq f(n) \leq (0.2470 + o(1)) n$. In particular $f(n) < n/4$ for all large n .

Theorem 1.1 says that the two natural constants of the problem, the worst finite independence ratio and the best measurable density, coincide; Matolcsi, Ruzsa, Varga and Zsámboki [9, Conjecture 1] had conjectured that the finite independence ratio is never below $1/4$, which together with $m_1 < 1/4$ would have made the two constants differ. Determining the numerical value of m_1 remains open.

Write

$$\alpha^* = \inf \left\{ \frac{\alpha(P)}{|P|} : P \subset \mathbb{R}^2 \text{ finite, nonempty} \right\}.$$

Theorem 1.1 follows from three statements: $f(n)/n \rightarrow \alpha^*$ with $f(n) \geq \alpha^*n$ (Section 3); $\alpha^* \geq m_1$ (Larman–Rogers, Section 4); and $m_1 \geq \alpha^*$ (Section 5). The last is the new part. It is a single-label version of the transfer in Section 4 of the OpenAI preprint [10], which turns an arbitrary proper colouring into a measurable weak colouring; the one change needed is in how the averaging is done (Remark 5.4).

Formalization. All of the above is formalized in Lean 4 with Mathlib, building on OpenAI’s formalization [11] of the five-colour paper; see Section 6.

2. DENSITIES

For a measurable $A \subset \mathbb{R}^2$ and a family of regions $(W_r)_r$ put

$$\bar{\delta}_W(A) = \limsup_{r \rightarrow \infty} \frac{|A \cap W_r|}{|W_r|},$$

where $|\cdot|$ is Lebesgue measure. We consider four families: discs $B(0, r)$ or squares $[-r, r]^2$, with r ranging over $\mathbb{R}_{>0}$ or over $\mathbb{N}_{\geq 1}$; below $\mathbb{N} = \{1, 2, \dots\}$. Let m_1^W be the supremum of $\bar{\delta}_W(A)$ over measurable unit-distance-free A . The database [4] uses discs with real r .

Remark 2.1. All four values m_1^W equal α^* , so m_1 is unambiguous. Indeed, Proposition 4.1 below gives $m_1^W \leq \alpha^*$ for each family, and Section 5 constructs, for discs and for squares, a measurable unit-distance-free A with $\limsup_{r \in \mathbb{N}} |A \cap W_r|/|W_r| \geq \alpha^*$; a limsup over $r \in \mathbb{R}$ is at least the limsup over $r \in \mathbb{N}$.

3. CONVERGENCE OF $f(n)/n$

Proposition 3.1. $f(n) \geq \alpha^*n$ for every n , and $f(n)/n \rightarrow \alpha^*$.

Proof. If $|P| = n$ then $\alpha(P) \geq \alpha^*n$ by definition; hence $f(n) \geq \alpha^*n$. Fix a finite P_0 with $|P_0| = s$ and $\alpha(P_0) = a$. For $N \geq s$ write $N = ks + r$ with $0 \leq r < s$. Take k translates of P_0 and r single points, placed so that distinct pieces are at distance more than $\text{diam } P_0 + 2$. The union has N points, its unit-distance graph is the disjoint union of the pieces’ graphs, and its independence number is $ka + r$. Hence $f(N) \leq ka + r \leq Na/s + s$, so $\limsup_N f(N)/N \leq a/s$. Taking the infimum over P_0 gives $\limsup_N f(N)/N \leq \alpha^*$. $\square$

4. THE BOUND $\alpha^* \geq m_1$

Proposition 4.1 (Larman–Rogers [8]). *Let A be measurable and unit-distance-free, and let W be any of the four families. Then $\alpha(P) \geq \bar{\delta}_W(A) |P|$ for every finite nonempty P . Hence $m_1^W \leq \alpha^*$.*

Proof. Let $P = \{p_1, \dots, p_n\}$, $\rho = \max_j |p_j|$, $\delta = \bar{\delta}_W(A)$. For each x , $(P+x) \cap A$ is an independent set of $G_{P+x} \cong G_P$, so $|(P+x) \cap A| \leq \alpha(P)$. By Tonelli,

$$\alpha(P) |W_r| \geq \int_{W_r} |(P+x) \cap A| dx = \sum_{j=1}^n |A \cap (W_r + p_j)| \geq n|A \cap W_r| - \sum_{j=1}^n |W_r \setminus (W_r + p_j)|.$$

For a disc, $W_r \setminus (W_r + p)$ lies in the annulus $\max(0, r - |p|) \leq |x| \leq r$, of area $\pi r^2 - \pi \max(0, r - |p|)^2 \leq 2\pi r|p|$; for a square it lies in two strips of total area at most $2r(|p_1| + |p_2|) \leq 2\sqrt{2}r|p|$. In both cases $|W_r \setminus (W_r + p)| \leq 2\pi r|p|$ and $|W_r| \geq \pi r^2$, so $\alpha(P) \geq n|A \cap W_r|/|W_r| - 2n\rho/r$. Letting $r \rightarrow \infty$ along a sequence realizing the limsup gives $\alpha(P) \geq n\delta$. $\square$

5. THE BOUND $m_1 \geq \alpha^*$

5.1. Setting. Identify $\mathbb{R}^2$ with $\mathbb{C}$. Let $E = \overline{\mathbb{Q}} \subset \mathbb{C}$ be the field of algebraic numbers, viewed as a countable discrete additive group, and $K = \{u \in E : |u| = 1\}$, a countable discrete multiplicative group. For $x, y \in E$ we have $|x - y| = 1$ iff $x - y \in K$. K is dense in the unit circle S^1 , since $\frac{(1-t^2)+2it}{1+t^2} \in K$ for $t \in \mathbb{Q}$.

Let $D = \widehat{E}$ be the compact (metrizable, as E is countable) dual group of E ; for $a, z \in E$ and $d \in D$ put $(ad)(z) = d(az)$ and $\psi_z(d) = d(z)$. The Euclidean-continuous characters (continuous for the topology E inherits from $\mathbb{C}$) $C = \{j(\xi) : \xi \in \mathbb{R}^2\}$, $j(\xi)(z) = e^{i\xi \cdot z}$, form a subgroup of D . It is Borel: D is a compact metrizable group, $j : \mathbb{R}^2 \rightarrow D$ is a continuous injective homomorphism (injectivity can be tested on $z \in \mathbb{Q} \cup i\mathbb{Q}$), so $C = \bigcup_{m \geq 1} j([-m, m]^2)$ is σ -compact, hence Borel.

Theorem 5.1 (Haar rigidity [10, Thm. 2.3]). *Every K -invariant Borel probability measure ν on D with $\nu(C) = 0$ is Haar measure. Equivalently (by uniqueness of Fourier coefficients on D), $\int_D \psi_z d\nu = 0$ for all $z \in E \setminus \{0\}$.*

The second form is proved in Lean as `OAI.PlaneFiveColor.Spectral.wild_fourier` [11]. We also use the following lemma, whose proof in [10] is a standard spectral argument (spectral theorem for the discrete group E , Bochner's theorem on $\mathbb{R}^2$, uniqueness of Fourier coefficients on D).

Lemma 5.2 ([10, Lemma 4.1]). *Let E act by measure-preserving maps on a standard probability space (Ω, μ) , with Koopman operators T_z on $L^2(\mu)$. Let $\mathcal{H}_c = \{v \in L^2(\mu) : \|T_z v - v\|_2 \rightarrow 0 \text{ as } z \rightarrow 0, z \in E\}$, the limit in the topology of $\mathbb{C}$. Then:*

- (i) $\mathcal{H}_c = L^2(\Omega, \mathcal{F}_c, \mu)$ for a sub- σ -algebra $\mathcal{F}_c$, so the orthogonal projection onto $\mathcal{H}_c$ is $\mathbb{E}(\cdot | \mathcal{F}_c)$;
- (ii) $\mathcal{H}_c$ is the range of the spectral projection of $\{T_z\}$ on C ;
- (iii) the restrictions $T_z|_{\mathcal{H}_c}$ extend to a strongly continuous unitary representation $(T_x)_{x \in \mathbb{R}^2}$ on $\mathcal{H}_c$ preserving the cone of nonnegative functions and fixing $\mathbf{1}$;
- (iv) if unitaries V_u ($u \in K$) satisfy $V_u T_z V_u^{-1} = T_{uz}$, then $V_u \mathcal{H}_c = \mathcal{H}_c$ and V_u commutes with the projection onto $\mathcal{H}_c$.

5.2. An invariant random independent set. Let $\Omega \subset \{0, 1\}^E$ be the set of independent sets of the unit-distance graph on E . Each edge constraint is clopen, so Ω is a closed subset of the compact metrizable space $\{0, 1\}^E$. Translations $s_z \omega = \omega - z$ ($z \in E$) and rotations $\rho_u \omega = u\omega$ ($u \in K$) are homeomorphisms of Ω , and $s_w \rho_u = \rho_u s_{u^{-1}w}$.

We use Følner sequences in E and K : finite sets F_N with $|F_N \Delta (F_N + w)|/|F_N| \rightarrow 0$ for every w (and the multiplicative analogue in K). They exist because E and K are countable abelian groups, hence amenable [2, Thm. G.5.1]. For E one can take explicitly: enumerate $E = \{e_0, e_1, \dots\}$, let $\Lambda_N \cong \mathbb{Z}^{d_N}$ be the subgroup generated by $e_0, \dots, e_N$ (E is torsion-free), fix a basis of Λ_N , and let F_N be the box of side L_N in these coordinates, with L_N so large that $|F_N \Delta (F_N + e_j)|/|F_N| < 1/N$ for $0 \leq j \leq N$. Since every $w \in E$ is an integer combination of finitely many e_j , this gives the Følner property for every w .

Lemma 5.3. *There is a Borel probability measure μ on Ω , invariant under all s_z and ρ_u , with $\mu(\{\omega : 0 \in \omega\}) \geq \alpha^*$.*

Proof. The space $\mathcal{P}(\Omega)$ of Borel probability measures is weak-* compact and metrizable (Riesz–Markov and Banach–Alaoglu), and for a homeomorphism g of Ω the map $\nu \mapsto g_* \nu$ is weak-* continuous. The set $\{0 \in \omega\}$ is clopen, so $\nu \mapsto \nu(0 \in \omega)$ is weak-* continuous.

Translations. Each F_N is a finite subset of $\mathbb{R}^2$; choose a maximum independent set $J_N \subset F_N$, so $|J_N| \geq \alpha^* |F_N|$ and $J_N - z \in \Omega$ for all z . Let $\nu_N = |F_N|^{-1} \sum_{z \in F_N} \delta_{J_N - z}$. Then $\nu_N(0 \in \omega) =$

$|J_N|/|F_N| \geq \alpha^*$ and $\|(s_w)_*\nu_N - \nu_N\|_{\text{TV}} \leq |F_N \triangle (F_N + w)|/|F_N| \rightarrow 0$. Any weak-* limit point μ_0 is therefore s_w -invariant for every w and satisfies $\mu_0(0 \in \omega) \geq \alpha^*$.

Rotations. Let G_M be Følner sets in K and $\mu_M = |G_M|^{-1} \sum_{u \in G_M} (\rho_u)_* \mu_0$. Each $(\rho_u)_* \mu_0$ is translation invariant, because $(s_w)_*(\rho_u)_* \mu_0 = (\rho_u)_*(s_{u^{-1}w})_* \mu_0 = (\rho_u)_* \mu_0$; and $(\rho_u)_* \mu_0(0 \in \omega) = \mu_0(0 \in \omega)$ since $0 \in u\omega \iff 0 \in \omega$. A weak-* limit point μ of (μ_M) is translation invariant, rotation invariant by the Følner property, and has $\mu(0 \in \omega) \geq \alpha^*$. $\square$

Remark 5.4. In [10, Prop. 4.2] a single colouring is averaged over Følner sets of $E \rtimes K$. Applied to a fixed maximum independent set J of a finite set and a finite averaging set $\Phi \subset E \rtimes K$, the empirical measure $\nu_\Phi = |\Phi|^{-1} \sum_{g \in \Phi} \delta_{gJ}$ has $\nu_\Phi(0 \in \omega) = \sum_w c(w) \mathbf{1}_J(w) / \sum_w c(w)$ with multiplicities $c(w) = \#\{g \in \Phi : g^{-1}0 = w\}$ that are not constant in w , i.e. the c -weighted proportion of J , for which no lower bound α^* is available a priori. Averaging first over translations (where each position is counted once) and then over rotations (which fix 0) avoids this.

5.3. Projection onto the continuous factor. Fix μ from Lemma 5.3. Since Ω is compact metrizable, (Ω, μ) is a standard probability space. Define unitaries on $L^2(\mu)$ by $T_z F(\omega) = F(\omega - z)$ and $V_u F(\omega) = F(u^{-1}\omega)$; then $T_z T_w = T_{z+w}$ and $V_u T_z V_u^{-1} = T_{uz}$. Let $f = \mathbf{1}[0 \in \omega]$. Then $T_z f = \mathbf{1}[z \in \omega]$, $V_u f = f$, $\mathbb{E}f \geq \alpha^*$, and since 0 and $u \in K$ are adjacent, $f \cdot T_u f = 0$ pointwise, so

$$(1) \quad \langle f, T_u f \rangle = 0 \quad (u \in K).$$

Let $p = \mathbb{E}(f \mid \mathcal{F}_c) \in \mathcal{H}_c$ and $b = f - p \in \mathcal{H}_c^\perp$ (Lemma 5.2). Then $0 \leq p \leq 1$ a.e., $\mathbb{E}p = \mathbb{E}f$, and $V_u p = p$, $V_u b = b$. Both p and b are real-valued.

Let λ_b be the spectral measure of b on D : $\langle b, T_z b \rangle = \int_D \psi_z d\lambda_b$. By Lemma 5.2(ii), $\lambda_b(C) = 0$. For $u \in K$,

$$\langle b, T_{uz} b \rangle = \langle b, V_u T_z V_u^{-1} b \rangle = \langle V_u^{-1} b, T_z V_u^{-1} b \rangle = \langle b, T_z b \rangle,$$

With $m_u : D \rightarrow D$, $m_u(d)(z) = d(uz)$, we have $\int_D \psi_z d((m_u)_* \lambda_b) = \int_D \psi_{uz} d\lambda_b$, so $(m_u)_* \lambda_b$ and λ_b have the same Fourier coefficients, and λ_b is K -invariant. If $b \neq 0$, Theorem 5.1 applied to $\lambda_b/\|b\|^2$ gives $\langle b, T_z b \rangle = 0$ for $z \neq 0$; this is trivial if $b = 0$. Since T_u preserves $\mathcal{H}_c$ and $\mathcal{H}_c^\perp$, $\langle p, T_u b \rangle = \langle b, T_u p \rangle = 0$. Expanding (1),

$$\langle p, T_u p \rangle = 0 \quad (u \in K).$$

By Lemma 5.2(iii), $(T_x)_{x \in \mathbb{R}^2}$ is a strongly continuous unitary representation on $\mathcal{H}_c \ni p$, so $u \mapsto \langle p, T_u p \rangle$ is continuous and, K being dense in S^1 , vanishes on S^1 . Hence for all $x \in \mathbb{R}^2$ and $u \in S^1$,

$$(2) \quad \int (T_x p)(T_{x+u} p) d\mu = \langle T_x p, T_{x+u} p \rangle = \langle p, T_u p \rangle = 0.$$

Moreover $0 \leq T_x p \leq 1$ a.e. (apply positivity to p and $\mathbf{1} - p$) and $\mathbb{E}[T_x p] = \langle T_x p, \mathbf{1} \rangle = \langle p, T_{-x} \mathbf{1} \rangle = \mathbb{E}p \geq \alpha^*$.

5.4. A measurable field and one sample.

Lemma 5.5. *Let $x \mapsto v_x$ be a continuous map from $\mathbb{R}^2$ to the real Hilbert space $L^2(\mu; \mathbb{R})$. There is a jointly measurable $H : \mathbb{R}^2 \times \Omega \rightarrow \mathbb{R}$ with $H(x, \cdot) = v_x$ μ -a.e. for every $x \in \mathbb{R}^2$.*

Proof. Fix a dense sequence (x_k) in $\mathbb{R}^2$ and a representative of each v_{x_k} . For $j \geq 1$ let $k_j(x)$ be the least k with $\|v_{x_k} - v_x\|_2 < 2^{-j}$; it exists by continuity and density, and $x \mapsto k_j(x)$ is Borel because each $x \mapsto \|v_{x_k} - v_x\|_2$ is continuous. Put $H_j(x, \omega) = v_{x_{k_j(x)}}(\omega) = \sum_k \mathbf{1}[k_j(x) = k] v_{x_k}(\omega)$, which is jointly measurable as a countable sum of products of measurable functions. For fixed x , $\sum_j \|H_j(x, \cdot) - v_x\|_1 \leq \sum_j \|H_j(x, \cdot) - v_{x_k}\|_2 < \sum_j 2^{-j} < \infty$ (μ is a probability measure), so

by Tonelli $\sum_j |H_j(x, \cdot) - v_x| < \infty$ a.e., and $H_j(x, \cdot) \rightarrow v_x$ a.e. Let $H = \limsup_j H_j$ where finite and 0 elsewhere. $\square$

Apply Lemma 5.5 to the real-valued $v_x = T_x p$ and replace H by $\min(\max(H, 0), 1)$. Fix the family W (discs or squares) and integer radii $r \in \mathbb{N}$. By (2) and Tonelli, for every $r \in \mathbb{N}$,

$$\int_{\Omega} \int_{B(0,r)} \int_{S^1} H(x, \omega) H(x+u, \omega) d\sigma(u) dx d\mu(\omega) = 0,$$

where σ is normalized arc-length measure on S^1 , and the inner double integral is nonnegative, so it vanishes for μ -a.e. ω ; intersecting over the countably many $r \in \mathbb{N}$ gives a μ -conull set Ω_1 on which it vanishes for every $r \in \mathbb{N}$. Let $X_r(\omega) = |W_r|^{-1} \int_{W_r} H(x, \omega) dx \in [0, 1]$. By Tonelli, $\mathbb{E} X_r = \mathbb{E} p \geq \alpha^*$, and by the reverse Fatou lemma ($X_r \leq 1$), $\mathbb{E}[\limsup_r X_r] \geq \alpha^*$; hence $\mu(\limsup_r X_r \geq \alpha^*) > 0$ (otherwise $\alpha^* - \limsup_r X_r > 0$ a.e., so its integral would be positive). Fix $\omega \in \Omega_1$ with $\limsup_r X_r(\omega) \geq \alpha^*$ and set $h = H(\cdot, \omega) : \mathbb{R}^2 \rightarrow [0, 1]$:

$$(3) \quad \int_{B(0,r)} \int_{S^1} h(x) h(x+u) d\sigma dx = 0 \quad (r \in \mathbb{N}), \quad \limsup_{r \in \mathbb{N}} \frac{1}{|W_r|} \int_{W_r} h \geq \alpha^*.$$

5.5. Density points. Let $A_0 = \{h > 0\}$. By (3) the nonnegative function $h(x)h(x+u)$ vanishes for $dx \otimes \sigma$ -a.e. $(x, u) \in B(0, r) \times S^1$, for every r ; since $\mathbf{1}_{A_0}(x)\mathbf{1}_{A_0}(x+u) > 0$ iff $h(x)h(x+u) > 0$, also $\int_{B(0,r)} \int_{S^1} \mathbf{1}_{A_0}(x)\mathbf{1}_{A_0}(x+u) d\sigma dx = 0$ for all r , and since $h \leq \mathbf{1}_{A_0}$, $\limsup_{r \in \mathbb{N}} |A_0 \cap W_r|/|W_r| \geq \alpha^*$. Let A be the set of Lebesgue density points of A_0 . It is Borel, and $|A \Delta A_0| = 0$ by the Lebesgue density theorem, so A has the same densities as A_0 .

Lemma 5.6. *A contains no two points at distance 1.*

Proof. Suppose $x, y = x + u \in A$ with $|u| = 1$. For $0 < \eta < 1$ let x' be uniform on $B(x, \eta)$ and, independently, u' uniform on the arc $\{e^{i\theta}u : |\theta| < \eta/2\}$, so $|u' - u| < \eta$. Then $\mathbb{P}(x' \notin A_0) = |B(x, \eta) \setminus A_0|/|B(x, \eta)| \rightarrow 0$. Given u' , $x' + u'$ is uniform on $B(x + u', \eta) \subset B(y, 2\eta)$, so $\mathbb{P}(x' + u' \notin A_0) \leq |B(y, 2\eta) \setminus A_0|/(\pi\eta^2) = 4|B(y, 2\eta) \setminus A_0|/|B(y, 2\eta)| \rightarrow 0$. So for small η , $\mathbb{P}(x' \in A_0, x' + u' \in A_0) > 0$. The law of (x', u') is absolutely continuous with respect to $dx \otimes \sigma$ and is supported in $B(0, r) \times S^1$ for an integer $r > |x| + 1$. Hence the set $\{(x', u') : x' \in A_0, x' + u' \in A_0\} \cap (B(0, r) \times S^1)$ has positive $dx \otimes \sigma$ -measure, contradicting $\int_{B(0,r)} \int_{S^1} \mathbf{1}_{A_0}(x)\mathbf{1}_{A_0}(x+u) d\sigma dx = 0$. $\square$

Thus A is a measurable unit-distance-free set with $\limsup_{r \in \mathbb{N}} |A \cap W_r|/|W_r| \geq \alpha^*$, for W = discs and for W = squares. This proves $m_1^W \geq \alpha^*$ for all four conventions, completing Remark 2.1 and, with Propositions 3.1 and 4.1, Theorem 1.1. Corollary 1.2 follows from [5, 1].

6. FORMALIZATION

The repository [12] contains a Lean 4 project (toolchain v4.34.1, Mathlib d13f23b). The directory `OAI/` is a verbatim copy of the 70-file import closure of OpenAI's five-colour theorem from [11] (Apache-2.0). Our files define

- `indepNum P` (independence number of the unit-distance graph on a `Finset C`),
- `f n`, the minimum of `indepNum P` over n -element P ,
- `upperDensity` (squares $[-m, m]^2$, $m \in \mathbb{N}$) and `m1`,

and prove `f_div_tendsto_m1` ($f(n)/n \rightarrow m_1$) and `m1_mul_le_f` ($m_1 n \leq f(n)$). The file `Disc.lean` defines `m1Ball` with open discs of real radius (the convention of [4]) and proves `f_div_tendsto_m1Ball`, `m1Ball_mul_le_f` and `m1Ball_eq_m1`. `#print axioms` reports only `propext`, `Classical.choice` and `Quot.sound` for all of these and for `wild_fourier`. The Lean proof follows Sections 3–5; the Følner sets in E are constructed explicitly as above, and the

weak-* limits are realized by ultrafilter limits of positive functionals and the Riesz–Markov theorem.

ACKNOWLEDGEMENTS

The negative answer to the second question of Erdős is due to Dúcz and Varga [6]. Theorem 5.1 and the transfer template are from [10]. AI tools were used in this work: the observation that [10, Section 4] applies to a single colour class, the write-up and the Lean formalization were produced with Claude (Anthropic), and the manuscript was audited with GPT (OpenAI).

REFERENCES

- [1] G. Ambrus, A. Csiszárík, M. Matolcsi, D. Varga, P. Zsámboki, *The density of planar sets avoiding unit distances*, Math. Program. (2023), doi:10.1007/s10107-023-02012-9; arXiv:2207.14179.
- [2] B. Bekka, P. de la Harpe, A. Valette, *Kazhdan’s Property (T)*, Cambridge Univ. Press, 2008.
- [3] B. Benkő, Lean formalization of [6], <https://github.com/bbeatrix/unit-distance-graph-independence-ratio>.
- [4] T. F. Bloom, *Erdős Problems*, <https://www.erdosproblems.com>, Problems #232 and #1070.
- [5] H. T. Croft, *Incidence incidents*, Eureka **30** (1967), 22–26.
- [6] Á. Dúcz, D. Varga, *A unit-distance graph in the plane with independence ratio below 1/4*, arXiv:2606.28157 (2026).
- [7] P. Erdős, *Some combinatorial and metric problems in geometry*, in: Intuitive Geometry (Siófok, 1985), Colloq. Math. Soc. János Bolyai **48**, North-Holland, 1987, 167–177.
- [8] D. G. Larman, C. A. Rogers, *The realization of distances within sets in Euclidean space*, Mathematika **19** (1972), 1–24.
- [9] M. Matolcsi, I. Z. Ruzsa, D. Varga, P. Zsámboki, *The fractional chromatic number of the plane is at least 4*, arXiv:2311.10069.
- [10] OpenAI, *The Euclidean plane is not five-colorable*, preprint, 23 September 2026, <https://github.com/openai/math/tree/adc7f1241b42e322a6451854ab7e4b4c146bf78a/preprints/The-Euclidean-plane-is-not-five-colorable-September-23-2026>.
- [11] OpenAI, *openai/math*, Lean development, <https://github.com/openai/math/tree/adc7f1241b42e322a6451854ab7e4b4c146bf78a/lean/OAI/Analysis/PlaneSpectrum>.
- [12] S. Li, <https://github.com/daizisheng/erdos-1070>, release v1.0.