

COPRIME PAIRS WITH EQUAL SUM OF DIVISORS, ASSUMING SMOOTH SHIFTED PRIMES

SHISHENG LI

ABSTRACT. Let $h(x)$ be the number of pairs $1 \leq a < b < x$ with $\gcd(a, b) = 1$ and $\sigma(a) = \sigma(b)$. Erdős asked whether $h(x) > x^{2-o(1)}$ (Erdős Problem #824). By an argument of Pollack and Pomerance, this follows if for every fixed $\delta > 0$ there are $y^{1-o(1)}$ primes $p \leq 5y$ with $p+1$ free of prime factors exceeding y^δ . We write this implication out with explicit parameters and formalize it in Lean 4. We then check that the argument of the recent OpenAI preprint *Weighted dilation graphs, smooth shifted primes and totient fibers*, which proves the corresponding statement for $p-1$, applies to $p+1$ after a change of sign in two of its sections. Consequently $h(x) = x^{2-o(1)}$, conditionally on the correctness of that preprint, which has not been refereed and whose main new estimates we have not independently verified.

1. INTRODUCTION

Let $\sigma(n)$ be the sum of the divisors of n and

$$h(x) = \#\{(a, b) : 1 \leq a < b < x, \gcd(a, b) = 1, \sigma(a) = \sigma(b)\}.$$

Trivially $h(x) < x^2$. Erdős [5] asked whether $h(x)/x \rightarrow \infty$; Problem #824 of [3] asks whether $h(x) > x^{2-o(1)}$. Pollack and Pomerance [7, §6] answered Erdős's question, proving $h(x) > x^{1.4}$ for large x . Following an idea of Erdős [4], their argument shows: if $\theta > 1$ is such that there are $y^\theta/(\log y)^{O(1)}$ primes $p \leq y^\theta$ with $P^+(p+1) \leq y$, then there are at least $x^{(2-\varepsilon)(1-1/\theta)+o(1)}$ coprime pairs, for every fixed $\varepsilon > 0$. Here $P^+(m)$ is the largest prime factor of m . With the Baker–Harman value $\theta \approx 3.377$ [1] this gives the exponent 1.4; an admissible θ for every $\theta > 1$ would give $x^{2-o(1)}$, as also remarked by Tao on the problem's forum [9]. A claim of $h(x) \geq x^{13/8-o(1)}$ by a different route was submitted to the forum in August 2026 [2].

For a fixed smoothness exponent, the best unconditional result is due to Lichtman [6]: for fixed $a \neq 0$ and $\delta > 15/(32\sqrt{e}) = 0.2843\dots$ there are $\gg x/(\log x)^C$ primes $x < p \leq 2x$ with $P^+(p-a) \leq x^\delta$. The corresponding statement for every fixed $\delta > 0$ is a conjecture of Erdős. For $a = 1$ it is claimed in the OpenAI preprint [10] (“WD” below):

Theorem 1.1 ([10, Theorem 1.2]). *For every fixed $0 < \delta < 1/4$, as $x \rightarrow \infty$, $\#\{p \text{ prime} : 2x < p \leq 5x, P^+(p-1) \leq x^\delta\} \geq x^{1-o(1)}$.*

The application to σ needs $p+1$ instead of $p-1$.

Definition 1.2. Hypothesis A^+ : for every $0 < \delta < 1/4$ and every $\eta > 0$ there is y_0 such that for every real $y \geq y_0$,

$$\#\{p \text{ prime} : p \leq 5y, P^+(p+1) \leq y^\delta\} \geq y^{1-\eta}.$$

Theorem 1.3. *If A^+ holds, then for every $\varepsilon > 0$ we have $h(x) > x^{2-\varepsilon}$ for all sufficiently large x . The pairs may moreover be taken with a, b squarefree.*

Theorem 1.3 is the argument of [7, §6] with $\theta \rightarrow \infty$. We give it in full in §3, with explicit parameters and under the weaker hypothesis of $y^{1-o(1)}$ (rather than $y/(\log y)^{O(1)}$) smooth shifted primes, because this is the version we have formalized in Lean 4 (§4).

Theorem 1.4. *The proof of Theorem 1.1 in [10], modified as described in §2, proves: for every fixed $0 < \delta < 1/4$, $\#\{p \text{ prime} : 2x - 2 < p \leq 5x, P^+(p+1) \leq x^\delta\} \geq x^{1-o(1)}$. In particular A^+ holds if the argument of [10] is correct.*

Corollary 1.5. *If the argument of [10] is correct, then $h(x) = x^{2-o(1)}$.*

The status of the corollary is exactly that of [10], an unrefereed preprint. Its Sections 3–5 introduce a new transference and shifted-correlation method; we have not verified them independently (§5). Our contribution is the observation that the sign change goes through, a check of the sections that it affects, and the formalization of Theorem 1.3.

2. SMOOTH SUCCESSORS FROM THE ARGUMENT OF WD

Theorem 1.4 is proved by the argument of [10] for Theorem 1.1, which detects primes of the form $2u + 1$ with u supported on x^δ -smooth integers; we apply it to primes of the form $2u - 1$ instead, so that $p + 1 = 2u$ is x^δ -smooth. The results of Sections 2–5 of [10] are used as stated. In Sections 6–7 of [10] (the Type II estimate and the prime extraction) one replaces $2u + 1$ by $2u - 1$ throughout and adjusts the congruences accordingly; the local densities and residue counts of the sieve are the same for both signs.

The one step that is not purely notational is the determinant parametrization in the Type II estimate. Two factorizations $mn = 2eh - 1$, $m'n = 2eh' - 1$ with $m \neq m'$ give $m' - m = -2ek$ and $h' - h = -kn$ for some integer $k \neq 0$, and then $mh' - m'h = -k(mn - 2eh) = k$. So the shifted endpoints are $z = m'h$ and $z + k = mh'$, as in [10], and the shift-cancellation theorem [10, Theorem 5.1], which holds for all $0 < |k| \leq (\log x)^C$, applies unchanged; the original argument already uses both signs of k .

3. PROOF OF THEOREM 1.3

It suffices to treat $0 < \varepsilon < 1$: for $\varepsilon \geq 1$ take any $\varepsilon' \in (0, 1)$; then $h(x) > x^{2-\varepsilon'} \geq x^{2-\varepsilon}$ for $x \geq 1$. Write $L = \log x$. Fix $\varepsilon \in (0, 1)$, a real $B \geq 8/\varepsilon$, $\eta = \varepsilon/16$ and $\delta = 1/(2B) \in (0, 1/4)$; these are fixed before $x \rightarrow \infty$. For large x put $y = L^B$,

$$\mathcal{P} = \{p \text{ prime} : p \leq 5y, P^+(p+1) \leq y^\delta\}, \quad M = |\mathcal{P}|.$$

By A^+ , $M \geq y^{1-\eta} = L^{B(1-\eta)}$ for all large x . Let $k = \lfloor L/\log(5y+1) \rfloor$. Then $(5y+1)^k \leq x$, and since $\log(5y+1) = B \log L + O(1)$,

$$(1) \quad k = \frac{L}{B \log L} (1 + o(1)), \quad k \leq L.$$

For large x , $M \geq L^2 \geq 2k$ because $B(1-\eta) > 2$.

3.1. The family. Let $\mathcal{A}$ be the set of products of k distinct primes of $\mathcal{P}$. Each $a \in \mathcal{A}$ is squarefree and $a < (5y+1)^k \leq x$, and distinct k -subsets give distinct products. So $N := |\mathcal{A}| = \binom{M}{k} \geq (M/k)^k$ and, by (1) and $\log k \leq \log L$,

$$(2) \quad \log N \geq k \log L (B(1-\eta) - 1) = L \left(1 - \eta - \frac{1}{B}\right) (1 + o(1)).$$

For $a = p_1 \cdots p_k$ we have $\sigma(a) = \prod_i (p_i + 1) \leq x$, and every prime factor of $\sigma(a)$ is at most $y^\delta = L^{1/2}$. The number U of $L^{1/2}$ -smooth integers in $[1, x]$ satisfies

$$(3) \quad U \leq \left(1 + \frac{L}{\log 2}\right)^{L^{1/2}}, \quad \log U = o(L),$$

since such an integer is $\prod_{q \leq L^{1/2}} q^{e_q}$ with $0 \leq e_q \leq L/\log 2$.

3.2. Many equal values. With $r(s) = \#\{a \in \mathcal{A} : \sigma(a) = s\}$, Cauchy's inequality gives

$$\#\{(a, b) \in \mathcal{A}^2 : \sigma(a) = \sigma(b)\} = \sum_s r(s)^2 \geq N^2/U.$$

3.3. Pairs sharing many primes. Let $t = \lceil \eta k \rceil$. The ordered pairs in $\mathcal{A}^2$ sharing at least t primes number at most $N \binom{k}{t} \binom{M}{k-t}$ (choose a , t of its primes, and the other $k-t$ primes of b), and

$$\binom{M}{k-t} / \binom{M}{k} = \prod_{i=0}^{t-1} \frac{k-i}{M-k+t-i} \leq \left(\frac{2k}{M}\right)^t, \quad \binom{k}{t} \leq 2^k.$$

By (1), (3) and $M \geq L^{B(1-\eta)}$,

$$\begin{aligned} \log(2^k (2k/M)^t \cdot 2U) &\leq (k+1) \log 2 + \log U - \eta k (B(1-\eta) \log L - \log(2L)) \\ &= -\eta L \left(1 - \eta - \frac{1}{B}\right) (1 + o(1)) + o(L), \end{aligned}$$

which is negative for large x . So at least $N^2/(2U)$ ordered pairs (a, b) have $\sigma(a) = \sigma(b)$ and share fewer than t primes; in particular $a \neq b$.

3.4. Removing the common factor. For such a pair let $g = \gcd(a, b)$, $u = a/g$, $v = b/g$. As a, b are squarefree, g, u, v are pairwise coprime, so $\sigma(g)\sigma(u) = \sigma(g)\sigma(v)$ and $\sigma(u) = \sigma(v)$, with $\gcd(u, v) = 1$, $u \neq v$, and $u, v < x$ squarefree. The integer g is a product of at most $t-1 < \eta k$ primes below $5y+1$, so $g \leq (5y+1)^{\eta k} \leq x^\eta$. Given the ordered pair (u, v) , the pair $(a, b) = (gu, gv)$ is determined by g ; each unordered pair $\{u, v\}$ therefore arises from at most $2x^\eta$ ordered pairs (a, b) . Hence

$$h(x) \geq \frac{N^2}{4Ux^\eta}, \quad \log h(x) \geq \left(2 - 3\eta - \frac{2}{B}\right)L - o(L) > (2 - \varepsilon)L$$

for large x , because $3\eta + 2/B \leq 3\varepsilon/16 + \varepsilon/4 < \varepsilon/2$. $\square$

Remark 3.1. Since $h(x)$ only depends on the integers below x , $h(x) = h(\lceil x \rceil)$. Hence the theorem for integer x , which is the form formalized in Lean, implies it for real x : for $0 < \varepsilon < 1$, $h(x) = h(\lceil x \rceil) > \lceil x \rceil^{2-\varepsilon} \geq x^{2-\varepsilon}$. Pollack and Pomerance count pairs with $b \leq x$; the difference is immaterial.

4. FORMALIZATION

The lower bound in Theorem 1.3 is formalized in Lean 4 (v4.33.1) with Mathlib (commit 0df444a) in the directory `lean/` of [8]. The definitions are

```
noncomputable def h (x : ℕ) : ℕ :=
  ((Finset.Ico 1 x ×ˢ Finset.Ico 1 x).filter
    (fun p => p.1 < p.2 ∧ Nat.Coprime p.1 p.2 ∧ sigma 1 p.1 = sigma 1 p.2)).card
def SmoothSuccessorHypWeak : Prop :=
  ∀ δ : ℝ, 0 < δ → δ < 1/4 → ∀ η : ℝ, 0 < η → ∃ y0 : ℝ, ∀ y : ℝ, y0 ≤ y →
    (y ^ (1 - η) : ℝ) ≤ (((Finset.Iic [5*y]_+).filter
      (fun p => p.Prime ∧ ∀ q ∈ (p+1).primeFactors, ((q:ℕ):ℝ) ≤ y ^ δ))).card : ℝ)
```

where `sigma 1` is Mathlib's $\sigma_1 = \sigma$; `SmoothSuccessorHypWeak` is Definition 1.2. The theorem `theorem erdos824_of_smoothSuccessorWeak (H : SmoothSuccessorHypWeak) :`

$$\forall \varepsilon : \mathbb{R}, 0 < \varepsilon \rightarrow \exists X : \mathbb{R}, \forall x : \mathbb{N}, X \leq x \rightarrow (x : \mathbb{R}) \wedge (2 - \varepsilon) < (h \ x : \mathbb{R})$$

formalizes the lower bound of Theorem 1.3 (not the squarefree addendum, which is proved only in §3). It has no `sorry`, and `#print axioms` reports only `propext`, `Classical.choice` and `Quot.sound`. The Lean proof follows §3 with $B = 16/\varepsilon$ and $\delta = \varepsilon/32$.

5. STATUS

The implication in Theorem 1.3 is unconditional and machine-checked. Theorem 1.4 rests on [10]. We rechecked Sections 6–7 of WD for the change of sign, but we have not independently verified Sections 3–5 (transference for weighted dilation graphs, construction of the ideal operator, lifted shift cancellation; about 2800 lines of source). Writing Lean statements for WD exposed several underspecified points. Examples: the hypothesis $\ell \geq 1$ is used in Section 4 but not stated in the definition of prime groups; Section 5 uses a coefficient form of the ideal-operator theorem [10, Theorem 4.1] that the theorem does not state. We have not assessed whether any of them affects the argument; this is part of the paper's explicit reliance on [10].

ACKNOWLEDGEMENTS

The reduction in §3 is due to Pollack and Pomerance [7], after Erdős; its relevance to Problem #824 was also pointed out by T. Tao. AI tools were used extensively in this work. Claude (Anthropic) checked the adaptation of [10] to $p + 1$, wrote the text and the Lean formalization. GPT models (OpenAI) were used to referee drafts. The preprint [10] is itself AI-generated. The author has checked the arguments presented here and takes responsibility for them.

REFERENCES

- [1] R. C. Baker and G. Harman, Shifted primes without large prime factors, *Acta Arith.* 83 (1998), 331–361.
- [2] Cam, A partial proof of Erdős Problem #824 with exponent $13/8$, submitted to [3] on 28 August 2026, <https://github.com/LeanMeanBean/Erdos-824>.
- [3] T. F. Bloom, Erdős problems, <https://www.erdosproblems.com>, Problem #824.
- [4] P. Erdős, On the normal number of prime factors of $p - 1$ and some related problems concerning Euler's φ -function, *Quart. J. Math. Oxford Ser. 6* (1935), 205–213.
- [5] P. Erdős, Remarks on number theory. II. Some problems on the σ function, *Acta Arith.* 5 (1959), 171–177.
- [6] J. D. Lichtman, Primes in arithmetic progressions to large moduli and shifted primes without large prime factors, arXiv:2211.09641.
- [7] P. Pollack and C. Pomerance, Some problems of Erdős on the sum-of-divisors function, *Trans. Amer. Math. Soc. Ser. B* 3 (2016), 1–26.
- [8] S. Li, Erdős #824: Lean formalization, <https://github.com/daizisheng/erdos-824>.
- [9] T. Tao, comment on Problem #824, Erdős problems forum, 11 August 2025, <https://www.erdosproblems.com/forum/thread/824>.
- [10] OpenAI, Weighted dilation graphs, smooth shifted primes and totient fibers, OpenAI Math Release preprint, 24 September 2026, <https://github.com/openai/math> (version of commit `adc7f12`).