

# Multivariable Calculus - Problem Sheet 2

Dom Hutchinson

October 17, 2018

## My Solution 4.1

$$\begin{aligned} f(\mathbf{r}) &= \mathbf{a} \circ \mathbf{r} \\ &= a_1x + a_2y + a_3z \\ \implies \nabla f(\mathbf{r}) &= (a_1, a_2, a_3) \end{aligned}$$

## My Solution 4.2

$$\begin{aligned} \mathbf{v}(\mathbf{r}) &= \nabla r^n \\ \text{Since } r &= \sqrt{x^2 + y^2 + z^2} \\ \implies \nabla \circ \mathbf{v}(\mathbf{r}) &= \nabla \circ \nabla r^n \\ &= \nabla \circ \nabla (x^2 + y^2 + z^2)^{n/2} \\ &= \nabla \circ \left( xn(x^2 + y^2 + z^2)^{\frac{n}{2}-1}, yn(x^2 + y^2 + z^2)^{\frac{n}{2}-1}, zn(x^2 + y^2 + z^2)^{\frac{n}{2}-1} \right) \\ &= \left( n(x^2 + y^2 + z^2)^{\frac{n}{2}-1} + xn(xn - 2x)(x^2 + y^2 + z^2)^{\frac{n}{2}-2} \right) \\ &\quad + \left( n(x^2 + y^2 + z^2)^{\frac{n}{2}-1} + yn(yn - 2y)(x^2 + y^2 + z^2)^{\frac{n}{2}-2} \right) \\ &\quad + \left( n(x^2 + y^2 + z^2)^{\frac{n}{2}-1} + zn(zn - 2z)(x^2 + y^2 + z^2)^{\frac{n}{2}-2} \right) \\ &= 3n(x^2 + y^2 + z^2)^{\frac{n}{2}-1} + n(x^n - 2x^2 + y^2n - 2y^2 + z^2n - 2z^2)(x^2 + y^2 + z^2)^{\frac{n}{2}-2} \\ &= n(x^2 + y^2 + z^2)^{\frac{n}{2}-2} (3(x^2 + y^2 + z^2) + (x^n - 2x^2 + y^2n - 2y^2 + z^2n - 2z^2)) \\ &= n(x^2 + y^2 + z^2)^{\frac{n}{2}-2} ((n+1)(x^2 + y^2 + z^2)) \end{aligned}$$

This equals zero if  $n = 0$  or  $n = -1$ .

## My Solution 4.3

$$\begin{aligned} \mathbf{v}(\mathbf{r}) &= \boldsymbol{\omega} \times \mathbf{r} \\ &= \begin{vmatrix} \mathbf{e}_1 & \mathbf{e}_2 & \mathbf{e}_3 \\ \omega_1 & \omega_2 & \omega_3 \\ x & y & z \end{vmatrix} \\ &= \left( \begin{vmatrix} \omega_2 & \omega_3 \\ y & z \end{vmatrix}, \begin{vmatrix} \omega_1 & \omega_3 \\ x & z \end{vmatrix}, \begin{vmatrix} \omega_1 & \omega_2 \\ x & y \end{vmatrix} \right) \\ &= (\omega_2z - \omega_3y, \omega_1z - \omega_3x, \omega_1y - \omega_2x) \\ \text{Then } \nabla \times \mathbf{v} &= \begin{vmatrix} \mathbf{e}_1 & \mathbf{e}_2 & \mathbf{e}_3 \\ \partial_x & \partial_y & \partial_z \\ \omega_2z - \omega_3y & \omega_1z - \omega_3x & \omega_1y - \omega_2x \end{vmatrix} \\ &= \left( \frac{\partial}{\partial y}(\omega_1y - \omega_2x) - \frac{\partial}{\partial z}(\omega_1z - \omega_3x), \right. \\ &\quad \left. \frac{\partial}{\partial x}(\omega_1y - \omega_2x) - \frac{\partial}{\partial z}(\omega_2z - \omega_3y), \right. \\ &\quad \left. \frac{\partial}{\partial x}(\omega_1z - \omega_3x) - \frac{\partial}{\partial y}(\omega_2z - \omega_3y) \right) \\ &= (\omega_1 - \omega_1, -\omega_2 - \omega_2, -\omega_3 - (-\omega_3)) \\ &= (0, -2\omega_2, 0) \end{aligned}$$

## My Solution 7

The cross product is anticommutative, but this identity shows it to be commutative.

The correct identity is

$$\nabla \circ (\mathbf{u} \times \mathbf{v}) = \mathbf{v} \circ (\nabla \times \mathbf{u}) - \mathbf{u} \circ (\nabla \times \mathbf{v})$$

### My Solution 8

$$\begin{aligned}
 [\nabla \times (\mathbf{u} \times \mathbf{v})]_i &= \varepsilon_{ijk} \frac{\partial}{\partial x_j} [\mathbf{u} \times \mathbf{v}]_k \\
 &= \varepsilon_{ijk} \frac{\partial}{\partial x_j} \varepsilon_{klm} (u_l v_m) \\
 &= \varepsilon_{ijk} \varepsilon_{klm} \frac{\partial}{\partial x_j} (u_l v_m) \\
 &= \varepsilon_{kij} \varepsilon_{klm} \left( \frac{\partial u_l}{\partial x_j} v_m + u_l \frac{\partial v_m}{\partial x_j} \right) \\
 &= (\delta_{il} \delta_{jm} - \delta_{im} \delta_{jl}) \left( \frac{\partial u_l}{\partial x_j} v_m + u_l \frac{\partial v_m}{\partial x_j} \right) \\
 &= \left( \frac{\partial u_i}{\partial x_j} v_j + u_i \frac{\partial v_j}{\partial x_j} \right) - \left( \frac{\partial u_j}{\partial x_j} v_i + u_j \frac{\partial v_i}{\partial x_j} \right) \\
 &= ([(\mathbf{v} \circ \nabla) \mathbf{u}]_i + [(\nabla \circ \mathbf{v}) \mathbf{u}]_i) - ([(\nabla \circ \mathbf{u}) \mathbf{v}]_i + [(\mathbf{u} \circ \nabla) \mathbf{v}]_i)
 \end{aligned}$$

This is true for  $i = 1, 2, 3$  so identity holds.