

Statistics 2 - Notes

Dom Hutchinson

October 30, 2019

Contents

1	Estimation	2
1.1	Introduction	2
2	The Likelihood Function	3
3	Maximum Likelihood Estimates	4
3.1	Determining MLEs - The Tractable Case	5
4	Statistics and Estimators	7
5	Probabilistic Convergence	9
5.1	Probabilistic Convergence & Estimators	11
6	The Fisher Information	12
7	Efficiency and The Cramer-Rao Bound	15
8	Asymptotic Distribution of the Maximum Likelihood Estimator	17
9	Confidence Sets Around the Maximum Likelihood Estimator	20
10	Asymptotic Approximation of Confidence Intervals	21
11	Estimating the Information for Maximum Likelihood Estimates	22
0	Appendix	26
0.1	Notation	26
0.2	R	26
0.3	Probability Distributions	26

1 Estimation

1.1 Introduction

Definition 1.1 - *Probabilty Space, $(\Omega, \mathcal{F}, \mathbb{P})$*

A mathematical construct for modelling the real world. A *Probabilty Space* has three elements

- i) Ω - Sample space.
- ii) \mathcal{F} - Set of events.
- iii) \mathbb{P} - Probability measure.

and must fulfil the following conditions

- i) $\Omega \in \mathcal{F}$;
- ii) $\forall A \in \mathcal{F} \implies A^c \in \mathcal{F}$;
- iii) $\forall A_0, \dots, A_n \in \mathcal{F} \implies \left(\bigcup_i A_i \right) \in \mathcal{F}$;
- iv) $\mathbb{P}(\Omega) = 1$; and,
- v) $\mathbb{P}\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} \mathbb{P}(A_i)$ for disjoint A_1, A_2, \dots (Countable Additivity).

Definition 1.2 - *Random Variable*

A function which maps an event in the sample space to a value *e.g.* $X : \Omega \rightarrow \mathbb{R}$.

Remark 1.1 - *Probability Density Function for iid Random Variable Vector*

For $\mathbf{X} \sim f_n(\cdot; \theta)$ where each component of \mathbf{X} is independent and identically distribution the probability density function of \mathbf{X} is

$$f_n(\mathbf{x}; \theta) = \prod_{i=1}^n f(x_i; \theta)$$

Definition 1.3 - *Expectation*

The mean value for a random variable. For rv X

$$\mathbb{E}(X) := \sum_{x \in \mathcal{X}} x f_X(x) \quad \& \quad \mathbb{E}(X) := \int_{\mathbb{R}} x f_X(x) dx$$

Theorem 1.1 - *Expection of a Function*

For a function $g : \mathbb{R} \rightarrow \mathbb{R}$ and rv X with pmf f_X

$$\mathbb{E}(g(X)) := \sum_{g(x) \in \mathcal{X}} x f_X(x) \quad \& \quad \mathbb{E}(g(X)) := \int_{\mathbb{R}} g(x) f_X(x) dx$$

Theorem 1.2 - *Expectation of a Linear Operator*

For rv X with pmf f_X & $a, b \in \mathbb{R}$

$$\mathbb{E}(aX + b) = a\mathbb{E}(X) + b$$

Definition 1.4 - *Variance*

For rv X

$$\text{Var}(X) := \mathbb{E}[(X - \mathbb{E}(X))^2] = \mathbb{E}(X^2) - \mathbb{E}(X)^2$$

Theorem 1.3 - Variance of a Linear Operator

For rv X and $a, b \in \mathbb{R}$

$$\text{Var}(aX + b) = a^2 \text{Var}(X)$$

Definition 1.5 - Moment of a Random Variable

For rv X the n^{th} moment of X is defined as $\mathbb{E}(X^n)$.

N.B. - $\mathbb{E}(X^n) \neq \mathbb{E}(X)^n$.

Definition 1.6 - Covariance

For rv X & Y

$$\text{Cov}(X, Y) := \mathbb{E}[(X - \mathbb{E}(X))(Y - \mathbb{E}(Y))] = \mathbb{E}(XY) - \mathbb{E}(X)\mathbb{E}(Y)$$

Theorem 1.4 - Properties of Covariance

Let X & Y be independent random variables

i) $\text{Cov}(X, X) = \text{Var}(X)$;

ii) $\text{Cov}(X, Y) = 0$

Theorem 1.5 - Variance of two Random Variables with linear operators

$$\text{Var}(aX + bY) = a^2 \text{Var}(X) + b^2 \text{Var}(Y) + 2ab \text{Cov}(X, Y)$$

Theorem 1.6 - Independent Random Variables

Random variables X_1, \dots, X_n are independent iff

$$\mathbb{P}(X_1 \leq a_1, \dots, X_n \leq a_n) = \prod_{i=1}^n \mathbb{P}(X_i \leq a_i) \quad \forall a_1, \dots, a_n \in \mathbb{R}$$

2 The Likelihood Function

Definition 2.1 - Likelihood Function

Define $\mathbf{X} \sim f_n(\cdot; \theta^*)$ for some unknown $\theta^* \in \Theta$ and let \mathbf{x} be an observation of \mathbf{X} .

A *Likelihood Function* is any function, $L(\cdot; \mathbf{x}) : \Theta \rightarrow [0, \infty)$, which is proportional to the PMF/PDF of the observed realisation \mathbf{x} .

$$L(\theta; \mathbf{x}) := C f_b(\mathbf{x}; \theta) \quad \forall C > 0$$

N.B. Sometimes this is called the *Observed Likelihood Function* since it is dependent on observed data.

Definition 2.2 - Log-Likelihood Function

Let $\mathbf{X} \sim f_n(\cdot; \theta^*)$ for some unknown $\theta^* \in \Theta$ and \mathbf{x} be an observation of \mathbf{X} .

The *Log-Likelihood Function* is the natural log of a *Likelihood Function*

$$\ell(\theta; \mathbf{x}) := \ln f_n(\mathbf{x}; \theta) + C, \quad C \in \mathbb{R}$$

Theorem 2.1 - Multidimensional Transforms

Let \mathbf{X} be a continuous random vector in \mathbb{R}^n with PDF $f_{\mathbf{X}}$; $g : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a continuous differentiable bijection; and, $h := g^{-1}$.

Then $\mathbf{Y} = g(\mathbf{X})$ is a continuous random vector and its PDF is

$$f_{\mathbf{Y}}(\mathbf{y}) = f_{\mathbf{X}}(h(\mathbf{y})) H_h(\mathbf{Y})$$

where

$$J_h := \left| \det \left(\frac{\partial h}{\partial \mathbf{y}} \right) \right|$$

Proposition 2.1 - *Invariance of Likelihood Function by bijective transformation of the observations independent of θ*

Let $g : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a bijective transformation which is independent of θ ; and $\mathbf{Y} := g(\mathbf{X})$.

Then \mathbf{Y} is a random variable with PDF/PMF

$$f_{\mathbf{Y}}(\mathbf{y}; \theta) \propto f_{\mathbf{X}}(g^{-1}(\mathbf{y}); \theta)$$

Hence, if $\mathbf{y} = g(\mathbf{x})$ then $L_{\mathbf{Y}}(\theta; \mathbf{y}) \propto L_{\mathbf{X}}(\theta; \mathbf{x})$

Proof 2.1 - *Proposition 2.1*

Let $g : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a bijective transformation which is independent of θ ; $h := g^{-1}$; \mathbf{X}, \mathbf{Y} be a rvs st $\mathbf{Y} := g(\mathbf{X})$.

i) *Discrete Case* - Consider the case when \mathbf{X} is a discrete rv. Then

$$\begin{aligned} f_{\mathbf{Y}}(\mathbf{y}; \theta) &= \mathbb{P}(\mathbf{Y} = \mathbf{y}; \theta) \\ &= \mathbb{P}(g^{-1}(\mathbf{Y}) = g^{-1}(\mathbf{y}); \theta) \\ &= \mathbb{P}(h(\mathbf{Y}) = h(\mathbf{y}); \theta) \\ &= \mathbb{P}(\mathbf{X} = h(\mathbf{y}); \theta) \\ &= f_{\mathbf{X}}(g^{-1}(\mathbf{y}); \theta) \end{aligned}$$

ii) *Continuous Case* - Consider the case when \mathbf{X} is a continuous rv.

Then, by **Theorem 2.1**

$$f_{\mathbf{Y}}(\mathbf{y}; \theta) = f_{\mathbf{X}}(g^{-1}(\mathbf{y}); \theta) J_{g^{-1}}(\mathbf{y})$$

Since $J_{g^{-1}}$ does not depend on θ this case is solved.

Thus in both cases $L_{\mathbf{Y}}(\theta; \mathbf{y}) = f_{\mathbf{Y}}(\mathbf{y}; \theta) \propto f_{\mathbf{X}}(g^{-1}(\mathbf{y}); \theta) = L_{\mathbf{X}}(\theta; \mathbf{x})$. □

3 Maximum Likelihood Estimates

Definition 3.1 - *Maximum Likelihood Estimate*

Let $\mathbf{X} \sim f_n(\cdot; \theta)$; and \mathbf{x} be a realisation of \mathbf{X} .

The *Maximum Likelihood Estimate* is the value $\hat{\theta} \in \Theta$ st

$$\forall \theta \in \Theta \quad f_n(\mathbf{x}; \hat{\theta}) \geq f_n(\mathbf{x}, \theta)$$

Equivalently

$$\forall \theta \in \Theta \quad L(\hat{\theta}; \mathbf{x}) \geq L(\theta; \mathbf{x}) \quad \text{or} \quad \ell(\hat{\theta}; \mathbf{x}) \geq \ell(\theta; \mathbf{x})$$

i.e. $\hat{\theta}(\mathbf{x}) := \operatorname{argmax}_{\theta} (L(\theta; \mathbf{x}))$.

Remark 3.1 - *The Maximum Likelihood Estimate may not be unique*

Example 3.1 - *MLE for Uniform Distribution*

Consider $\mathbf{X} \stackrel{\text{iid}}{\sim} U[0, \theta]$ for $\theta > 0$.

Then

$$\begin{aligned} L(\theta; \mathbf{x}) &\propto f_n(\mathbf{x}; \theta) \\ &= \prod_{i=1}^n \frac{1}{\theta} \mathbb{1}\{x_i \in [0, \theta]\} \\ &= \frac{1}{\theta^n} \prod_{i=1}^n \mathbb{1}\{x_i \in [0, \theta]\} \\ \implies \hat{\theta} &= \max\{x_i : x_i \in \mathbf{x}\} \end{aligned}$$

Remark 3.2 - MLE of Reparameterisation

Define $\tau(\theta) : \mathbb{R} \rightarrow \mathbb{R}$. Then

$$\hat{\tau} = \tau(\hat{\theta})$$

N.B. We often write \tilde{f} to represent the pmf when τ is taken as a parameter rather than θ . *i.e.* $f(x; \theta) = \tilde{f}(x; \tau(\theta))$.

Theorem 3.1 - Invariance of MLE under bijective Reparameterisation

Let $g : \Theta \rightarrow G$ be a bijective transformation of the statistical parameter θ .

Let $\mathbf{X} \sim f(\cdot; \theta) = \tilde{f}(\cdot; g(\theta))$ for some θ , and let \mathbf{x} be a realisation of \mathbf{X} .

If $\hat{\theta}$ is an MLE of θ then $\hat{\tau} = g(\hat{\theta})$ is an MLE of τ .

Proof 3.1 - Theorem 3.1

This is a proof by contradiction.

Suppose $\exists \tau^* \in G$ st $\tilde{f}(x; \tau^*) > \tilde{f}(x; \hat{\tau})$. We know that $\forall \theta \in \Theta$, $f(x; \theta) = \tilde{f}(x; g(\theta))$ and $\forall \tau \in G$, $f(x; g^{-1}(\tau)) = \tilde{f}(x; \tau)$.

We deduce that

$$\begin{aligned} f(x; g^{-1}(\tau^*)) &= \tilde{f}(x; \tau^*) \\ &> \tilde{f}(x; \hat{\tau}) \text{ by assumption} \\ &= f(x; g^{-1}(\hat{\tau})) \\ &= f(x; \hat{\theta}) \end{aligned}$$

This contradicts the assumption that $\hat{\theta}$ is an maximum likelihood estimate of θ .

□

Remark 3.3 - Not all Reparameterisations are Bijective

When reparameterisations $g : \mathbb{R} \rightarrow \mathbb{R}$ is not bijective it is helpful to consider the *induced likelihood*

$$L^*(\tau; \mathbf{x}) := \max_{\theta \in G_\tau} L(\theta; \mathbf{x}) \text{ where } G_\tau := \{\theta : g(\theta) = \tau\}$$

Since this reduces the domain to only where g is bijective.

3.1 Determining MLEs - The Tractable Case**Proposition 3.1 - Differentiable Likelihood in the continuous case - Multivariate**

When $L(\theta; \mathbf{x})$ is differentiable one can find MLEs by considering its extrema. This is done equating & solving the cases when the gradient is zero, *i.e.* $\nabla L(\theta; \mathbf{x}) = 0$, and then checking whether this is a maximum or minimum point.

A point is a local minimum if the Hessian at the point is *Negative Definite* *i.e.* $\mathbf{x}^T \mathbf{A} \mathbf{x} < 0 \forall \mathbf{x} \neq \mathbf{0}$.

Example 3.2 - MLE of Normal Distribution

Let $\mathbf{X} \stackrel{\text{iid}}{\sim} \mathcal{N}(\mu, \sigma^2)$

$$\begin{aligned}
 L(\mu, \sigma^2; \mathbf{x}) &= \prod_{i=1}^n \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x_i - \mu)^2}{2\sigma^2}} \\
 \Rightarrow \ell(\mu, \sigma^2; \mathbf{x}) &= C - \frac{n}{2} \ln(\sigma^2) - \frac{1}{2\sigma^2} \sum_{i=1}^n (x_i - \mu)^2 \\
 \Rightarrow \nabla \ell(\mu, \sigma^2; \mathbf{x}) &= \left(\frac{-1}{\sigma^2} \sum_{i=1}^n (x_i - \mu), \quad -\frac{n}{2\sigma^2} + \frac{1}{2\sigma^4} \sum_{i=1}^n (x_i - \mu)^2 \right) \\
 \text{Setting } \frac{-1}{\sigma^2} \sum_{i=1}^n (x_i - \mu) &= 0 \\
 \Rightarrow \hat{\mu} &= \frac{1}{n} \sum_{i=1}^n x_i = \bar{x} \\
 \text{Setting } -\frac{n}{2\sigma^2} + \frac{1}{2\sigma^4} \sum_{i=1}^n (x_i - \mu)^2 &= 0 \\
 \Rightarrow \hat{\sigma}^2 &= \frac{1}{n} \sum_{i=1}^n (x_i \hat{\mu})^2
 \end{aligned}$$

We now want to check whether $(\hat{\mu}, \hat{\sigma}^2)$ is a minimum.

$$\begin{aligned}
 \nabla^2 \ell(\mu, \sigma^2; \mathbf{x}) &= \begin{pmatrix} \frac{\partial^2 \ell(\mu, \sigma^2; \mathbf{x})}{\partial \mu^2} & \frac{\partial^2 \ell(\mu, \sigma^2; \mathbf{x})}{\partial \mu \partial \sigma^2} \\ \frac{\partial^2 \ell(\mu, \sigma^2; \mathbf{x})}{\partial \mu \partial \sigma^2} & \frac{\partial^2 \ell(\mu, \sigma^2; \mathbf{x})}{\partial (\sigma^2)^2} \end{pmatrix} \\
 &= \begin{pmatrix} -\frac{n}{\hat{\sigma}^2} & 0 \\ 0 & -\frac{n}{2\hat{\sigma}^4} \end{pmatrix}
 \end{aligned}$$

Since $\begin{pmatrix} z_1 & z_2 \end{pmatrix} \begin{pmatrix} -a & 0 \\ 0 & -b \end{pmatrix} \begin{pmatrix} z_1 \\ z_2 \end{pmatrix} = -az_1^2 - bz_2^2 < 0 \forall a, b > 0$ and we have $\frac{n}{\hat{\sigma}^2}, \frac{n}{2\hat{\sigma}^4} > 0$ then we can conclude that $\nabla^2 \ell$ is negative definite.

Thus $\hat{\mu} = \bar{x}$ & $\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (x_i \hat{\mu})^2$ is an MLE for the normal distribution.

Example 3.3 - MLE for Capture-Recapture Model

Suppose you are wanting to calculate the unknown size of a population, n . The Capture-Recapture Model is one technique that can be used. You tag $t \leq n$ members of the population; wait for a while; then recapture $c \leq n$ members of which $x \leq \min\{t, c\} \leq n$ are tagged.

With t, c, x known produce a MLE for n .

We first work out the associated probability distribution for X , the population size. We have

- i) $\binom{t}{x}$ ways of choosing x members among the tagged ones;
- ii) $\binom{n-t}{c-x}$ ways of choosing the remaining members among the non-tagged ones;
- iii) $\binom{n}{c}$ ways of choosing c members in a population of n individuals.

Thus

$$f_X(x; n) = \frac{\binom{t}{x} \binom{n-t}{c-x}}{\binom{n}{c}}$$

This means that $X \sim \text{Hypergeometric}(t, n, c)$ with t & c known.

Now we calculate the MLE for X

$$\begin{aligned}
 L(n; x) &= f_X(x; n) \\
 &= \frac{\binom{t}{x} \binom{n-t}{c-x}}{\binom{n}{c}} \\
 &= \frac{x!(t-x)! (c-x)! (n-t-c+x)!}{t! (n-t)! c! (n-c)!}
 \end{aligned}$$

Now we consider $L(n; x) = 0$ when $x > \min\{t, c\}$. We want to identify values of n for which $L(n; x) \geq L(n-1; x)$.

Consider $n-1 \geq \min\{t, c\} \implies L(n-1; x) > 0$

$$\begin{aligned}
 \text{Let } r(n) &:= \frac{L(n; x)}{L(n-1; x)} \\
 &= \frac{n-t}{n-t-c+x} \frac{n-c}{n} \\
 \implies 1 &\leq r(n) \\
 \Leftrightarrow 1 &\leq \frac{n-t}{n-t-c+x} \frac{n-c}{n} \\
 \Leftrightarrow n(n-t-c+x) &\leq (n-t)(n-c) \\
 \Leftrightarrow n^2 - nt - cn + xn &\leq n^2 - nt - cn + ct \\
 \Leftrightarrow xn &\leq ct \\
 \Leftrightarrow x &\leq \frac{ct}{n}
 \end{aligned}$$

So $L(n; x)$ is increasing for $n \leq \lfloor \frac{ct}{x} \rfloor$ & decreasing for $n > \lfloor \frac{ct}{x} \rfloor$.
Consequently $\hat{n}_{\text{MLE}}(x) = \lfloor \frac{ct}{x} \rfloor$

4 Statistics and Estimators

Definition 4.1 - Statistic

Given some data \mathbf{x} a statistic is a function of the data $T(\mathbf{x})$.

N.B. A statistic cannot depend on an unknown statistical parameter.

Definition 4.2 - Estimate

Let $\mathbf{X} \sim f_n(\cdot; \theta^*)$ with $\theta^* \in \Theta$ and \mathbf{x} be a realisation of \mathbf{X} .

An *Estimate* θ^* is a statistic $\hat{\theta}(\mathbf{x}) = T(\mathbf{x})$ which is intended to approximate the real value of θ^* .

N.B. An *Estimate* is a real value & thus is hard to evaluate.

Definition 4.3 - Estimator

Let $\mathbf{X} \sim f_n(\cdot; \theta^*)$ with $\theta^* \in \Theta$ and \mathbf{x} be a realisation of \mathbf{X} .

An *Estimator* of θ^* is $\hat{\theta}$ where $\hat{\theta}(\mathbf{x})$ is an *estimate*.

N.B. We call $T(\mathbf{X})$ an estimator. This is a random variable.

Definition 4.4 - Distribution of an Estimator

Let $\mathbf{X} \sim f_n(\cdot; \theta^*)$ with $\theta^* \in \Theta \subseteq \mathbb{R}$.

If $\hat{\theta}(\mathbf{X})$ is a real-valued random variable, we can write its CDF as

$$\begin{aligned}
 F_{\hat{\theta}(\mathbf{X})}(t; \theta^*) &= \mathbb{P}(\hat{\theta}(\mathbf{X}) \leq t; \theta^*) \\
 &= \int_{\chi^n} \mathbb{1}\{\hat{\theta}(\mathbf{x}) \leq t\} f_n(\mathbf{x}; \theta^*) d\mathbf{x}
 \end{aligned}$$

Remark 4.1 - Estimator depends upon true value

The distribution of $\hat{\theta}(\mathbf{X})$ depends on the distribution of \mathbf{X} which in turn depends upon the distribution of θ^* .

Thus the distribution of an estimator depends on the true parameter of the variable it is estimating.

Remark 4.2 - Estimator Distribution & Sample Size

As sample size increases the distribution of an estimator may converge to a more standard distribution (e.g. Normal, Poisson).

Definition 4.5 - Bias

Bias is a measure of how much an estimator deviates from the true value, on average.

$$\begin{aligned}\text{Bias}(\hat{\theta}; \theta^*) &:= \mathbb{E}(\hat{\theta}(\mathbf{X}) - \theta^*; \theta^*) \\ &= \mathbb{E}(\hat{\theta}; \theta^*) - \mathbb{E}(\theta^*; \theta^*) \\ &= \mathbb{E}(\hat{\theta}; \theta^*) - \theta^*\end{aligned}$$

Definition 4.6 - Unbiased Estimator

An *Estimator*, $\hat{\theta}$, is said to be *Unbiased* if $\forall \theta \in \Theta$, $\text{Bias}(\hat{\theta}; \theta) = 0$. Equivalently $\mathbb{E}(\hat{\theta}; \theta) = \theta$.

Definition 4.7 - Mean Square Error

The *Mean Square Error* of an estimator is the mean of the squared error associated with rv $\hat{\theta}$.

$$MSE(\hat{\theta}; \theta^*) := \mathbb{E}[(\hat{\theta}(\mathbf{X}) - \theta^*)^2; \theta^2]$$

Proposition 4.1 - Simplification of MSE Formula

The MSE is a combination of variance & bias.

$$\begin{aligned}MSE(\hat{\theta}; \theta^*) &= \mathbb{E}[(\hat{\theta}(\mathbf{X}) - \theta^*)^2; \theta^2] \\ &= \mathbb{E}\left[\left\{\hat{\theta} - \mathbb{E}(\hat{\theta}; \theta^*)\right\}^2; \theta^*\right] + \left(\mathbb{E}(\hat{\theta} - \theta^*; \theta^*)\right)^2 \\ &= \text{Var}(\hat{\theta}; \theta^*) + \text{Bias}(\hat{\theta}; \theta^*)^2\end{aligned}$$

Example 4.1 - Sample mean as an Estimator

Let $\mathbf{X} \stackrel{\text{iid}}{\sim} \text{Poisson}(\lambda^*)$.

Suppose we are using the sample mean, $\hat{\lambda}(\mathbf{x}) := \frac{1}{n} \sum_{i=1}^n x_i$, as an estimate of λ^* . We first want to show this estimator is *Unbiased*

$$\begin{aligned}\mathbb{E}(\hat{\lambda}; \lambda) &= \mathbb{E}\left(\frac{1}{n} \sum_{i=1}^n X_i; \lambda\right) \\ &= d_n^1 \sum_{i=1}^n \mathbb{E}(X_i; \lambda) \\ &= \frac{1}{n} n \lambda \\ &= \lambda\end{aligned}$$

Thus $\hat{\lambda}$ is unbiased.

Now we consider the MSE of $\hat{\lambda}$

$$\begin{aligned}MSE(\hat{\lambda}; \lambda) &= \text{Var}(\hat{\lambda}; \lambda) \\ &= \text{Var}\left(\frac{1}{n} \sum_{i=1}^n X_i; \lambda\right) \\ &= \frac{1}{n^2} \sum_{i=1}^n \text{Var}(X_i; \lambda) \\ &= \frac{1}{n^2} n \lambda \\ &= \frac{\lambda}{n}\end{aligned}$$

This shows that as the sample size increases the MSE of $\hat{\lambda}$ converges to 0.

5 Probabilistic Convergence

Remark 5.1 - Motivation

Here we consider the properties of a maximum likelihood estimators as the sample size increases.

Theorem 5.1 - Markov's Inequality

For a *non-negative* random variable X and a constant $a > 0$

$$\mathbb{P}(X \geq a) \leq \frac{\mathbb{E}(X)}{a}$$

Proof 5.1 - Markov's Inequality

Consider continuous X . We have

$$\begin{aligned} a\mathbb{P}(X \geq a) &= a \int_a^\infty f_X(x) dx \\ &\leq \int_a^\infty x f_X(x) dx \\ &\leq \int_0^\infty x f_X(x) dx \\ &= \mathbb{E}(X) \\ \implies a\mathbb{P}(X \geq a) &= \mathbb{E}(X) \\ \implies \mathbb{P}(X \geq a) &\leq \frac{\mathbb{E}(X)}{a} \end{aligned}$$

□

Theorem 5.2 - Chebyshev's Inequality

Let $\mu = \mathbb{E}(X)$ and $\sigma^2 = \text{Var}(X)$. Then

$$\forall a > 0, \mathbb{P}(|X - \mu| \geq a) \leq \frac{\sigma^2}{a^2}$$

Proof 5.2 - Chebyshev's Inequality

We have

$$\begin{aligned} \mathbb{P}(|X - \mu| \geq a) &= \mathbb{P}(|X - \mu|^2 \geq a^2) \\ &\leq \frac{\mathbb{E}((X - \mu)^2)}{a^2} \text{ By Markov's Inequality} \\ &= \frac{\sigma^2}{a^2} \end{aligned}$$

□

Definition 5.1 - Convergence in Probability

We say the sequence of random variables $\{Z_n\}_{n \in \mathbb{N}}$ converges in probability to the random variable Z if

$$\forall \varepsilon > 0, \lim_{n \rightarrow \infty} \mathbb{P}(|Z_n - Z| > \varepsilon) = 0$$

N.B. This is denoted $Z_n \rightarrow_{\mathbb{P}} Z$.

N.B. The random variables $\{Z_n\}_{n \in \mathbb{N}}$ & Z must be in the same probability space.

Theorem 5.3 - Weak Law of Large Numbers

If $\{X_n\}_{n \in \mathbb{N}}$ are independent & identically distributed and $\mathbb{E}(X_1) = \mu < \infty$ then

$$Z_n = \frac{1}{n} \sum_{i=1}^n X_i \rightarrow_{\mathbb{P}} \mu$$

N.B. This is an example of Convergence in Probability.

Definition 5.2 - Convergence in Distribution

We say the sequence of random variables $\{Z_n\}_{n \in \mathbb{N}}$ converges in distribution to random variable Z if

$$\forall z \in \mathbb{R} \text{ where } \mathbb{P}(Z \leq z) \text{ is continuous, } \lim_{n \rightarrow \infty} \mathbb{P}(Z_n \leq z) = \mathbb{P}(Z \leq z)$$

N.B. This is denoted $Z_n \rightarrow_{\mathcal{D}} Z$.

N.B. The random variables $\{Z_n\}_{n \in \mathbb{N}}$ & Z need not be in the same probability space.

Remark 5.2 - Equivalent Statements to Convergence in Distribution

Saying that $Z_n \rightarrow_{\mathcal{D}} Z$ is equivalent to saying that

$$\forall z \in \mathbb{R} \text{ where } F_Z(z) \text{ is continuous, } \lim_{n \rightarrow \infty} F_{Z_n}(z) = F_Z(z)$$

Theorem 5.4 - Central Limit Theorem

If $\{X_n\}_{n \in \mathbb{N}}$ are independent & identically distributed, $\mathbb{E}(X_1) = \mu < \infty$ and $\text{Var}(X_1) = \sigma^2 < \infty$ then

$$\frac{\sqrt{n}}{\sigma}(Z_n - \mu) \rightarrow_{\mathcal{D}} Z \sim \text{Normal}(0, 1)$$

Theorem 5.5 - Convergence in Probability & Distribution

Convergence in probability \implies Convergence in distribution, **but** the opposite is not necessarily true.

Theorem 5.6 - Convergence in Probability & Distribution to a Constant

Convergence in distribution to a constant **and** convergence in probability to a constant are equivalent.

Example 5.1 -

Let $X \sim \text{Bernoulli}(\frac{1}{2})$ and $\{X_n\}_{n \in \mathbb{N}}$ be a sequence of random variables where $X_i := (1 - X) + \frac{1}{n}$. We have

$$F_X(x) = \begin{cases} 0 & , x < 0 \\ \frac{1}{2} & , x \in [0, 1) \\ 1 & , x \geq 1 \end{cases} \quad F_{X_n}(x) = \begin{cases} 0 & , x < \frac{1}{n} \\ \frac{1}{2} & , x \in [\frac{1}{n}, 1 + \frac{1}{n}) \\ 1 & , x \geq 1 + \frac{1}{n} \end{cases}$$

Clearly $F_{X_n}(x) \rightarrow F_X(x)$ at all points at which F_X is continuous (i.e. $x \in \mathbb{R} \setminus \{0, 1\}$).

Thus $X_n \rightarrow_{\mathcal{D}} X$.

Theorem 5.7 - Continuous Mapping Theorem

Let $g : Z \rightarrow G$ be a *continuous* function. Then

- i) If $Z_n \rightarrow_{\mathbb{P}} Z$, then $g(Z_n) \rightarrow_{\mathbb{P}} g(Z)$;
- ii) If $Z_n \rightarrow_{\mathcal{D}} Z$, then $g(Z_n) \rightarrow_{\mathcal{D}} g(Z)$

Theorem 5.8 - Slutsky's Theorem

Let $\{Y_n\}_{n \in \mathbb{N}}$ & $\{Z_n\}_{n \in \mathbb{N}}$ be sequences of random variables, Y be a random variable & $c \in \mathbb{R} \setminus \{0\}$ be a constant.

If $Y_n \rightarrow_{\mathcal{D}} Y$ and $Z_n \rightarrow_{\mathcal{D}} c$, then

- i) $Y_n + Z_n \rightarrow_{\mathcal{D}} Y + c$;
- ii) $Y_n Z_n \rightarrow_{\mathcal{D}} Yc$; and,
- iii) $\frac{Y_n}{Z_n} \rightarrow_{\mathcal{D}} \frac{Y}{c}$.

Definition 5.3 - Convergence in Quadratic Mean

Let $\{Z_n\}_{n \in \mathbb{N}}$ be a sequence of random variables & Z be a random variable.

We say that $\{Z_n\}_{n \in \mathbb{N}}$ Converges in Quadratic Mean to the random variable Z if

$$\lim_{n \rightarrow \infty} \mathbb{E}[(Z_n - Z)^2] = 0$$

N.B. This is denoted $Z_n \rightarrow_{qm} Z$.

Theorem 5.9 - If $Z_n \rightarrow_{qm} Z$ then $Z_n \rightarrow_{\mathbb{P}} Z$

Proof 5.3 - Theorem 5.9

Fix any $\varepsilon > 0$. We have

$$\begin{aligned} \mathbb{P}(|Z_n - Z| > \varepsilon) &= \mathbb{P}(|Z_n - Z|^2 > \varepsilon^2) \\ &\leq \frac{1}{\varepsilon^2} \mathbb{E}[(Z_n - Z)^2] \text{ by Markov's Inequality} \\ &\rightarrow 0 \text{ since } Z_n \rightarrow_{qm} Z. \end{aligned}$$

Hence $Z_n \rightarrow_{\mathbb{P}} Z$. □

5.1 Probabilistic Convergence & Estimators**Definition 5.4 - Consistency of a Sequence of Estimators**

A sequence of estimators, $\{\hat{\theta}_n(\cdot) : \chi^n \rightarrow \Theta\}$, are said to be *Consistent* if

$$\forall \theta \in \Theta \text{ with } \mathbf{X}_n \sim f_n(\cdot; \theta), \hat{\theta}_n(\mathbf{X}_n) \rightarrow_{\mathbb{P}(\cdot; \theta)} \theta$$

Remark 5.3 - Consistency of a Sequence of Estimators

- i) In numerous situations one will talk about the consistency of *the* estimator, *e.g.* for the MLE, but also for the mean, etc. This implicitly refers to the corresponding sequence of MLEs, sequence of means, etc.
- ii) Note the $\mathbb{P}(\cdot; \theta)$ in the limit above, and in particular the dependence on θ . This is often omitted in practice, you should however not forget what the symbols actually mean.
- iii) Quadratic mean / Mean Square convergence \implies consistency.
That is, if the MSE of the estimator converges to 0, the estimator is consistent.

Example 5.2 - Consistency of Flipping Coins

Let $\mathbf{X} \stackrel{\text{iid}}{\sim} \text{Bernoulli}(\theta^*)$ for some $\theta^* \in [0, 1]$.

The maximum likelihood estimate and method of moments for $\hat{\theta}_n$ are the sample mean.

$$\hat{\theta}_n(X_1, \dots, X_n) = \frac{1}{n} \sum_{i=1}^n X_i$$

By the *Weak Law of Large Numbers* we have that *consistency* of $\{\hat{\theta}_n\}$, since $\mathbb{E}(X_1) = \theta^*$.

Example 5.3 - Crude Confidence Interval when Flipping Coins

Let $\mathbf{X} \stackrel{\text{iid}}{\sim} \text{Bernoulli}(\theta^*)$ for some $\theta^* \in [0, 1]$ and define $\hat{\theta}_n := \hat{\theta}_n(X_1, \dots, X_n)$.

We shall produce a *confidence interval* for θ^* .

$$\mathbb{E}(\hat{\theta}_n; \theta^*) = \theta^* \quad \text{and} \quad \text{Var}(\hat{\theta}_n; \theta^*) = \frac{\theta^*(1 - \theta^*)}{n}$$

$$\begin{aligned}
\mathbb{P}\left(|\hat{\theta}_n - \theta^*| \geq \varepsilon; \theta^*\right) &\leq \frac{\theta^*(1-\theta^*)}{n\varepsilon^2} \quad \text{by Chebyshev's Inequality} \\
\text{We don't know } \theta^*, \text{ but can deduce that } \theta^*(1-\theta^*) &\leq \frac{1}{4} \\
\implies \mathbb{P}\left(|\hat{\theta}_n - \theta^*| \geq \varepsilon; \theta^*\right) &\leq \frac{1}{4n\varepsilon^2} \\
&\text{Define } \alpha := \frac{1}{4n\varepsilon^2} \\
\implies \mathbb{P}\left(|\hat{\theta}_n - \theta^*| \geq \frac{1}{2\sqrt{n\alpha}}; \theta^*\right) &\leq \alpha \\
\implies \mathbb{P}\left(\hat{\theta}_n - \frac{1}{2\sqrt{n\alpha}} < \theta^* < \hat{\theta}_n + \frac{1}{2\sqrt{n\alpha}}; \theta^*\right) &\geq 1 - \alpha
\end{aligned}$$

This means the random interval $(\hat{\theta}_n - \frac{1}{2\sqrt{n\alpha}}, \hat{\theta}_n + \frac{1}{2\sqrt{n\alpha}}; \theta^*)$ contains θ^* with probability $1 - \alpha$.

We can note that the interval decreases as n increases, and increases as α decreases. *N.B.* $\hat{\theta}_n$ is a random variable, while θ^* is not.

Example 5.4 - Asymptotically Exact Confidence Interval when Flipping Coins

This is an improvement on the bound produced in **Example 5.3**.

Let $\mathbf{X} \stackrel{\text{iid}}{\sim} \text{Bernoulli}(\theta^*)$ for some $\theta^* \in [0, 1]$, $W \sim \text{Normal}(0, 1)$ and define $\hat{\theta}_n := \hat{\theta}_n(X_1, \dots, X_n)$. We shall show that

$$\frac{\sqrt{n}(\hat{\theta}_n - \theta^*)}{\sqrt{\hat{\theta}_n(1 - \hat{\theta}_n)}} \rightarrow_D W$$

We know that $\text{Var}(X_1) = \theta^*(1 - \theta^*)$.

By the *Weak Law of Large Numbers* $\hat{\theta}_n \rightarrow_{\mathbb{P}} \theta^*$.

By the *Central Limit Theorem*

$$\frac{\sqrt{n}(\hat{\theta}_n - \theta^*)}{\sqrt{\hat{\theta}_n(1 - \hat{\theta}_n)}} \rightarrow_D W$$

$$\text{Define } Y_n = \frac{\sqrt{n}(\hat{\theta}_n - \theta^*)}{\sqrt{\theta^*(1 - \theta^*)}} \text{ and } Z_n = \frac{\sqrt{\theta^*(1 - \theta^*)}}{\sqrt{\hat{\theta}_n(1 - \hat{\theta}_n)}}.$$

By the *Continuous Mapping Theorem* tells us that $Z_n \rightarrow_D 1$ and $Z_n \rightarrow_{\mathbb{P}} 1$.

Hence, by *Slutsky's Theorem*

$$\frac{\sqrt{n}(\hat{\theta}_n - \theta^*)}{\sqrt{\hat{\theta}_n(1 - \hat{\theta}_n)}} = Y_n Z_n \rightarrow_D W$$

This gives us random interval

$$\left(\hat{\theta}_n - z_{\alpha/2} \sqrt{\frac{\hat{\theta}_n(1 - \hat{\theta}_n)}{n}}, \hat{\theta}_n + z_{\alpha/2} \sqrt{\frac{\hat{\theta}_n(1 - \hat{\theta}_n)}{n}} \right)$$

This interval captures θ^* asymptotically (in n) with probability $1 - \alpha$.

N.B. $z_{\alpha} = \Phi^{-1}(1 - \alpha)$ where Φ is the cumulative density function of a $\text{Normal}(0, 1)$.

6 The Fisher Information

Remark 6.1 - Motivation

In the next part of the content we shall show that given $\mathbf{X}_n \stackrel{\text{iid}}{\sim} f(\cdot; \theta^*)$ then for sufficiently regular models

- i) There exists a lower bound on the achievable performance of any estimate of θ^* .
- ii) A scaled & centered sequence of maximum likelihood estimators $\{\hat{\theta}_n(\mathbf{X}_n)\}$ become asymptotically normal as $n \rightarrow \infty$.

Remark 6.2 - Measuring Performance of Estimator

We measure the performance of an estimator $\hat{\theta}$ in terms of variance, since its mean should be θ^* . Lower variance indicates better performance.

Definition 6.1 - The Score Function

Let $\ell(\theta; x) := \ln f(x; \theta)$.

The *Score Function* is a measure of the sensitivity of the likelihood function wrt θ

$$\ell'(\theta; x) := \frac{d}{d\theta} \ell(\theta; x) = \frac{\frac{d}{d\theta} \ln f(x; \theta)}{\ln f(x; \theta)} = \frac{\ln L'(\theta; x)}{\ln L(\theta; x)}$$

Remark 6.3 - θ^* is a turning point of $\ell(\theta; x)$

Note that under the *Fisher Information Regularity Conditions* we have that $\forall \theta \in \Theta$

$$\begin{aligned} \mathbb{E}(\ell'(\theta; X); \theta) &= \int_S \frac{\frac{d}{d\theta} f(x; \theta)}{f(x; \theta)} f(x; \theta) dx \\ &= \int_S \frac{d}{d\theta} f(x; \theta) dx \\ &= \frac{d}{d\theta} \int_S f(x; \theta) dx \\ &= \frac{d}{d\theta} (1) \\ &= 0 \end{aligned}$$

This shows that we expect the derivative to equal 0 at θ^* . Further, this means θ^* is a turning point of the log-likelihood function (hopefully a maximum).

Example 6.1 - Application of Remark 6.3

Let $X \sim \text{Poisson}(\theta)$. Then $f_X(x; \theta) = \frac{\theta^x}{x!} e^{-\theta} \mathbf{1}\{x \in \mathbb{N}\}$.

$$\begin{aligned} \implies \ell(\theta; x) &= -\theta + x \ln \theta - \ln x! \\ \implies \ell'(\theta; x) &= -1 + \frac{x}{\theta} \\ \implies \mathbb{E}(\ell'(\theta; X); \theta) &= -1 + \frac{\theta}{\theta} \\ &= 0 \end{aligned}$$

Definition 6.2 - Fisher Information Regularity Conditions

Let Θ be an open interval in \mathbb{R} and $f(x; \theta)$ be a pmf/pdf.

Below are conditions which a model is required to meet in order to be considered sufficiently regular such that *Fisher Information* can be drawn from it.

- i) Both $L'(\theta; x) = \frac{d}{d\theta} f(x; \theta)$ and $L''(\theta; x) = \frac{d^2}{d\theta^2} f(x; \theta)$ exist for any $x \in \mathcal{X}$.
- ii) $\forall \theta \in \Theta$ the set $S := \{x \in \mathcal{X} : f(x; \theta) > 0\}$ does not depend on $\theta \in \Theta$.
- iii) The identity below exists

$$\int_S \frac{d}{d\theta} f(x; \theta) dx = \frac{d}{d\theta} \int_S f(x; \theta) dx = 0$$

Definition 6.3 - Fisher Information

Fisher Information is a technique for measuring the amount of information that an observable random variable X carries about an unknown parameter θ upon which the probability of X depends.

Let $X \sim f(\cdots; \theta)$. Then the *Fisher Information* for any $\theta \in \Theta$ is

$$I(\theta) := \mathbb{E}(\ell'(\theta; X)^2; \theta) \geq 0$$

N.B. This is the *Expectation of the score, squared* \equiv *Second moment of the score*.

Remark 6.4 - Fisher Information

- i) *Fisher Information* is a function of the parameter, θ , not the data, X .
- ii) $I(\theta)$ can be thought of as being the average *information* brought by a single observation X about θ , assuming $X \sim f(\cdot; \theta)$.
- iii) Since $\forall \theta \in \Theta$, $\mathbb{E}(\ell'(\theta; X); \theta) = 0$ then

$$I(\theta) = \text{Var}(\ell'(\theta; X); \theta)$$

The variance of the score.

Example 6.2 - Fisher Information of Poisson

Let $X \sim \text{Poisson}(\theta)$.

From **Example 6.1** we know that $\ell'(\theta; x) = -1 + \frac{x}{\theta}$. Then

$$\begin{aligned} I(\theta) &= \text{Var}(\ell'(\theta; X); \theta) \\ &= \text{Var}\left(-1 + \frac{X}{\theta}; \theta\right) \\ &= \text{Var}\left(\frac{X}{\theta}; \theta\right) \\ &= \frac{1}{\theta^2} \text{Var}(X; \theta) \\ &= \frac{1}{\theta^2} \cdot \theta \text{ since } X \sim \text{Poisson}(\theta) \\ &= \frac{1}{\theta} \end{aligned}$$

Theorem 6.1 - Alternative Expression of Fisher Information

Let $f(x; \theta)$ be a pmf/pdf which satisfies the conditions of **Definition 6.2**. If

$$\forall \theta \in \Theta \quad \int_{\mathcal{X}} \frac{d^2}{d\theta^2} f(x; \theta) dx = \frac{d}{d\theta} \int_{\mathcal{X}} \frac{d}{d\theta} f(x; \theta) dx$$

Then

$$I(\theta) = -\mathbb{E}\left(\frac{d^2}{d\theta^2} \ell(\theta; X); \theta\right)$$

N.B. $\frac{d}{d\theta} \int_{\mathcal{X}} \frac{d}{d\theta} f(x; \theta) dx = 0$ by the regularity conditions.

Proof 6.1 - Theorem 6.1

By the *Quotient Rule*

$$\begin{aligned} \frac{d^2}{d\theta^2} \ell(\theta; x) &= \frac{d}{d\theta} \frac{\frac{d}{d\theta} f(x; \theta)}{f(x; \theta)} \\ &= \frac{\frac{d^2}{d\theta^2} f(x; \theta)}{f(x; \theta)} - \left(\frac{\frac{d}{d\theta} f(x; \theta)}{f(x; \theta)} \right)^2 \end{aligned}$$

Consequently

$$\begin{aligned} \mathbb{E}\left(\frac{d^2}{d\theta^2} \ell(\theta; X); \theta\right) &= \int_S \frac{\frac{d^2}{d\theta^2} f(x; \theta)}{f(x; \theta)} f(x; \theta) dx - \int_S \left(\frac{\frac{d}{d\theta} f(x; \theta)}{f(x; \theta)} \right)^2 f(x; \theta) dx \\ &= \int_S \frac{d^2}{d\theta^2} f(x; \theta) dx - \int_S \ell'(\theta; x)^2 f(x; \theta) dx \\ &= 0 - \mathbb{E}(\ell'(\theta; X)^2; \theta) \\ &= -I(\theta) \\ \Rightarrow \quad I(\theta) &= -\mathbb{E}\left(\frac{d^2}{d\theta^2} \ell(\theta; X); \theta\right) \end{aligned}$$

□

7 Efficiency and The Cramer-Rao Bound

Definition 7.1 - IID Score Function

Let $\mathbf{X} \stackrel{\text{iid}}{\sim} f(\cdot; \theta)$ for some $\theta \in \Theta$. Then the *Score Function* is

$$\ell'_n(\theta; \mathbf{x}) := \frac{d}{d\theta} \ell_n(\theta; \mathbf{x}) \text{ where } \ell_n(\theta; \mathbf{x}) := \ln f_n(\mathbf{x}; \theta) = \sum_{i=1}^n \ell(\theta; x_i)$$

N.B. $\frac{d}{d\theta} \ell_n(\theta; \mathbf{x}) = \frac{d}{d\theta} \sum \ell(\theta; x_i) = \sum \ell'(\theta; x_i)$.

Definition 7.2 - IID Fisher Information

Let $\mathbf{X} \stackrel{\text{iid}}{\sim} f(\cdot; \theta)$ for some $\theta \in \Theta$. Then the *Fisher Information* is

$$I_n(\theta) := \mathbb{E}(\ell'_n(\theta; \mathbf{X})^2; \theta) = \text{Var}(\ell'_n(\theta; \mathbf{X}); \theta)$$

Theorem 7.1 - Relationship between IID Fisher Information & Fisher Information

Consider the situation where $\forall \theta \in \Theta, f_n(\mathbf{x}; \theta) = \prod_{i=1}^n f(x_i; \theta)$. Then

$$\forall \theta \in \Theta, I_n(\theta) = nI(\theta)$$

Proof 7.1 - Theorem 7.1

Let $\mathbf{X} \stackrel{\text{iid}}{\sim} f(\cdot; \theta)$. Then

$$\begin{aligned} I_n(\theta) &= \text{Var}(\ell'_n(\theta; \mathbf{X}); \theta) \\ &= \text{Var}\left(\sum_{i=1}^n \ell'(\theta; X_i); \theta\right) \\ &= n\text{Var}\left(\sum_{i=1}^n \ell'(\theta; X_1); \theta\right) \\ \implies I_n(\theta) &= nI(\theta) \end{aligned}$$

□

Theorem 7.2 - Cauchy-Schwarz Inequality for Expectation

Let X & Y be real-valued random variables in the same probability space. Then

$$\mathbb{E}(XY)^2 \leq \mathbb{E}(X^2)\mathbb{E}(Y^2)$$

Proof 7.2 - Theorem 7.2

If $\mathbb{E}(Y^2) = 0$ then $\mathbb{P}(Y = 0) = 1$ so $\mathbb{E}(XY) = 0$ and the statement holds.

Thus, assume $\mathbb{E}(Y^2) > 0$ and define $\lambda := \frac{\mathbb{E}(XY)}{\mathbb{E}(Y^2)}$. Then

$$\begin{aligned} 0 &\leq \mathbb{E}(X - \lambda Y)^2 \\ &= \mathbb{E}(X^2) - 2\lambda\mathbb{E}(XY) + \lambda^2\mathbb{E}(Y^2) \\ &= \mathbb{E}(X^2) - 2\frac{\mathbb{E}(XY)^2}{\mathbb{E}(Y^2)} + \frac{\mathbb{E}(XY)^2}{\mathbb{E}(Y^2)} \\ &= \mathbb{E}(X^2) - \frac{\mathbb{E}(XY)^2}{\mathbb{E}(Y^2)} \\ \implies \mathbb{E}(XY)^2 &\leq \mathbb{E}(X^2)\mathbb{E}(Y^2) \end{aligned}$$

□

Theorem 7.3 - Covariance Inequality

Let X and Y be real-valued random variables in the same probability space. Then

$$\text{Cov}(X, Y)^2 \leq \text{Var}(X)\text{Var}(Y)$$

Proof 7.3 - Theorem 7.3

Let $W = X - \mathbb{E}(X)$ and $Z = Y - \mathbb{E}(Y)$ giving $\mathbb{E}(WZ) = \text{Cov}(X, Y)$, $\mathbb{E}(W^2) = \text{Var}(X)$ and $\mathbb{E}(Z^2) = \text{Var}(Y)$.

By applying the *Cauchy-Schwarz inequality* we get

$$\text{Cov}(X, Y)^2 = \mathbb{E}(WZ)^2 \leq \mathbb{E}(W^2)\mathbb{E}(Z^2) = \text{Var}(X)\text{Var}(Y) \iff \text{Cov}(X, Y)^2 \leq \text{Var}(X)\text{Var}(Y)$$

Remark 7.1 - Correlation value

The result in **Theorem 7.3** is the reason why correlation is valued in $[-1, 1]$.

$$\text{Corr}(X, Y) = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}}$$

Theorem 7.4 - Cramer-Rao Inequality - Scalar Parameter

Let $\mathbf{X}_n \stackrel{\text{iid}}{\sim} f(\cdot; \theta)$ and assume the *Fisher Information Regularity Conditions* hold.

Let $\hat{\theta}_n(\cdot)$ be an estimator of θ with expectation $m(\theta) := \mathbb{E}(\hat{\theta}_n(\mathbf{X}_n); \theta)$ which satisfies

$$\forall \theta \in \Theta, \underbrace{\frac{d}{d\theta} \int \hat{\theta}_n(\mathbf{x}) f_n(\mathbf{x}; \theta) d\mathbf{x}}_{\mathbb{E}(\hat{\theta}_n)} = \int \hat{\theta}_n(\mathbf{x}) \frac{d}{d\theta} f_n(\mathbf{x}; \theta) d\mathbf{x}$$

Then

$$\forall \theta \in \Theta, \quad \text{Var}(\hat{\theta}_n(\mathbf{X}_n); \theta) \geq \frac{m'(\theta)^2}{nI(\theta)}$$

Proof 7.4 - Theorem 7.4

We notice that

$$\begin{aligned} m'(\theta) &= \frac{d}{d\theta} \mathbb{E}(\hat{\theta}_n(\mathbf{X}_n); \theta) \\ &= \frac{d}{d\theta} \int_{S^n} \hat{\theta}_n(\mathbf{x}_n) f_n(\mathbf{x}_n; \theta) d\mathbf{x}_n \end{aligned}$$

The clever part of this proof is to observe that

$$\begin{aligned} \text{Var}(\hat{\theta}_n(\mathbf{X}_n); \theta) nI(\theta) &= \text{Var}(\hat{\theta}_n(\mathbf{X}_n); \theta) \text{Var}(\ell'_n(\theta; \mathbf{X}_n); \theta) \\ &\geq \text{Cov}(\hat{\theta}_n(\mathbf{X}_n), \ell'_n(\theta; \mathbf{X}_n); \theta)^2 \text{ by Covariance Inequality} \end{aligned}$$

Thus

$$\begin{aligned} \text{Cov}(\hat{\theta}_n(\mathbf{X}_n), \ell'_n(\theta; \mathbf{X}_n); \theta)^2 &= \mathbb{E}(\hat{\theta}_n(\mathbf{X}_n) \ell'_n(\theta; \mathbf{X}_n); \theta) - \mathbb{E}(\hat{\theta}_n(\mathbf{X}_n); \theta) \mathbb{E}(\ell'_n(\theta; \mathbf{X}_n); \theta) \\ &= \mathbb{E}(\hat{\theta}_n(\mathbf{X}_n) \ell'_n(\theta; \mathbf{X}_n); \theta) - \mathbb{E}(\hat{\theta}_n(\mathbf{X}_n); \theta) \times 0 \\ &= \mathbb{E}(\hat{\theta}_n(\mathbf{X}_n) \ell'_n(\theta; \mathbf{X}_n); \theta) \\ &= \int_{S^n} \hat{\theta}_n(\mathbf{x}_n) \ell'_n(\theta; \mathbf{x}_n) f_n(\mathbf{x}_n; \theta) d\mathbf{x}_n \\ &= \int_{S^n} \hat{\theta}_n(\mathbf{x}_n) \frac{\frac{d}{d\theta} f_n(\mathbf{x}_n; \theta)}{f_n(\mathbf{x}_n; \theta)} f_n(\mathbf{x}_n; \theta) d\mathbf{x}_n \\ &= \int_{S^n} \hat{\theta}_n(\mathbf{x}_n) \frac{d}{d\theta} f_n(\mathbf{x}_n; \theta) \\ &= \frac{d}{d\theta} \int_{S^n} \hat{\theta}_n(\mathbf{x}_n) f_n(\mathbf{x}_n; \theta) d\mathbf{x}_n \text{ by regularity assumption} \\ &= m'(\theta) \\ \implies \text{Var}(\hat{\theta}_n(\mathbf{X}_n); \theta) nI(\theta) &\geq m'(\theta)^2 \end{aligned}$$

Proposition 7.1 - Useful result from Cramer-Rao Inequality

If $\hat{\theta}_n(\mathbf{X}_n)$ is an unbiased estimator (i.e. $m(\theta) = \theta$) then

$$\text{Var}(\hat{\theta}_n(\mathbf{X}_n); \theta) = \text{MSE}(\hat{\theta}_n(\mathbf{X}_n); \theta) \geq \frac{1}{nI(\theta)}$$

This shows there is a lower bound on the possible performance of an estimator.

Definition 7.3 - Efficient Estimator

An *Estimator* is said to be *Efficient* when its variance is equal to the *Cramer-Rao lower bound* $\forall \theta^*$.

Example 7.1 - Efficient Coin Flipping

Let $\mathbf{X} \stackrel{\text{iid}}{\sim} \text{Bernoulli}(\theta)$ with $\theta \in [0, 1]$, this corresponds to flipping a coin n times and considering each flip the random variable $X : \{H, T\} \rightarrow \{0, 1\}$ such that $X(H) = 1$ and $X(T) = 0$ with probability distribution such that $\mathbb{P}(X = 1; \theta) = \theta$ and $\mathbb{P}(X = 0; \theta) = 1 - \theta$. We consider the intuitive estimator of θ

$$\hat{\theta}_n := \hat{\theta}_n(\mathbf{X}_n) := \frac{1}{n} \sum_{i=1}^n X_i$$

The estimator is unbiased $\forall n \in \mathbb{N}$ and its variance is

$$\text{Var}(\hat{\theta}_n; \theta) = \frac{\text{Var}(X_1; \theta)}{n} = \frac{\mathbb{E}(X_1^2; \theta) - \mathbb{E}(X_1; \theta)^2}{n} = \frac{\theta - \theta^2}{n} = \frac{\theta(1 - \theta)}{n}$$

Now we consider the *Cramer-Rao bound*

$$\begin{aligned} \text{We find } L(\theta; x) &= \theta^x (1 - \theta)^{1-x} \\ \implies \ell(\theta; x) &= x \ln \theta + (1 - x) \ln(1 - \theta) \\ \implies \ell'(\theta; x) &= \frac{x}{\theta} - \frac{1-x}{1-\theta} \\ \implies \ell''(\theta; x) &= -\frac{x}{\theta^2} - \frac{1-x}{(1-\theta)^2} \end{aligned}$$

Thus we can use $I(\theta) = -\mathbb{E}(\ell''(\theta; X); \theta)$

$$\begin{aligned} \implies I(\theta) &= -\mathbb{E}\left(-\frac{X}{\theta^2} - \frac{1-X}{(1-\theta)^2}; \theta\right) \\ &= \mathbb{E}\left(\frac{X}{\theta^2} + \frac{1-X}{(1-\theta)^2}; \theta\right) \\ &= \frac{\theta}{\theta^2} + \frac{1-\theta}{(1-\theta)^2} \\ &= \frac{1}{\theta} + \frac{1}{1-\theta} \\ &= \frac{1}{\theta(1-\theta)} \\ I_n(\theta) &= nI(\theta) \text{ Since } X_1, X_2, \dots \text{ are iid} \end{aligned}$$

The *Cramer-Rao bound* for the variance is

$$\frac{1}{nI(\theta)} = \frac{\theta(1 - \theta)}{n}$$

Thus our estimator is efficient.

8 Asymptotic Distribution of the Maximum Likelihood Estimator

Theorem 8.1 -

Suppose that $\mathbf{X}_n \stackrel{\text{iid}}{\sim} f(\cdot; \theta^*)$ for some $\theta^* \in \Theta$ and assume that

- i) The sequence of maximum likelihood estimators $\{\hat{\theta}_n(\mathbf{X}_n)\}$ is consistent;
- ii) The *Fisher Information Regularity Conditions* (**Definition 6.2**) hold and $I(\theta^*) = -\mathbb{E}[\ell''(\theta; X); \theta] > 0$.

- iii) $\exists C(\cdot) : \mathcal{X} \rightarrow [0, \infty)$ such that $\mathbb{E}(C(X_1); \theta^*) < \infty$, $\Xi \subset \Theta$ an open set containing θ^* and $\Delta(\cdot) : \Xi \rightarrow [0, \infty)$ continuous at 0 st $\Delta(0) = 0$, st $\forall \theta, \theta', x \in \Xi^2 \times \mathcal{X}$.

$$|\ell''(\theta; x) - \ell''(\theta'; x)| \leq C(x)\Delta(\theta - \theta')$$

Then $\forall \theta^* \in \Theta$

$$\sqrt{nI(\theta^*)}(\hat{\theta}_n(\mathbf{X}_n) - \theta^*) \rightarrow_{\mathcal{D}(\cdot; \theta^*)} Z \sim \text{Normal}(0, 1)$$

Theorem 8.2 -

Under the conditions of **Theorem 8.1**, with $\hat{\theta}_n := \hat{\theta}_n(\mathbf{X})$ the maximum likelihood estimator

$$\ell'_n(\hat{\theta}_n; \mathbf{X}) = \ell'_n(\theta^*; \mathbf{X}) + (\hat{\theta}_n - \theta^*)\{\ell''_n(\theta^*; \mathbf{X}) + R_n\}$$

where $\frac{1}{n}R_n \rightarrow_{\mathbb{P}(\cdot; \theta^*)} 0$.

Proof 8.1 - Theorem 8.1

By **Theorem 8.2** $\ell'_n(\hat{\theta}_n; \mathbf{X}) = \ell'_n(\theta^*; \mathbf{X}) + (\hat{\theta}_n - \theta^*)\{\ell''_n(\theta^*; \mathbf{X}) + R_n\}$ where $\frac{1}{n}R_n \rightarrow_{\mathbb{P}(\cdot; \theta^*)} 0$.

Since $\hat{\theta}_n$ is the maximum likelihood estimator & the *Fisher Information Regularity Conditions* hold, the score at $\ell'(\hat{\theta}_n; X) = 0$.

Hence, $0 = \ell''(\hat{\theta}_n; X) = \ell''(\theta; X) + (\hat{\theta}_n - \theta^*)\{\ell''(\theta; X) + R_n\}$.

Rearranging & rescaling by \sqrt{n} gives

$$\sqrt{n}(\hat{\theta}_n - \theta^*) = \frac{\frac{1}{\sqrt{n}}\ell'(\theta^*; X)}{-\frac{1}{\sqrt{n}}\{\ell''(\theta^*; X) + R_n\}} =: \frac{U_n}{V_n - \frac{R_n}{n}}$$

Recall that $\ell'_n(\theta^*; X) = \sum_{i=1}^n \ell'(\theta; X_i)$ and $\ell''_n(\theta^*; X) = \sum_{i=1}^n \ell''(\theta^*; X_i)$.

Since $\mathbb{E}(\ell'(\theta^*; X_i); \theta^*) = 0$ and $\text{Var}(\ell'(\theta^*; X_i); \theta^*) = I(\theta^*)$

$\Rightarrow U_n \rightarrow_{\mathcal{D}(\cdot; \theta^*)} U \sim \text{Normal}(0, I(\theta^*))$ by the *Central Limit Theorem*.

We observed that $V_n \rightarrow_{\mathbb{P}(\cdot; \theta^*)} I(\theta^*)$ by the *Weak Law of Large Numbers* since $\mathbb{E}(-\ell''(\theta^*; X_i); \theta^*) = I(\theta^*)$.

It follows that $V_n - \frac{1}{n}R_n \rightarrow_{\mathbb{P}(\cdot; \theta^*)} I(\theta^*)$ by *Slutsky's Theorem*.

Using *Slutsky's Theorem* again

$$\sqrt{n}(\hat{\theta}_n - \theta^*) = \frac{U_n}{V_n - \frac{1}{n}R_n} \rightarrow_{\mathcal{D}(\cdot; \theta^*)} \frac{\sqrt{I(\theta^*)}}{I(\theta^*)} Z \text{ where } Z \sim \text{Normal}(0, 1)$$

We can rewrite this as

$$\sqrt{nI(\theta^*)}(\hat{\theta}_n - \theta^*) \rightarrow_{\mathcal{D}(\cdot; \theta^*)} Z \sim \text{Normal}(0, 1)$$

Proof 8.2 - Theorem 8.2

This is a non-examinable, sketch proof of Theorem 8.2.

By the regularity conditions and the mean value theorem

$$\frac{\ell'_n(\theta; \mathbf{x}) - \ell'_n(\theta^*; \mathbf{x})}{\theta - \theta^*} = \ell''_n(\tilde{\theta}; \mathbf{x})$$

for some $\tilde{\theta} \in (\theta, \theta^*)$. Hence, we deduce that

$$\begin{aligned} \ell'_n(\theta; \mathbf{x}) - \ell'_n(\theta^*; \mathbf{x}) &= (\theta - \theta^*)\ell''_n(\tilde{\theta}; \mathbf{x}) \\ &= (\theta - \theta^*)\{\ell''_n(\theta^*; \mathbf{x}) + [\ell''_n(\tilde{\theta}; \mathbf{x}) - \ell''_n(\theta^*; \mathbf{x})]\} \\ &= (\theta - \theta^*)\{\ell''_n(\theta; \mathbf{x}) + R_n(\theta, \theta^*, \mathbf{x})\} \end{aligned}$$

Now we replace θ with the maximum likelihood estimator $\hat{\theta}_n := \hat{\theta}_n(\mathbf{X})$. We find

$$\ell'(\hat{\theta}_n; \mathbf{X}) = \ell'_n(\theta^*; \mathbf{X}) + (\hat{\theta}_n - \theta^*)\{\ell''_n(\theta^*; \mathbf{X}) + R_n(\hat{\theta}_n, \theta^*, \mathbf{x})\}$$

and we need to analyse R_n .

Since $\hat{\theta}_n \rightarrow_{\mathbb{P}(\cdot; \theta^*)} \theta^*$ we can take n large enough that $\mathbb{P}(\hat{\theta}_n \in \Xi; \theta^*)$ with arbitrarily high probability.

On the event $\{\hat{\theta} \in \Xi\}$ and we have $\{\tilde{\theta}_n \in \Xi\}$ since $\tilde{\theta}_n \in (\hat{\theta}_n, \theta^*)$ and

$$\begin{aligned} |\frac{1}{n}R_n| &= \frac{1}{n} |\ell''_n(\tilde{\theta}_n; \mathbf{X}) - \ell''_n(\theta^*; \mathbf{X})| \\ &= \frac{1}{n} \left| \sum_{i=1}^n \ell''(\tilde{\theta}_n; X_i) - \ell''(\theta^*; X_i) \right| \\ &\leq \frac{1}{n} \sum_{i=1}^n \left| \ell''(\tilde{\theta}_n; X_i) - \ell''(\theta^*; X_i) \right| \\ &\leq \Delta(\tilde{\theta}_n - \theta^*) \left\{ \frac{1}{n} \sum_{i=1}^n C(X_i) \right\} \end{aligned}$$

from the smoothness condition on ℓ'' .

From the *Weak Law of Large Numbers*

$$\frac{1}{n} \sum_{i=1}^n C(X_i) \rightarrow_{\mathbb{P}(\cdot; \theta^*)} \mathbb{E}(C(X_1); \theta^*) < \infty$$

and from the consistency of $\{\hat{\theta}_n\}$ and $\{\tilde{\theta}_n\}$ and continuity of $\Delta(\cdot)$ we have by the *Continuous Mapping Theorem*

$$\Delta(\tilde{\theta}_n - \theta^*) \rightarrow_{\mathbb{P}(\cdot; \theta^*)} 0$$

Hence, $\frac{1}{n}R_n \rightarrow_{\mathbb{P}(\cdot; \theta^*)} 0$ □

Definition 8.1 - Asymptically Efficient

A sequence of estimators $\{\hat{\theta}_n(\mathbf{X})\}$ is *Asymptotically Efficient* if either its mean-squared error converges to the *Cramer-Rao Lower Bound*

$$\forall \theta \in \Theta, n\text{MSE}(\hat{\theta}_n(\mathbf{X}_n); \theta) \xrightarrow{n \rightarrow \infty} \frac{1}{I(\theta)}$$

or $\hat{\theta}_n$ is *Asymptotically Normally Distributed* in the sense of **Theorem 8.1**

$$\forall \theta \in \Theta, \sqrt{nI(\theta)}(\hat{\theta} - \theta) \rightarrow_{\mathcal{D}(\cdot; \theta)} Z$$

N.B. The variance of $\frac{Z}{\sqrt{nI(\theta^*)}}$ is exactly $\frac{1}{nI(\theta)}$.

Theorem 8.3 -

Under the conditions of **Theorem 8.1** the maximum likelihood estimator is *asymptotically efficient*.

Definition 8.2 - Regular Statistical Model

Any *Statistical Model* which satisfies the condition of **Theorem 8.1** is a *Regular Statistical Model*.

Remark 8.1 - Why use MLE over others

Due to the *Asymptotic Efficiency* of maximum likelihood estimators it is better to use them in *Regular Statistical Models*.

9 Confidence Sets Around the Maximum Likelihood Estimator

Definition 9.1 - Coverage of an Interval

Let $\mathbf{X} \sim f_n(\cdot; \theta)$, $\theta \in \Theta = \mathbb{R}$, $L(\cdot) : \mathcal{X}^n \rightarrow \Theta$ and $U(\cdot) : \mathcal{X}^n \rightarrow \Theta$ where $\forall \mathbf{x} \in \mathcal{X}^n$, $L(\mathbf{x}) < U(\mathbf{x})$. Then, $\forall \theta \in \Theta$ the coverage $C_{\mathcal{I}}(\theta)$ of the random interval $\mathcal{I}(\mathbf{X}) := [L(\mathbf{X}), U(\mathbf{X})]$ at θ is

$$C_{\mathcal{I}}(\theta) := \mathbb{P}(\theta \in [L(\mathbf{X}), U(\mathbf{X})]; \theta) = \mathbb{P}(L(\mathbf{X}) \leq \theta \leq U(\mathbf{X}); \theta)$$

Remark 9.1 - Coverage of an Interval in Words

$C_{\mathcal{I}}(\theta)$ is the probability that the deterministic quantity θ falls into the random interval $\mathcal{I}(\mathbf{X})$ under the probability distribution $\mathbb{P}(\cdot; \theta)$ where $\mathbf{X} \sim f_n(\cdot; \theta)$.

Remark 9.2 - Multi-Dimensional Coverage

We can extend *Coverage of an Interval* to the multi-dimensional case by considering confidence sets and then considering the probability $\mathbb{P}(\theta \in \mathcal{I}(\mathbf{X}); \theta)$.

Definition 9.2 - Confidence Interval

$\forall \alpha \in [0, 1]$ we say that an interval $\mathcal{I}(\mathbf{X}) := [L(\mathbf{X}), U(\mathbf{X})]$ is a $1 - \alpha$ confidence interval if $\forall \theta \in \Theta$ its coverage is at least $1 - \alpha$ or more formally $\inf_{\theta \in \Theta} C_{\mathcal{I}}(\theta) \geq 1 - \alpha$.

Remark 9.3 - Exact Confidence Interval

If $C_{\mathcal{I}}(\theta) = 1 - \alpha \forall \theta \in \Theta$ then \mathcal{I} is an exact $1 - \alpha$ confidence interval.

Definition 9.3 - Observed Confidence Interval

For an interval $\mathcal{I}(\cdot) = [L(\cdot), U(\cdot)]$ with $L : \mathcal{X}^n \rightarrow \Theta$ and $U : \mathcal{X}^n \rightarrow \Theta$, and a realisation \mathbf{x} , the corresponding *Observed Confidence Interval* is $\mathcal{I}(\mathbf{x})$.

N.B. Nothing interesting can be said about the probability that $\theta \in \mathcal{I}(\mathbf{x})$ since θ and $\mathcal{I}(\mathbf{x})$ are deterministic.

Notation 9.1 - Quantile of Normal(0, 1)

For any $\beta \in (0, 1)$ let $z_{\beta} \in \mathbb{R}$ be such that for $Z \sim \text{Normal}(0, 1)$, $1 - \Phi(z_{\beta}) = \mathbb{P}(Z > z_{\beta}) = \beta$.

Example 9.1 - Confidence interval for the mean of a Normal Distribution

Let $\mathbf{X} \stackrel{\text{iid}}{\sim} \text{Normal}(\mu, \sigma^2)$ for $\theta = (\mu, \sigma^2) \in \mathbb{R} \times \mathbb{R}^{\geq 0}$ and where σ^2 is known.

Consider the estimator $\hat{\mu}_n = \hat{\mu}_n(\mathbf{X}) = \frac{1}{n} \sum_{i=1}^n X_i$ of μ . Then we know that the following non-asymptotic result holds.

We have $\frac{1}{n} \sum_{i=1}^n X_i \sim \text{Normal}(\mu, \frac{\sigma^2}{n})$. Thus

$$\frac{\frac{1}{n} \sum_{i=1}^n X_i - \mu}{\sqrt{\sigma^2/n}} \sim \text{Normal}(0, 1)$$

Then

$$\begin{aligned} \forall \alpha \in (0, 1) \quad , \quad & \mathbb{P} \left(z_{1-\alpha/2} \leq \frac{\hat{\mu}_n(\mathbf{X}) - \mu}{\sqrt{\sigma^2/n}} \leq z_{\alpha/2}; \mu \right) \\ &= \mathbb{P} \left(\frac{\hat{\mu}_n(\mathbf{X}) - \mu}{\sqrt{\sigma^2/n}} \leq z_{\alpha/2} \right) - \mathbb{P} \left(\frac{\hat{\mu}_n(\mathbf{X}) - \mu}{\sqrt{\sigma^2/n}} \leq z_{1-\alpha/2} \right) \\ &= \left(1 - \frac{\alpha}{2} \right) - \left(1 - \left(1 - \frac{\alpha}{2} \right) \right) \\ &= 1 - \alpha \end{aligned}$$

By symmetry we notice that $z_{1-\frac{\alpha}{2}} = -z_{\alpha/2}$.

By rearranging we have the equivalence of events

$$\left\{ -z_{\alpha/2} \leq \frac{\hat{\mu}_n(\mathbf{X}) - \mu}{\sqrt{\sigma^2/n}} \leq z_{\alpha/2} \right\} = \left\{ \hat{\mu}_n(\mathbf{X}) - z_{\alpha/2} \frac{\sigma}{\sqrt{n}} \leq \mu \leq \hat{\mu}_n(\mathbf{X}) + z_{\alpha/2} \frac{\sigma}{\sqrt{n}} \right\}$$

To rearrange we separate into two events & treat then separately

$$\begin{aligned}\left\{\frac{\hat{\mu}_n(\mathbf{X}) - \mu}{\sigma/\sqrt{n}} \leq z_{\alpha/2}\right\} &= \left\{\frac{\hat{\mu}_n(\mathbf{X})}{\sigma/\sqrt{n}} - z_{\alpha/2} \leq \frac{\mu}{\sigma/\sqrt{n}}\right\} \\ &= \left\{\mu \geq \hat{\mu}_n(\mathbf{X}) - \frac{\sigma}{\sqrt{n}} z_{\alpha/2}\right\}\end{aligned}$$

Similarly

$$\begin{aligned}\left\{-z_{\alpha/2} \leq \frac{\hat{\mu}_n(X) - \mu}{\sigma/\sqrt{n}}\right\} &= \left\{\frac{\mu}{\sigma/\sqrt{n}} \leq \frac{\hat{\mu}_n(X)}{\sigma/\sqrt{n}} + z_{\alpha/2}\right\} \\ &= \left\{\mu \leq \hat{\mu}_n(X) + z_{\alpha/2} \frac{\sigma}{\sqrt{n}}\right\}\end{aligned}$$

So the interval $\mathcal{I}(X) = [L(X), U(X)]$ where $L(\mathbf{x}) = \bar{x} - z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$ and $U(\mathbf{x}) = \bar{x} + z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$ is an $1 - \alpha$ exact confidence interval.

Remark 9.4 - Confidence Intervals with unknown σ^2

When σ^2 is unknown we can define $\{\hat{\sigma}_n^2\}_{n \in \mathbb{N}}$ to be a consistent sequence of estimators of σ^2 (e.g. the sample variance)

$$\hat{\sigma}_n^2 := \frac{1}{n-1} \sum_{i=1}^n (X_i - \hat{\mu}_n(\mathbf{X}))^2$$

10 Asymptotic Approximation of Confidence Intervals

Theorem 10.1 -

Assume $\mathbf{X} \sim f(\cdot; \theta^*)$. Let $\{\hat{\theta}_n\}_{n \in \mathbb{N}}$ be a consistent sequence of estimators of θ^* and assume that $\{\hat{\theta}_n\}$ is asymptotically normal in the sense that

$$\exists \sigma^2 > 0 \text{ st } \frac{\hat{\theta}_n(\mathbf{X}) - \theta^*}{\sqrt{\sigma^2/n}} \rightarrow_{\mathcal{D}(\cdot; \theta^*)} Z \sim \text{Normal}(0, 1)$$

Then $\forall \alpha \in (0, 1)$, $\mathcal{I}_n(\mathbf{X}) = [L_n(\mathbf{X}), U_n(\mathbf{X})]$ is an asymptotically exact $1 - \alpha$ confidence interval, where $L_n(\mathbf{x}) := \hat{\theta}_n(\mathbf{x}) - z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$ and $U_n(\mathbf{x}) := \hat{\theta}_n(\mathbf{x}) + z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$.

Proof 10.1 - Theorem 10.1

Let $\{W_n\}_{n \in \mathbb{N}}$ be defined by $W_n := \frac{\hat{\theta}_n(X) - \theta^*}{\sqrt{\sigma^2/n}}$.

Since $W_n \rightarrow_{\mathcal{D}(\cdot; \theta^*)} Z \sim \text{Normal}(0, 1)$ we have

$$\begin{aligned}\mathbb{P}(-z_{\alpha/2} \leq W_n \leq z_{\alpha/2}) &= F_{W_n}(z_{\alpha/2}) - F_{W_n}(-z_{\alpha/2}) \\ &\xrightarrow{n \rightarrow \infty} \Phi(z_{\alpha/2}) - \Phi(-z_{\alpha/2}) \\ &= 1 - \alpha\end{aligned}$$

Similarly to before we have the equivalence of events

$$\{-z_{\alpha/2} \leq W_n \leq z_{\alpha/2}\} = \left\{\hat{\theta}_n - z_{\alpha/2} \frac{\sigma}{\sqrt{n}} \leq \theta^* \leq \hat{\theta}_n + z_{\alpha/2} \frac{\sigma}{\sqrt{n}}\right\}$$

So $\lim_{n \rightarrow \infty} \mathbb{P}\left(\hat{\theta}_n(X) - z_{\alpha/2} \frac{\sigma}{\sqrt{n}} \leq \theta^* \leq \hat{\theta}_n(X) + z_{\alpha/2} \frac{\sigma}{\sqrt{n}}; \theta^*\right) = 1 - \alpha$

Remark 10.1 - Theorem 10.1

The confidence interval is only asymptotically exact. For finite n , the overage of the confidence interval will be different from $1 - \alpha$ but the difference will converge to 0 as n increases. In practice σ^2 may be unknown, in these cases substitute for a consistent sequence of estimators of σ^2 .

Theorem 10.2 -

Assume $\mathbf{X} \sim f(\cdot; \theta^*)$ let $\{\hat{\theta}_n\}_{n \in \mathbb{N}}$ be a consistent sequence of estimators of θ^* and assume that $\{\hat{\theta}_n\}$ is asymptotically normal in the sense that

$$\exists \sigma^2 > 0 \text{ st } \frac{\hat{\theta}_n(\mathbf{X}) - \theta^*}{\sqrt{\sigma^2/n}} \xrightarrow{\text{mathcal{D}}(\cdot; \theta^*)} Z \sim \text{Normal}(0, 1)$$

Assume also that $\{\hat{\sigma}_n^2\}_{n \in \mathbb{N}}$ is a consistent sequence of estimators of σ^2 . Then $\forall \alpha \in (0, 1)$, $\mathcal{I}_n(\mathbf{X}) = [L_n(\mathbf{X}), U_n(\mathbf{X})]$ is an asymptotically exact $1 - \alpha$ confidence interval, where $L_n(\mathbf{x}) := \hat{\theta}_n(\mathbf{x}) - z_{\alpha/2} \sqrt{\hat{\sigma}_n^2(\mathbf{x})/n}$ and $U_n(\mathbf{x}) := \hat{\theta}_n(\mathbf{x}) + z_{\alpha/2} \sqrt{\hat{\sigma}_n^2(\mathbf{x})/n}$.

Proof 10.2 - Theorem 10.2

$$\text{Define } W_n := \frac{\hat{\theta}_n - \theta^*}{\sqrt{\hat{\sigma}_n^2(X)/n}} = \frac{\hat{\theta}_n(X) - \theta^*}{\sqrt{\sigma^2/n}} - \sqrt{\frac{\sigma^2}{\hat{\sigma}_n^2(X)}}.$$

By consistency of $\{\hat{\sigma}_n^2\}_{n \in \mathbb{N}}$ and the *Continuous Mapping Theorem*

$$\sqrt{\frac{\sigma^2}{\hat{\sigma}_n^2(X)}} \xrightarrow{\mathbb{P}(\cdot; \theta^*)} 1$$

By *Slutsky's Theorem*

$$W_n \xrightarrow{\mathcal{D}(\cdot; \theta^*)} Z \sim \text{Normal}(0, 1)$$

The rest of the proof is the same as for **Theorem 10.1**.

Remark 10.2 - Theorem 10.2

For a given n the quality of the normal approximation will be affected by this additional approximation. One may find that for less accurate estimators of σ^2 , the n required for the confidence interval to have almost the right coverage will be higher.

11 Estimating the Information for Maximum Likelihood Estimates

Remark 11.1 - Applying Theorem 10.2 to sequences of MLEs for regular statistical models

When dealing with *Maximum Likelihood Estimators* for regular statistical models we have that $\sigma^2 = 1/I(\theta^*)$ thus

$$\sqrt{nI(\theta^*)}(\hat{\theta}_n - \theta^*) \xrightarrow{\mathcal{D}(\cdot; \theta^*)} Z \sim \text{Normal}(0, 1)$$

However the *Fisher Information* is unknown so we consider two cases

- i) When the expectation, $I(\theta^*) = -\mathbb{E}(\ell''(\theta^*; X_1); \theta^*)$, can be calculated. In this case we replace θ^* with $\hat{\theta}_n$ in the equation.
- ii) When the expectation **cannot** be calculated we invoke the *Weak Law of Large Numbers* and consider the sequence of estimators, $J_n(\hat{\theta}_n) := -\frac{1}{n} \sum_{i=1}^n \ell''(\hat{\theta}_n; X_i)$.

Theorem 11.1 - Case i)

Assume $\{\hat{\theta}_n\}$ is a sequence of *Maximum Likelihood Estimators* st $\hat{\theta}_n \xrightarrow{\mathbb{P}(\cdot; \theta^*)} \theta^*$ and I is a continuous function of θ . Then $I(\hat{\theta}_n) \xrightarrow{\mathbb{P}(\cdot; \theta^*)} I(\theta^*)$.

N.B. The proof of this follows directly from the *Continuous Mapping Function*.

Remark 11.2 - Theorem 11.1

It is only necessary for I to be continuous in the neighbourhood of θ^* . This is due to an extension of the *Continuous Mapping Theorem* that states

If $X_n \rightarrow_{\mathbb{P}} X$ and g is a function with discontinuity set D then
 $\mathbb{P}(X \in D) = 0 \implies (X_n) \rightarrow_{\mathbb{P}} g(X).$

Theorem 11.2 - Case ii)

Assume that $\{\hat{\theta}_n\}$ is a sequence of *Maximum Likelihood Estimators* st

- i) $\hat{\theta}_n \rightarrow_{\mathbb{P}(\cdot; \theta^*)} \theta^*$;
- ii) $I(\theta) = -\mathbb{E}(\ell''(\theta; X); \theta) \forall \theta \in \Theta$
- iii) $\exists C : \mathcal{X} \rightarrow [0, \infty)$ st $\mathbb{E}(C(X_1); \theta^*) < \infty$, $\Xi \subset \Theta$ is an open set containing θ^* and $\Delta(\cdot) : \Xi \rightarrow [0, \infty)$ is continuous at 0 st $\Delta(0) = 0$, and st $\forall \theta, \theta', x \in \Xi^2 \times \mathcal{X} \mid \ell''(\theta; x) - \ell''(\theta'; x) \mid \leq C(x)\Delta(\theta - \theta')$

Then

$$J_n(\hat{\theta}_n) \rightarrow_{\mathbb{P}(\cdot; \theta^*)} I(\theta^*)$$

Proof 11.1 - Theorem 11.2

Consider the following decomposition

$$\begin{aligned} J_n(\hat{\theta}) - I(\theta^*) &= -\frac{1}{n} \sum_{i=1}^n \ell''(\hat{\theta}_n; X_i) - I(\theta^*) \\ &= T_1 + T_2 \\ \text{Where } T_1 &= -\frac{1}{n} \sum_{i=1}^n \ell''(\hat{\theta}_n; X_i) + \frac{1}{n} \sum_{i=1}^n \ell''(\theta^*; X_i) \\ \text{and } T_2 &= -\left\{ \frac{1}{n} \sum_{i=1}^n \ell''(\theta^*; X_i) \right\} - I(\theta^*) \end{aligned}$$

Now the first term can be upper bounded as follows (for sufficiently large n , with arbitrary large probabilitiy the second inequality holds)

$$\begin{aligned} |T_1| &= \left| -\frac{1}{n} \sum_{i=1}^n \ell''(\hat{\theta}_n; X_i) + \frac{1}{n} \sum_{i=1}^n \ell''(\theta^*; X_i) \right| \\ &\leq \frac{1}{n} \sum_{i=1}^n \left| \ell''(\hat{\theta}_n; X_i) - \ell''(\theta^*; X_i) \right| \\ &\leq \Delta(\theta_n - \theta^*) \frac{1}{n} \sum_{i=1}^n C(X_i) \end{aligned}$$

By the *Weak Law of Large Numbers*

$$\frac{1}{n} \sum_{i=1}^n C(X_i) \rightarrow_{\mathbb{P}(\cdot; \theta^*)} \mathbb{E}(C(X_1); \theta^*)$$

by the assumed consistency of $\{\hat{\theta}_n\}_{n \in \mathbb{N}}$ and continuity of Δ we have that

$$\Delta(\hat{\theta}_n - \theta^*) \rightarrow_{\mathbb{P}(\cdot; \theta^*)} 0$$

Consequently $T_1 \xrightarrow{n \rightarrow \infty}_{\mathbb{P}(\cdot; \theta^*)} 0$.

By the *Weak Law of Large Numbers* we have

$$\begin{aligned} &-\frac{1}{n} \sum_{i=1}^n \ell''(\theta^*; X_i) \rightarrow_{\mathbb{P}(\cdot; \theta^*)} I(\theta^*) \\ \implies T_2 &= -\frac{1}{n} \sum_{i=1}^n \ell''(\theta^*; X_i) - I(\theta^*) \rightarrow_{\mathbb{P}(\cdot; \theta^*)} 0 \end{aligned}$$

Since $T_1 \xrightarrow{n \rightarrow \infty}_{\mathbb{P}(\cdot; \theta^*)} 0$ and $T_2 \xrightarrow{n \rightarrow \infty}_{\mathbb{P}(\cdot; \theta^*)} 0$ we deduce from the earlier decomposition that

$$J_n(\hat{\theta}_n) \rightarrow_{\mathbb{P}(\cdot; \theta^*)} I(\theta^*)$$

□

Remark 11.3 - Summary

Whenever **Theorem 8.1** holds for a sequence of *Maximum Likelihood Estimators*

$$i.e. \sqrt{nI(\theta^*)}(\hat{\theta}_n - \theta^*) \rightarrow_{\mathcal{D}(\cdot; \theta^*)} Z \sim \text{Normal}(0, 1)$$

we can replace $I(\theta^*)$ with one of two options

i) $I(\hat{\theta}_n)$ whenever

- (a) $I(\theta)$ is continuous in a neighbourhood of θ^* ; and,
- (b) The interval $[L(\mathbf{X}), U(\mathbf{X})]$ with $L(\mathbf{x}) := \hat{\theta}_n - z_{\alpha/2} \sqrt{nI(\hat{\theta})}$ and $U(\mathbf{x}) := \hat{\theta}_n + z_{\alpha/2} \sqrt{nI(\hat{\theta})}$ is an asymptotically exact $1 - \alpha$ confidence interval for θ^* .

ii) $J_n(\hat{\theta}_n) := -\frac{1}{n} \sum_{i=1}^n \ell''(\hat{\theta}_n; X_i)$ whenever

- (a) The assumptions of **Theorem 11.2** hold; and,
- (b) The interval $[L(\mathbf{X}), U(\mathbf{X})]$ with $L(\mathbf{x}) := \hat{\theta}_n - z_{\alpha/2} \sqrt{nJ_n(\hat{\theta}_n)}$ and $U(\mathbf{x}) := \hat{\theta}_n + z_{\alpha/2} \sqrt{nJ_n(\hat{\theta}_n)}$ is an asymptotically exact $1 - \alpha$ confidence interval for θ^* .

Example 11.1 - Coin Flipping

Here the new results for this chapter are applied in order to simplify methods used in previous examples when finding confidence intervals & upper bounds on θ^* .

The sequence of estimators $\hat{\theta}_n := \frac{1}{n} \sum_{i=1}^n X_i$ is consistent by the *Weak Law of Large Numbers* and the conditions for asymptotic normality hold $\forall \theta \in \Theta$. Hence

$$\sqrt{nI(\theta^*)}(\hat{\theta}_n - \theta^*) \rightarrow_{\mathcal{D}(\cdot; \theta^*)} Z \sim \text{Normal}(0, 1)$$

We can compute the *Fisher Information* $\forall \theta \in \Theta$. We have

$$\begin{aligned} \ell'(\theta(x)) &= \frac{x}{\theta} - \frac{1-x}{1-\theta} \\ \text{and } \ell''(\theta; x) &= -\frac{x}{\theta^2} - \frac{1-x}{(1-\theta)^2} \\ \implies I(\theta) &= \frac{1}{\theta} + \frac{1}{1-\theta} \\ &= \frac{1}{\theta(1-\theta)} \end{aligned}$$

In practice θ^* is unknown so we replace $I(\theta^*)$ with $I(\hat{\theta}_n)$ to give the asymptotically exact confidence interval, $[L(\mathbf{X}), U(\mathbf{X})]$ where

$$L(\mathbf{X}) = \hat{\theta}_n - z_{\alpha/2} \sqrt{\frac{\hat{\theta}_n(1-\hat{\theta}_n)}{n}} \text{ and } U(\mathbf{X}) = \hat{\theta}_n + z_{\alpha/2} \sqrt{\frac{\hat{\theta}_n(1-\hat{\theta}_n)}{n}}$$

If we did not know how to compute $I(\theta)$ we could instead compute

$$\begin{aligned} J_n(\hat{\theta}_n) &= -\frac{1}{n} \sum_{i=1}^n \ell''(\hat{\theta}_n; X_i) \\ &= -\frac{1}{n} \sum_{i=1}^n \left\{ -\frac{X_i}{\hat{\theta}_n^2} - \frac{1-X_i}{(1-\hat{\theta}_n)^2} \right\} \\ &= \frac{1}{\hat{\theta}_n^2} \left(\frac{1}{n} \sum_{i=1}^n X_i \right) + \frac{1}{(1-\hat{\theta}_n)^2} \left(1 - \frac{1}{n} \sum_{i=1}^n X_i \right) \\ &= \frac{\hat{\theta}_n}{\hat{\theta}_n^2} + \frac{1-\hat{\theta}_n}{(1-\hat{\theta}_n)^2} \\ &= \frac{1}{\hat{\theta}_n(1-\hat{\theta}_n)} \end{aligned}$$

In this case $J_n(\hat{\theta}_n) = I(\hat{\theta}_n)$, this is not always true.

Definition 11.1 - *Observed Fisher Information*

Let $\mathbf{X} \stackrel{\text{iid}}{\sim} f(\cdot; \theta^*)$ be a vector of n random variables.

The *Observed Fisher Information* at θ is

$$nJ_n(\theta) = -\ell''(\theta; \mathbf{X}) = -\sum_{i=1}^n \ell''(\theta; X_i)$$

N.B. $\mathbb{E}(J_n(\theta^*); \theta^*) = I(\theta^*)$ and that it differs from the *Fisher Information* (under the *Fisher Information Regularity Conditions* by not being an expectation.

0 Appendix

Definition 0.1 - Gradient

$$\nabla f(\boldsymbol{\theta}; \mathbf{x}) := \left(\frac{\partial f(\boldsymbol{\theta}; \mathbf{x})}{\partial \theta_1}, \dots, \frac{\partial f(\boldsymbol{\theta}; \mathbf{x})}{\partial \theta_n} \right)$$

Definition 0.2 - Hessian

$$\nabla^2 f(\boldsymbol{\theta}; \mathbf{x}) := \begin{pmatrix} \frac{\partial^2 f(\boldsymbol{\theta}; \mathbf{x})}{\partial \theta_1^2} & \frac{\partial^2 f(\boldsymbol{\theta}; \mathbf{x})}{\partial \theta_1 \partial \theta_2} & \cdots & \frac{\partial^2 f(\boldsymbol{\theta}; \mathbf{x})}{\partial \theta_1 \partial \theta_n} \\ \frac{\partial^2 f(\boldsymbol{\theta}; \mathbf{x})}{\partial \theta_1 \partial \theta_2} & \frac{\partial^2 f(\boldsymbol{\theta}; \mathbf{x})}{\partial \theta_2^2} & \cdots & \frac{\partial^2 f(\boldsymbol{\theta}; \mathbf{x})}{\partial \theta_2 \partial \theta_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 f(\boldsymbol{\theta}; \mathbf{x})}{\partial \theta_1 \partial \theta_n} & \frac{\partial^2 f(\boldsymbol{\theta}; \mathbf{x})}{\partial \theta_2 \partial \theta_n} & \cdots & \frac{\partial^2 f(\boldsymbol{\theta}; \mathbf{x})}{\partial \theta_n^2} \end{pmatrix}$$

0.1 Notation

Notation	Denotes
$Z_n \rightarrow_{\mathbb{P}} Z$	$\{Z_n\}_{n \in \mathbb{N}}$ converges in <i>Probability</i> to random variable Z .
$Z_n \rightarrow_{\mathcal{D}} Z$	$\{Z_n\}_{n \in \mathbb{N}}$ converges in <i>Distribution</i> to random variable Z .
$Z_n \rightarrow_{qm} Z$	$\{Z_n\}_{n \in \mathbb{N}}$ converges in <i>Quadratic Mean</i> to random variable Z .
$\theta \in \Theta \subseteq \mathbb{R}^{d_\theta}$	Scalar or vector parameter characterising a probability distribution
$\hat{\theta}$	Estimation for the value of the parameter θ
θ^*	True value of the parameter θ
\mathbb{P}	Probability measure $\mathbb{P} : \mathcal{F} \rightarrow [0, 1]$
Ω	Sample space
X	Scalar random variable
\mathcal{F}	Sigma field (Set of events)
χ	Support of rv X . A set χ is definitely in it <i>i.e.</i> $\mathbb{P}(X \in \chi; \theta) = 1$
\mathbf{X}	Vector consisting of scalar random variables

0.2 R

Command	Result
<code>hist(a)</code>	Plots a histogram of the values in array a
<code>mean(a)</code>	Returns the mean value of array a
<code>rbinom(s, n, p)</code>	Samples n of $Bi(n, p)$ random variables
<code>rep(v, n)</code>	Produces an array of size n where each entry has value v
<code>x ← v</code>	Maps value v to variable x

0.3 Probability Distributions

Definition 0.3 - Binomial Distribution

Let X be a discrete random variable modelled by a *Binomial Distribution* with n events and rate of success p .

$$\begin{aligned} p_X(k) &= \binom{n}{k} p^k (1-p)^{n-k} \\ \mathbb{E}(X) &= np \quad \& \quad \text{Var}(X) = np(1-p) \end{aligned}$$

Definition 0.4 - Gamma Distribution

Let T be a continuous random variable modelled by a *Gamma Distribution* with shape parameter

α & scale parameter λ . Then

$$\begin{aligned} f_T(x) &= \frac{\lambda^\alpha x^{\alpha-1} e^{-\lambda x}}{\Gamma(\alpha)} \quad \text{for } x > 0 \\ \mathbb{E}(T) &= \frac{\alpha}{\lambda} \quad \& \quad \text{Var}(T) = \frac{\alpha}{\lambda^2} \end{aligned}$$

N.B. $\alpha, \lambda > 0$.

Definition 0.5 - Exponential Distribution

Let T be a continuous random variable modelled by a *Exponential Distribution* with parameter λ . Then

$$\begin{aligned} f_T(t) &= \mathbf{1}\{t \geq 0\} \cdot \lambda e^{-\lambda t} \\ F_T(t) &= \mathbf{1}\{t \geq 0\} \cdot (1 - e^{-\lambda t}) \\ \mathbb{E}(X) &= \frac{1}{\lambda} \quad \& \quad \text{Var}(X) = \frac{1}{\lambda^2} \end{aligned}$$

N.B. Exponential Distribution is used to model the wait time between decays of a radioactive source.

Definition 0.6 - Normal Distribution

Let X be a continuous random variable modelled by a *Normal Distribution* with mean μ & variance σ^2 .

Then

$$\begin{aligned} f_X(x) &= \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}} \\ F_X(x) &= \frac{1}{\sqrt{2\pi\sigma^2}} \int_{-\infty}^x e^{-\frac{(y-\mu)^2}{2\sigma^2}} dy \\ M_X(\theta) &= e^{\mu\theta + \sigma^2\theta^2(1/2)} \\ \mathbb{E}(X) &= \mu \quad \& \quad \text{Var}(X) = \sigma^2 \end{aligned}$$

Definition 0.7 - Poisson Distribution

Let X be a discrete random variable modelled by a *Poisson Distribution* with parameter λ . Then

$$\begin{aligned} p_X(k) &= \frac{e^{-\lambda} \lambda^k}{k!} \quad \text{For } k \in \mathbb{N}_0 \\ \mathbb{E}(X) &= \lambda \quad \& \quad \text{Var}(X) = \lambda \end{aligned}$$

N.B. Poisson Distribution is used to model the number of radioactive decays in a time period.