Arthritis-friendly design principles

"arthritis people friendly design" *Generated by LeapSpace*

Arthritis-Friendly Design at a Glance

Key Findings Table

Key finding	Population/context	Evidence
The most consistent design principle is reducing force and joint strain: lower grip demand, less awkward wrist/finger positioning, lighter tools, slip resistance, and simplified interactions improve arthritis-friendliness across physical and digital products [1] [2] [3] [4] [5] [6].	People with rheumatoid arthritis (RA), hand osteoarthritis (OA), workers with arthritis, digital users with dexterity limits	[1] [2] [3] [4] [5] [6]
Adaptive devices can improve pain, hand function, occupational performance, and independence, especially for eating, cooking, toileting, bathing, and hand-related daily tasks [7] [8] [9] [10].	RA and hand OA in daily living	[7] [8] [9] [10]
Personalization matters more than one-size-fits-all design: fit, acceptability, and individualized assessment predict compliance and satisfaction [11] [12] [13] [14] [15].	Assistive devices, workplace accommodation, personalized aids	[11] [12] [13] [14] [15]
The evidence base is stronger for multimodal programs combining education, joint protection, exercise/orthoses, and adaptive tools than for isolated product features [9] [16] [17] [18].	RA and OA rehabilitation/occupational therapy	[9] [16] [17] [18]
Reviews and trials conflict: some studies report functional and pain benefits, but high-level syntheses often find only small or clinically unimportant average effects, and some products such as arthritis gloves show no meaningful benefit [19] [20] [21].	Joint-protection and assistive-product evidence base	[19] [20] [21]

Key finding	Population/context	Evidence
For digital tools, **large adjustable text, high contrast, large spaced tap targets, fewer screens, and alternatives to writing** are preferred; navigation and data entry remain major pain points [5] [6] [22] [23] [24].	Arthritis-related apps and touchscreen workflows	[5] [6] [22] [23] [24]
Co-design is increasingly the dominant development approach: patients, clinicians, and developers shape needs assessment, prototyping, and testing, usually improving acceptability and usability, though long-term effectiveness is less established [25] [26] [27] [28] [29].	Arthritis services, apps, education tools, toolkits	[25] [26] [27] [28] [29]

Direct Answer

People-friendly design for arthritis should mainly do five things:

1. **Reduce force demand** with built-up or easy-grip handles, lighter tools, anti-slip surfaces, and designs that avoid tight grasping or twisting [1] [2] [3] [4].
2. **Protect joints** by minimizing awkward wrist/finger angles, repetition, compression, and heavy lifting [20] [30] [31] [32].
3. **Match the person and task**, because usability and continued use depend heavily on individualized fit and acceptability rather than generic prescription [11] [12] [13] [14].
4. **Simplify digital interaction** with large text, high contrast, larger buttons, fewer steps, and alternatives to handwriting or complex navigation [5] [6] [22] [24].
5. **Use co-design** with people who have arthritis and clinicians to improve relevance, usability, and acceptance [25] [27] [28] [29].

Practical implication: the best-supported approach is not a single “arthritis product,” but a **design system** spanning home tools, work accommodations, and digital interfaces that consistently lowers force, lowers complexity, and allows personalization [1] [5] [25] [32].

Confidence level: High - the cross-cutting principles recur across multiple themes and study types, even though feature-level product trials are limited [1] [2] [4] [5] [25].

Study Scope

This report synthesizes evidence on arthritis-friendly design across five domains: ergonomic physical products, adaptive daily-living tools, workplace accommodations, accessible digital interfaces, and participatory co-design methods [25] [30] [33] [34]. The underlying studies include randomized trials, quasi-experimental studies, systematic reviews, case reports, surveys, usability studies, and implementation studies, largely focused on RA and hand or knee OA [16] [26] [32] [35] [36].

Assumptions & Limitations

- The evidence base contains **few controlled head-to-head product trials** isolating exact design variables such as handle diameter, angle, texture, or weight against standard products in arthritis populations [4] [37].
- Many positive findings come from **bundled interventions** that combine education, behavior change, therapy, orthoses, and assistive devices, making feature-specific attribution difficult [9] [16] [17].
- Long-term adherence evidence specifically for **older adults or people with severe arthritis-related impairment** remains limited in this set [38] [39] [40].
- Outcome measurement is fragmented: broad clinical measures are common, but direct design-usability metrics such as grip effort, comfort, and feature-level burden are less standardized [41] [42] [43].

Suggested Further Research

- Run controlled trials comparing specific tool/interface features against standard designs in arthritis populations [4] [44].
- Develop arthritis-specific usability measures covering pain during task, grip effort, joint-angle burden, comfort, and sustained use [41] [43].
- Study long-term adherence and abandonment of adaptive tools and digital systems in older adults and people with severe hand impairment [38] [39] [40].
- Separate the effects of education/training from the effects of design itself in multimodal programs [16] [17].
- Improve representativeness in co-design samples to reduce local or selective bias [45].

Full Report Body

Foundational Context

“People-friendly design” for arthritis is best understood as design that reduces the mismatch between a task’s physical or cognitive demands and the user’s pain, stiffness, weakness, fatigue, or limited dexterity [30] [33] [34]. In the retrieved evidence, this usually appears through three linked frameworks:

- **Joint protection**: distribute load, reduce effort, avoid prolonged or awkward positions, pace tasks, and modify tools or environments to lower strain [20] [30] [46].
- **Assistive enablement**: use devices or altered methods to preserve independence in self-care, cooking, work, and communication [33] [47] [48] [49].
- **Accessible interaction**: reduce motor precision demands in digital systems through larger, simpler, and more forgiving interfaces [5] [6] [22].

Taken together, these findings suggest that arthritis-friendly design is less about a specific product category than about a repeatable set of design rules: lower force, safer posture, fewer steps, and better fit to fluctuating function [1] [5] [32].

Confidence level: High - these core concepts are repeated across reviews, intervention studies, and usability work [22] [30] [33].

Ergonomic Product Design and Joint Protection

Low-force and joint-protective design features

The most consistently supported ergonomic principle is reducing force and joint-angle burden during hand use [1] [2] [3]. In RA, people preferred **narrower, slightly textured saucepan handles**, suggesting that tactile control and efficient grasp geometry matter [1]. Built-up utensil handles reduced required finger-joint flexion relative to standard handles, supporting designs that lower extreme small-joint positioning during eating tasks [2]. One experiment found a **42.25-mm grasp distance** minimized muscle load, reinforcing that geometry can materially alter strain [3].

Broader ergonomic guidance in the reviewed material recommends **lighter weight, easier grasping, convenient length, slip-resistant surfaces, rounded or curved edges, and reduced tissue compression** for kitchen tools [4]. These recommendations align with studies that measure output force, tool weight, handle properties, posture, muscular load, local grip pressure, and wrist angles as relevant design outcomes [42] [43].

A useful caution is that not all commonly prescribed products help: arthritis/compression gloves did not improve pain, function, or stiffness compared with placebo and were not cost-effective [21].

Confidence level: High - the directional principle is well supported, though exact optimal dimensions are not broadly established across products [1] [2] [3] [4].

Comparisons with standard household and kitchen tools

Evidence directly comparing arthritis-adapted tools with standard tools is limited but practically suggestive [4] [44]. In RA, specially designed tools such as **breadsaws, potato peelers, and scissors** reduced pain compared with standard tools [7] [8]. Reviews also report assistive devices being especially effective for **eating, cooking, and toileting** difficulties [8].

For eating and cooking, the literature describes **jar openers, adapted knives, electric can openers, ergonomic peelers, built-up and angled cutlery, and ergonomic handled cutlery** as improving comfort, independence, control, and maneuverability while reducing wrist strain or shock from weak or painful grasping [44] [50] [51]. However, much of this evidence comes from surveys, qualitative studies, and reviews rather than robust randomized product-comparison trials [44].

This creates a tension in the literature: clinically, adapted tools often seem useful; methodologically, the feature-level evidence remains thin [7] [44].

Confidence level: Medium - practical benefit is plausible and repeatedly reported, but controlled comparative evidence is sparse [7] [8] [44].

Joint-protection education plus environmental modification

Evidence is stronger when design changes are embedded in broader occupational therapy or joint-protection programs [9] [16] [17] [18]. In moderate-severe RA, an educational-behavioral joint-protection program that included adapting the surrounding environment reduced pain and disability and improved several AIMS2 domains at about 8 months versus control [16]. In CMC1 OA, multimodal occupational therapy combining education, exercises, orthoses, and assistive devices improved pain, grip strength, MAP-Hand, and QuickDASH at 3 to 4 months [17]. In hand OA, assistive devices outperformed leaflet-based joint-protection information over 30 to 90 days for hand function, occupational performance, and pain [9]. Standard care incorporating orthoses, joint-protection education, and adaptive equipment also improved activity pain and patient-specific function in thumb CMC OA [18].

But high-level syntheses are more cautious. One overview found that high-quality reviews showed no clinically important benefit for pain, hand function, or grip strength, whereas lower-quality reviews reported improvements across pain, function, fatigue, depression, self-efficacy, and symptoms [20]. A meta-analysis found only small or clinically unimportant average effects overall, with no short-term pain benefit and no long-term grip-strength benefit [19].

The conflict is therefore explicit: **applied programs often look beneficial, but pooled average effects are modest** [16] [17] [19] [20].

Confidence level: Medium-High - benefits are supported in several intervention studies, but effect size certainty is weakened by contradictory syntheses and multimodal designs [16] [17] [19].

Workplace ergonomics and accommodations

At work, arthritis-friendly design extends beyond objects to task structure and accommodation [15] [32]. A randomized trial showed that a workplace ergonomic intervention for workers with RA/OA reduced arthritis-related work difficulties over two years and improved physical functioning and pain compared with written education alone [32]. Reported work risks commonly included **heavy physical work, repetition, awkward posture, and lifting**, especially for hand-based tasks [31] [52].

Individualized ergonomic assessment and vocational counseling generated recommendations with high implementation rates in arthritis-specific work accommodation programs [15]. For measurement, the **WALS** and **RA WIS** were rated among the best tools for at-work productivity loss and were endorsed for worker productivity measurement [53] [54].

Evidence from broader office ergonomics suggests moderate support for arm/forearm supports, stretching programs, mouse-use feedback, and strong evidence for workplace-based resistance training, while standard workstation adjustment alone showed no clear benefit [55]. Evidence for alternative mouse designs alone was inconsistent [56].

Confidence level: High - arthritis-specific work difficulty reduction through ergonomic intervention is directly supported, especially when accommodations are individualized [15] [32].

Adaptive Tools for Daily Living

Devices for self-care and personal tasks

Assistive devices for arthritis are mainly used to support self-care, hand-related activities, and independence [33] [47] [48] [49]. In hand OA, assistive devices for daily activities improved occupational performance and hand function relative to a guideline leaflet, with pain improvement at 30 and 90 days [9]. In RA, a long-handled bath sponge had an 86% usage rate after prescription, while non-users were less favorable toward the device and perceived greater personal control over pain/disability, indicating that acceptance shapes adherence [13]. Across RA assistive-device users, satisfaction was generally high and abandonment limited, though sustained use varied by device type [14].

Dressing aids are associated with reduced pain and better function in older adults, and dressing difficulty tracks worse functional and mental status [10]. For a more narrowly studied example, an eye-drop device reduced both observed and reported difficulty compared with a standard bottle in RA, though overall evidence in this category is very limited [57] [58].

Confidence level: Medium-High - several device categories show benefit, but the number of direct trials is small [9] [13] [57].

Eating and cooking aids

Cooking and eating aids are among the most consistently useful categories in the evidence [8] [44]. Built-up handled utensils reduce finger flexion demands [2]. Adapted kitchen tools such as jar openers, peelers, ergonomic cutlery, and adapted knives are reported to improve independence and reduce strain [44] [50] [51]. In RA, specially designed kitchen-type tools reduced pain compared with standard versions [7] [8].

Outcome measurement in this area typically relies on patient-reported function scales such as AIMS-2 hand/finger function, DHI, HAQ-related hand items, openability ratings for food packaging, plus pain, grip strength, and task performance [59] [60] [61] [62] [63] [64].

The implication is practical: arthritis-friendly kitchen design should minimize twisting, sustained pinch, and shock transfer while improving leverage and grasp security [2] [50].

Confidence level: Medium - directionally consistent, but many studies use user reports rather than tightly controlled comparative testing [7] [44].

Client-centered and 3D-printed devices

Personalized devices are a promising extension of arthritis-friendly design [35]. A client-centered occupational therapy case report described 3D-printed devices for opening beverage bottles, locking/unlocking doors, and writing, with user input guiding device choice and modification [35]. After fitting, the participant reported decreased pain, improved occupational performance, and satisfaction [35]. Reviews also suggest 3D printing can address fit and comfort problems seen with traditional splints and orthoses, including pain, swelling, and pressure issues [65].

Taken together, these findings suggest personalization is especially valuable for tasks that are highly individual or poorly served by mass-market products [35] [65].

Confidence level: Low-Medium - promising but currently supported mainly by case-level and review evidence [35] [65].

Accessible Digital Interfaces

Usability features for arthritis-friendly apps and forms

Digital accessibility for arthritis remains an emerging field, but several interface preferences recur consistently [34] [66] [67] [68]. A digital app for inflammatory arthritis was described as highly usable partly because it used **simple minimal design, high-contrast colors, and adjustable large fonts** [22]. Across studies relevant to dexterity limitations, **small fonts, low contrast, long text, and irrelevant icons** were problematic, while **large touch targets, enough spacing, fewer screens, and voice/keyboard alternatives** were preferred [5] [6] [69]. In a rheumatology mobile workflow tool, users liked larger fonts and tapping instead of writing or circling, but still reported problems with navigation, data entry, pain assessment, documentation, and instructions [23]. A hand-OA digital joint-protection program also identified navigation as a major barrier [24].

One study found mobile/touchscreen interfaces could be preferred over paper for clinic data capture, with high ease-of-use ratings and no technical problems [66]. Another hand-exercise app for RA reported high usability and satisfaction after user-centered development [67].

Confidence level: High - the direction of preferred interface features is consistent across multiple usability studies [5] [6] [22] [23] [24].

Alternative input methods and device tradeoffs

Alternative input methods can help, but tradeoffs are common [70] [71] [72]. In RA, reduced hand function was only weakly linked to typing speed, whereas mouse speed was slower than nonimpaired norms; touch-typing training mattered more than impairment for typing speed [70]. Across accessibility studies, personalized keyboards, chorded on-screen input, and some alternative devices improved usability or performance, but not uniformly [73] [74] [75]. Eye- and head-based systems showed classic speed-error tradeoffs: some options were faster but more error-prone, while others reduced errors at the cost of speed [71] [76]. Reviews found that some alternative pointing devices reduced biomechanical loading and many alternative keyboards improved musculoskeletal outcomes, but touchscreen/gestural controls were not yet suitable replacements in general [72].

For physical controls, one study found arthritis participants applied less torque and had more difficulty with larger knobs, implying that “bigger” is not universally better for rotary interfaces [77].

Confidence level: Medium - useful trends exist, but evidence is heterogeneous and often not arthritis-specific alone [72] [77].

Co-designed digital self-management tools

User-centered digital development is now common in arthritis tools [27] [36] [67] [78]. The MijnReuma Reade app explicitly reported two pre-deployment pilot/usability rounds involving patients, rheumatologists, and IT experts [36]. The Remote Monitoring of RA app was co-designed with patients, clinicians, and researchers via interviews and focus groups, though the exact number of iterations was not reported [78]. A web-based RA self-management program also used co-design with patients and professionals [79]. The MyRA app used focus groups, co-creation sessions, heuristic review, and SUS evaluation, achieving a mean SUS of 84.4 [27]. A Turkish hand-exercise app used focus groups with therapists and people with RA, then prototype testing and revision, followed by 4 to 6 weeks of use with high usability and satisfaction [67].

The evidence indicates that co-design improves pre-deployment usability and acceptability, but stronger evidence on downstream pain, function, and quality-of-life outcomes is still limited [25] [78].

Confidence level: Medium-High - development methods are well described, but effectiveness beyond usability remains less certain [25] [27] [36].

Participatory Co-Design and Evaluation Outcomes

Co-design methods and stakeholder roles

A systematic review found co-designed inflammatory-arthritis interventions commonly used **needs assessment, ideation, prototyping, and pilot testing** [25]. Patient participation was often active and early: defining needs, shaping research questions, reviewing materials, testing prototypes, and advising on usability and communication [29] [80] [81] [82]. Clinicians contributed workflow, content, feasibility, and service-specific knowledge [29] [81] [82].

The clearest service-redesign framework in the set is the participatory health research approach used in the IMPACT project, where a steering committee including academics, people with arthritis, advocacy members, physiotherapists, GPs, and an orthopaedic surgeon shaped outcomes, recruitment, analysis, and dissemination [29].

Barriers include representativeness, literacy, technical constraints, local specificity, and communication problems [45] [83] [84]. Facilitators include iterative feedback, flexible participation formats, and balancing multiple user perspectives [25] [85] [86].

Confidence level: High - methods and stakeholder patterns are repeated across multiple studies and reviews [25] [29] [45].

Examples of co-designed arthritis interventions

Examples span toolkits, monitoring apps, decision aids, education tools, vocational interventions, and rehabilitation programs [26] [27] [28] [87]. The **My Knee** OA toolkit used systematic review, web-content appraisal, concept mapping, expert review, and three workshops to refine accuracy and usability [26]. The **ANSWER** methotrexate decision aid used animated patient stories, interactive preference questions, think-aloud testing, SUS, observation, and content analysis; users found it easy to use, with a mean SUS of 81.2 [28]. Multimedia education tools for knee OA, osteoporosis, and RA produced significant knowledge gains and high acceptability/comprehension in pilot testing [88]. A PT-led vocational intervention for inflammatory arthritis had positive feasibility findings but required PT training for delivery [87]. Co-design workshops with OA consumers showed strong engagement and changed the research focus, supporting equal-partnership design feasibility [89].

These projects show that co-design is not only feasible but productive in shaping content, format, and task priorities [90] [91].

Confidence level: High - there are multiple well-described examples with convergent feasibility and acceptability findings [26] [28] [87].

Outcome measures for design evaluation

Outcome measurement is a strength and weakness of the field [41] [92] [93]. Strong measures exist for pain, disability, quality of life, and work function:

- **Pain VAS**: repeatedly used and sensitive to change [92] [94] [95].
- **HAQ**: core disability measure that explicitly accounts for aids/devices [93] [96].
- **EQ-5D**: widely used for health-related quality of life [94] [95] [97] [98].
- **MAP-Hand**: task-specific hand activity performance [41].
- **WALS / RA WIS**: work limitations and instability [53] [54].
- **PIADS and QUEST/QUEST 2**: device impact and satisfaction measures used in assistive-technology evaluation [99].

However, many studies still rely on broad clinical endpoints rather than direct design-usability outcomes such as grip effort, completion burden, comfort, or feature-level interaction difficulty [41] [42] [43].

Confidence level: High - measurement fragmentation is directly supported across several outcome-focused sources [41] [43] [53].

Sustained use, adherence, and implementation barriers

For physical devices, sustained use can be fairly good when devices are acceptable and well-matched. In RA, 91% of devices provided after joint-protection education were still in use in one study [100]. A bath sponge had 86% usage after prescription, with non-use linked to less favorable attitudes [13]. Across assistive-device users, abandonment was generally limited but varied by category [14].

For digital systems, the arthritis-specific long-term evidence is thin, but adjacent implementation studies in older adults with functional limitations consistently identify **training/support, usability, maintenance, and fit with daily life** as key to adherence [38] [39] [40] [101] [102] [103]. This pattern likely applies to arthritis as well, but that extension is an inference rather than a directly established arthritis-specific finding.

Inferred: Digital arthritis tools are likely to have better sustained use when onboarding is simple, support is available, and interaction burden remains low over time [23] [24] [101].
Confidence level: Low - plausible cross-source synthesis, but not directly tested in arthritis-specific long-term studies [24] [101].

Cross-Source Synthesis: What Arthritis-Friendly Design Looks Like in Practice

Across the evidence base, arthritis-friendly design consistently converges on four practical principles:

1. **Low-force activation**: reduce gripping, pinching, twisting, and sustained load [1] [3] [4].
2. **Joint-safe geometry**: reduce extreme finger flexion and awkward wrist postures [2] [43].
3. **Interaction simplicity**: reduce steps, screens, text burden, and navigation complexity [5] [22] [24].

4. **Customization**: adapt the product, workspace, or interface to the user's actual impairments and priorities [11] [12] [15] [35].

Taken together, these findings imply that the best arthritis-friendly solutions are usually **systems**, not isolated objects: for example, a kitchen task may improve most when the person has an adapted handle, altered workflow, pacing strategy, and possibly digital prompting or instructional support [8] [30] [68].

Confidence level: High - the synthesis directly reflects repeated patterns across physical, workplace, and digital evidence [1] [5] [11] [30].

Practical Design Guidance Derived from the Evidence

Physical product design

- Use built-up or efficient-diameter handles and avoid requiring tight grasp [2] [3].
- Prefer light weight and slip-resistant surfaces [4].
- Reduce twisting and pinch-heavy interactions where possible [7] [50].
- Favor rounded edges and lower compression surfaces [4].

Assistive device provision

- Assess the specific task and user priorities before prescribing [11] [12].
- Expect better use when the device feels acceptable, relevant, and easy to integrate [13] [14].
- Reassess after fitting because continued use varies by device category [14].

Workplace design

- Target repetition, posture, lifting, and task layout rather than only workstation furniture [31] [32] [55].
- Use individualized ergonomic recommendations and accommodation planning [15].

Digital interface design

- Large adjustable text, high contrast, clear spacing, and large targets are preferred [5] [22].
- Minimize screens and navigation depth [6] [24].
- Avoid handwriting-heavy or precision-heavy inputs when alternatives exist [23] [69].
- Consider alternative input options when hand pain or dexterity fluctuates [72] [75].

Confidence level: High - each recommendation is directly supported, although exact parameter values are rarely standardized [4] [5].

Conclusions

The clearest answer from the evidence is that arthritis-friendly design should be built around **force reduction, joint protection, simplified interaction, and individualized fit** [1] [5] [11] [32]. Adaptive tools and ergonomic interventions often improve pain, function, and independence in daily life and work, but the field relies heavily on multimodal programs rather than clean product-feature comparisons [9] [16] [17]. Digital arthritis design is advancing through user-centered methods and has identifiable usability rules, yet navigation and long-term effectiveness remain underdeveloped [23] [24] [25]. Co-design has become a practical best practice for relevance and acceptability, though proof of sustained clinical impact is still incomplete [25] [27] [29].

Overall, the evidence supports a move from generic accessibility toward **arthritis-specific, low-strain, personalized design systems** across products, workplaces, and digital tools [1] [22] [25] [35].

Overall confidence: Medium-High - the broad design principles are strongly supported across domains, but the evidence is weakened by limited head-to-head product trials, heterogeneous interventions, and fragmented usability measurement [4] [19] [41].

References

1. Torrens, G.E., Hann, J., Webley, M., (...), Sutherland, I.A. (2000) Hand performance assessment of ten people with Rheumatoid Arthritis when using a range of specified saucepans. *Disability and Rehabilitation*. https://doi.org/10.1080/096382800296971
2. Jordon, M., Levine, D., Richards, J., (...), Brit, M. (2026) Biomechanical assessment of grasping postures in individuals with rheumatoid arthritis when holding adaptive silverware. *Frontiers in Rehabilitation Sciences*. https://doi.org/10.3389/fresc.2026.1770139
3. Liu, C.-H., Yip, K.-S., Fan, S.-C. (2017) Optimal grasp distance and muscle loads for people with rheumatoid arthritis using carpometacarpal and metacarpophalangeal immobilization orthoses. *American Journal of Occupational Therapy*. https://doi.org/10.5014/ajot.2017.017681
4. Chen, H., Anton, N., Holden, R.J., Yu, D. (2024) Kitchen ergonomics in health and healthcare: A rapid scoping review. *Human Factors in Healthcare*. https://doi.org/10.1016/j.hfh.2024.100069
5. Kim, H., Xie, B. (2017) Health literacy in the eHealth era: A systematic review of the literature. *Patient Education and Counseling*. https://doi.org/10.1016/j.pec.2017.01.015
6. Radcliffe, Erin, Lippincott, Ben, Anderson, Raeda, Jones, Mike (2021) A Pilot Evaluation of mHealth App Accessibility for Three Top-Rated Weight Management Apps by People with Disabilities. *International Journal of Environmental Research and Public Health*. https://doi.org/10.3390/ijerph18073669
7. Nordenskiöld, U. (1997) Daily activities in women with rheumatoid arthritis. Aspects of patient education, assistive devices and methods for disability and impairment assessment. *Scandinavian Journal of Rehabilitation Medicine, Supplement*. https://www.scopus.com/pages/publications/0030671421

8. Nordenskiöld, U. (1997) Daily activities in women with rheumatoid arthritis. Aspects of patient education, assistive devices and methods for disability and impairment assessment.. *Scandinavian journal of rehabilitation medicine. Supplement*. https://www.scopus.com/pages/publications/0031307093
9. Amaral, D.S., Branco Pinto Duarte, A.L.B.P., Barros, S.S., (...), Marques, C.D.L. (2018) Assistive devices: an effective strategy in non-pharmacological treatment for hand osteoarthritis—randomized clinical trial. *Rheumatology International*. https://doi.org/10.1007/s00296-017-3892-1
10. Mann, W.C., Kimble, C., Justiss, M.D., (...), Wu, S.S. (2005) Problems with dressing in the frail elderly. *American Journal of Occupational Therapy*. https://doi.org/10.5014/ajot.59.4.398
11. Davalbhakta, S., Oswal, A., Phatak, S. (2020) Assistive devices: Regaining mobility in myositis. *Indian Journal of Rheumatology*. https://doi.org/10.4103/injr.injr_90_20
12. Ghasemzadeh, R., Kamali, M. (2010) Assistive technology: Use and service delivery. *Iranian Rehabilitation Journal*. https://www.scopus.com/pages/publications/84905259540
13. Rogers, J.C., Holm, M.B., Perkins, L. (2002) Trajectory of assistive device usage and user and non-user characteristics: Long-handled bath sponge. *Arthritis Care and Research*. https://doi.org/10.1002/art.10788
14. de Boer, I.G., Peeters, A.J., Ronday, H.K., (...), Vliet-Vlieland, T.P.M. (2009) Assistive devices: Usage in patients with rheumatoid arthritis. *Clinical Rheumatology*. https://doi.org/10.1007/s10067-008-0989-7
15. Lacaille, D., White, M.A., Rogers, P.A., (...), Esdaile, J.M. (2008) A proof-of-concept study of the "Employment and Arthritis: Making it Work" program. *Arthritis Care and Research*. https://doi.org/10.1002/art.24197
16. Masiero, S., Boniolo, A., Wassermann, L., (...), Punzi, L. (2007) Effects of an educational-behavioral joint protection program on people with moderate to severe rheumatoid arthritis: A randomized controlled trial. *Clinical Rheumatology*. https://doi.org/10.1007/s10067-007-0615-0
17. Tveter, A.T., Osterås, N., Nossum, R., (...), Kjeken, I. (2022) Short-Term Effects of Occupational Therapy on Hand Function and Pain in Patients With Carpometacarpal Osteoarthritis: Secondary Analyses From a Randomized Controlled Trial. *Arthritis Care and Research*. https://doi.org/10.1002/acr.24543
18. Pisano, K., Wolfe, T., Lubahn, J., Cooney, T. (2023) Effect of a stabilization exercise program versus standard treatment for thumb carpometacarpal osteoarthritis: A randomized trial. *Journal of Hand Therapy*. https://doi.org/10.1016/j.jht.2022.03.009
19. Bobos, P., Nazari, G., Szekeres, M., (...), MacDermid, J.C. (2019) The effectiveness of joint-protection programs on pain, hand function, and grip strength levels in patients with hand arthritis: A systematic review and meta-analysis. *Journal of Hand Therapy*. https://doi.org/10.1016/j.jht.2018.09.012

20. Bobos, P., MacDermid, J.C., Nazari, G., (...), Grewal, R. (2021) Joint protection programmes for people with osteoarthritis and rheumatoid arthritis of the hand: An overview of systematic reviews. *Physiotherapy Canada*. https://doi.org/10.3138/ptc-2019-0037

21. Hammond, A., Prior, Y., Cotterill, S., (...), Firth, J. (2021) Clinical and cost effectiveness of arthritis gloves in rheumatoid arthritis (A-GLOVES): randomised controlled trial with economic analysis. *BMC Musculoskeletal Disorders*. https://doi.org/10.1186/s12891-020-03917-8

22. Mattli, Renato, Weber, Manuel, Raab, Anja Maria, (...), Schmitt, Kai-Uwe (2025) Digital Multidomain Lifestyle Intervention for Community-Dwelling Older Adults: A Mixed Methods Evaluation. *International Journal of Public Health*. https://doi.org/10.3389/ijph.2025.1608014

23. Yen, P.-Y., Lara, B., Lopetegui, M., (...), Curtis, J.R. (2016) Usability and workflow evaluation of "RhEumAtic disease activity" (READY): A mobile application for rheumatology patients and providers. *Applied Clinical Informatics*. https://doi.org/10.4338/ACI-2016-03-RA-0036

24. Pouliopoulou, Dimitra V., D'Alessandro, Victoria, Billias, Nicole, (...), Bobos, Pavlos (2026) Iterative usability testing of a digital joint protection program for people with hand osteoarthritis. *PLoS ONE*. https://doi.org/10.1371/journal.pone.0342571

25. Lim, H.W., Bundele, A., Phang, K.F., (...), Sumner, J. (2026) Co-designing interventions for patients with inflammatory arthritis: a systematic review. *Systematic Reviews*. https://doi.org/10.1186/s13643-026-03152-7

26. Goff, A.J., De Oliveira Silva, D., Ezzat, A.M., (...), Barton, C.J. (2023) Co-design of the web-based 'My Knee' education and self-management toolkit for people with knee osteoarthritis. *Digital Health*. https://doi.org/10.1177/20552076231163810

27. Roumen, C., Hochstenbach, L.M., den Dungen, P.V., (...), Van Tubergen, A. (2025) Development of an Application for Self-Monitoring to Empower Patients With Rheumatoid Arthritis (MyRA). *Journal of Rheumatology*. https://doi.org/10.3899/jrheum.2024-1071

28. Li, L.C., Adam, P.M., Townsend, A.F., (...), Backman, C.L. (2013) Usability testing of ANSWER: A web-based methotrexate decision aid for patients with rheumatoid arthritis. *BMC Medical Informatics and Decision Making*. https://doi.org/10.1186/1472-6947-13-131

29. Toomey, Clodagh M, Bhardwaj, Avantika, Browne, Jacqui, (...), Kennedy, Norelee (2024) Guideline-based exercise management for hip and knee osteoarthritis: a cross-sectional comparison of healthcare professional and patient beliefs in Ireland. *BMJ Open*. https://doi.org/10.1136/bmjopen-2023-080646

30. Hammond, A. (2009) Joint protection: Enabling change in musculoskeletal conditions multiple sclerosis. *International Handbook of Occupational Therapy Interventions*. https://doi.org/10.1007/978-0-387-75424-6_33

31. da Costa, B.R., Vieira, E.R. (2010) Risk factors for work-related musculoskeletal disorders: A systematic review of recent longitudinal studies. *American Journal of Industrial Medicine*. https://doi.org/10.1002/ajim.20750

32. Baldwin, D., Johnstone, B., Ge, B., (...), Sharp, G. (2012) Randomized prospective study of a work place ergonomic intervention for individuals with rheumatoid arthritis and osteoarthritis. *Arthritis Care and Research*. https://doi.org/10.1002/acr.21699
33. Jeffreson, P. (1997) Arthritis: Using assistive devices to promote independence. *International Journal of Therapy and Rehabilitation*. https://doi.org/10.12968/bjtr.1997.4.10.14355
34. Mollard, E., Michaud, K. (2019) Mobile Apps for Rheumatoid Arthritis: Opportunities and Challenges. *Rheumatic Disease Clinics of North America*. https://doi.org/10.1016/j.rdc.2019.01.011
35. Janson, R., Burkhart, K., Firchau, C., (...), Garabrant, A. (2020) Three-dimensional printed assistive devices for addressing occupational performance issues of the hand: A case report. *Journal of Hand Therapy*. https://doi.org/10.1016/j.jht.2020.03.025
36. Seppen, B.F., Wiegel, J., L'Ami, M.J., (...), Bos, W.H. (2020) Feasibility of self-monitoring rheumatoid arthritis with a smartphone app: Results of two mixed-methods pilot studies. *JMIR Formative Research*. https://doi.org/10.2196/20165
37. Roll, S.C., Yo, S.H. (2022) (Re-)Defining ergonomics in hand therapy: Applications for the management of upper extremity osteoarthritis. *Journal of Hand Therapy*. https://doi.org/10.1016/j.jht.2022.06.006
38. Zhou, L., Luo, T., Li, N., (...), Hou, S. (2026) Digital technology for older adults with functional disabilities at home: a scoping review. *BMC Geriatrics*. https://doi.org/10.1186/s12877-026-07290-4
39. Auger, C., Hajj, N.E., Bier, N., (...), Ahmed, S. (2024) Intervention mapping of a mobility outcomes monitoring system for geriatric patients. *Gerontechnology*. https://doi.org/10.4017/GT.2024.23.S.1066.OPP
40. Picciafoco, L., Galati, S., Colombo, M., (...), Rolandi, E. (2026) Should Digital Technology Skills Be Considered Among the Instrumental Activities of Daily Living of Older Adults? Instruments Available and Open Questions. *Lecture Notes in Computer Science*. https://doi.org/10.1007/978-3-032-30424-7_11
41. Kjeken, Ingvild, Bergsmark, Kjetil, Haugen, Ida K., (...), Slatkowsky-Christensen, Barbara (2021) Task shifting in the care for patients with hand osteoarthritis. Protocol for a randomized controlled non-inferiority trial. *BMC Musculoskeletal Disorders*. https://doi.org/10.1186/s12891-021-04019-9
42. Murphy, P., McGorry, R., Teare, P., Brogmus, G. (2000) Design and performance of a manual task evaluator. *International Journal of Industrial Ergonomics*. https://doi.org/10.1016/S0169-8141(99)00016-5
43. Browne, A., O'Sullivan, L. (2012) A medical hand tool physical interaction evaluation approach for prototype testing using patient care simulators. *Applied Ergonomics*. https://doi.org/10.1016/j.apergo.2011.08.002
44. Hammond, A. (2008) Rehabilitation in musculoskeletal diseases. *Best Practice and Research: Clinical Rheumatology*. https://doi.org/10.1016/j.berh.2008.02.003

45. Aitken, Sarah Joy, James, Sophie, Lawrence, Amy, (...), Naganathan, Vasi (2024) Codesign of health technology interventions to support best-practice perioperative care and surgical waitlist management. *BMJ Health and Care Informatics*. https://doi.org/10.1136/bmjhci-2023-100928

46. Hammond, A. (2015) Joint protection: Enabling change in musculoskeletal conditions. *International Handbook of Occupational Therapy Interventions, Second Edition*. https://doi.org/10.1007/978-3-319-08141-0_42

47. Shipham, I., Pitout, S.J. (2003) Rheumatoid arthritis: hand function, activities of daily living, grip strength and essential assistive devices.. *Curationis*. https://doi.org/10.4102/curationis.v26i3.857

48. Hassel, K., Pinnig, M. (2023) Occupational Therapy and Assistive Devices in Patients with Rheumatoid Disorders of the Hand. *Aktuelle Rheumatologie*. https://doi.org/10.1055/a-1993-4360

49. Mann, W.C., Hurren, D., Tomita, M. (1995) Assistive devices used by home-based elderly persons with arthritis.. *The American journal of occupational therapy. : official publication of the American Occupational Therapy Association*. https://doi.org/10.5014/ajot.49.8.810

50. Marcus, Jacqueline B. (2019) Cooking Aids, Tableware Tips and Dining Advice. *Aging, Nutrition and Taste*. https://doi.org/10.1016/B978-0-12-813527-3.00011-9

51. Bennett, R., Demmers, T.A., Plourde, H., (...), Kakinami, L. (2019) Identifying Barriers of Arthritis-Related Disability on Food Behaviors to Guide Nutrition Interventions. *Journal of Nutrition Education and Behavior*. https://doi.org/10.1016/j.jneb.2019.06.030

52. Gignac, M.A.M., Cao, X., Tang, K., Beaton, D.E. (2011) Examination of arthritis-related work place activity limitations and intermittent disability over four-and-a-half years and its relationship to job modifications and outcomes. *Arthritis Care and Research*. https://doi.org/10.1002/acr.20456

53. Beaton, D.E., Tang, K., Gignac, M.A.M., (...), Bombardier, C. (2010) Reliability, validity, and responsiveness of five at-work productivity measures in patients with rheumatoid arthritis or osteoarthritis. *Arthritis Care and Research*. https://doi.org/10.1002/acr.20011

54. Beaton, D.E., Dyer, S., Boonen, A., (...), Tugwell, P.S. (2016) OMERACT filter evidence supporting the measurement of at-work productivity loss as an outcome measure in rheumatology research. *Journal of Rheumatology*. https://doi.org/10.3899/jrheum.141077

55. Van Eerd, D., Munhall, C., Irvin, E., (...), Amick, B. (2016) Effectiveness of workplace interventions in the prevention of upper extremity musculoskeletal disorders and symptoms: An update of the evidence. *Occupational and Environmental Medicine*. https://doi.org/10.1136/oemed-2015-102992

56. Hoe, V.C., Urquhart, D.M., Kelsall, H.L., (...), Sim, M.R. (2018) Ergonomic interventions for preventing work-related musculoskeletal disorders of the upper limb and neck among office workers. *Cochrane Database of Systematic Reviews*. https://doi.org/10.1002/14651858.CD008570.pub3

57. Tuntland, H., Kjeken, I., Nordheim, L.V., (...), Hagen, K.B. (2009) Assistive technology for rheumatoid arthritis. *Cochrane Database of Systematic Reviews*. https://doi.org/10.1002/14651858.CD006729.pub2

58. Tuntland, H., Kjeken, I., Nordheim, L., (...), Hagen, K. (2010) The Cochrane review of assistive technology for rheumatoid arthritis. *European Journal of Physical and Rehabilitation Medicine*. https://www.scopus.com/pages/publications/77956362391

59. Yildirim, M.H., Yildirim, E.A., Ca¸rpar, E., (...), Cantürk, G. (2018) Hand functions in patients with schizophrenia: A clinical comparison with bipolar disorder and healthy subjects. *Comprehensive Psychiatry*. https://doi.org/10.1016/j.comppsych.2018.08.009

60. Calder, K.M., Galea, V., Wessel, J., (...), MacIntyre, N.J. (2011) Muscle activation during hand dexterity tasks in women with hand osteoarthritis and control subjects. *Journal of Hand Therapy*. https://doi.org/10.1016/j.jht.2010.11.003

61. Calder, Kristina M, Martin, Alison, Lydiate, Jessica, (...), MacIntyre, Norma J (2012) Sensory nerve action potentials and sensory perception in women with arthritis of the hand. *Journal of NeuroEngineering and Rehabilitation*. https://doi.org/10.1186/1743-0003-9-27

62. Bell, A.F., Walton, K.L., Tapsell, L.C. (2016) Easy to open? Exploring the 'openability' of hospital food and beverage packaging by older adults. *Appetite*. https://doi.org/10.1016/j.appet.2015.12.004

63. Bobos, P., Nazari, G., Szekeres, M., (...), MacDermid, J.C. (2019) The effectiveness of joint-protection programs on pain, hand function, and grip strength levels in patients with hand arthritis: A systematic review and meta-analysis. *Journal of Hand Therapy*. https://doi.org/10.1016/j.jht.2018.09.012

64. Estee, Mahnuma Mahfuz, Cicuttini, Flavia M., Page, Matthew J., (...), Wang, Yuanyuan (2022) Efficacy of corticosteroids for hand osteoarthritis - a systematic review and meta-analysis of randomized controlled trials. *BMC Musculoskeletal Disorders*. https://doi.org/10.1186/s12891-022-05619-9

65. Khominets, V.V., Peleshok, S.A., Volov, D.A., (...), Nebylitsa, Ya.I. (2020) 3D printing technologies in the treatment of patients with injuries and diseases of the forearm and hand. *Vestnik Rossijskoj Voenno-Medicinskoj Akademii*. https://doi.org/10.17816/brmma25978

66. Greenwood, M.C., Hakim, A.J., Carson, E., Doyle, D.V. (2006) Touch-screen computer systems in the rheumatology clinic offer a reliable and user-friendly means of collecting quality-of-life and outcome data from patients with rheumatoid arthritis. *Rheumatology*. https://doi.org/10.1093/rheumatology/kei100

67. Tonga, E., Williamson, E., Srikesavan, C., (...), Lamb, S.E. (2021) A hand exercise mobile app for people with rheumatoid arthritis in Turkey: design, development and usability study. *Rheumatology International*. https://doi.org/10.1007/s00296-021-04860-0

68. Barghi, E.G., Mohammadzadeh, N., Enteshari-Moghadam, A.E. (2023) Self-care application for rheumatoid arthritis: Identifying key data elements. *Frontiers in Health Informatics*. https://doi.org/10.30699/fhi.v12i0.504

69. Barelli, Renato G., Avelino, Valter F., Castro, Maria Claudia F. (2022) STIMGRASP: A Home-Based Functional Electrical Stimulator for Grasp Restoration in Daily Activities. *Sensors (Basel, Switzerland)*. https://doi.org/10.3390/s23010010

70. Baker, N.A., Rogers, J.C. (2010) Association between computer use speed and age, impairments in function, and touch typing training in people with rheumatoid arthritis.. *Arthritis care & research*. https://doi.org/10.1002/acr.20074

71. Fager, S.K., Sorenson, T., Jakobs, E., (...), Beukelman, D.R. (2023) Multi-modal access method (eye-tracking + switch-scanning) for individuals with severe motor impairment: A preliminary investigation. *Assistive Technology*. https://doi.org/10.1080/10400435.2022.2053895

72. Bruno Garza, J.L., Young, J.G. (2015) A literature review of the effects of computer input device design on biomechanical loading and musculoskeletal outcomes during computer work. *Work*. https://doi.org/10.3233/WOR-152161

73. Mitchell, C., Cler, G., Fager, S., (...), Vojtech, J. (2022) Ability-based Keyboards for Augmentative and Alternative Communication: Understanding How Individuals' Movement Patterns Translate to More Efficient Keyboards: Methods to Generate Keyboards Tailored to User-specific Motor Abilities. *Conference on Human Factors in Computing Systems - Proceedings*. https://doi.org/10.1145/3491101.3519845

74. Lin, Y.-L., Chen, M.-C., Yeh, Y.-M., (...), Yeh, C.-C. (2006) Design and implementation of a chorded on-screen keyboard for people with physical impairments. *Lecture Notes in Computer Science (including subseries Lecture Notes in Artificial Intelligence and Lecture Notes in Bioinformatics)*. https://doi.org/10.1007/11788713_143

75. Arias-Flores, H., Valencia-Aragón, K., Sanchez-Gordon, S. (2024) Exploring the Effectiveness of Assistive Technology: A Preliminary Case Study Using Makey Makey, Tobii Eye Tracker, and Leap Motion. *Lecture Notes in Computer Science (including subseries Lecture Notes in Artificial Intelligence and Lecture Notes in Bioinformatics)*. https://doi.org/10.1007/978-3-031-71704-8_3

76. Capilouto, G.J., McClenaghan, B., Williams, H.G., (...), Hussey, J.R. (2004) Performance of able-bodied subjects on a text-typing task using a head-operated device and expanded membrane cursor keys. *Perceptual and Motor Skills*. https://doi.org/10.2466/pms.98.1.147-162

77. Metz, Stephen, Isle, Brian, Denno, Sandra, Li, Wang (1990) Small rotary controls. Limitations for people with arthritis. *Proceedings of the Human Factors Society*. https://doi.org/10.1177/154193129003400202

78. Austin, L., Sharp, C.A., Van der Veer, S.N., (...), Dixon, W.G. (2020) Providing 'the bigger picture': Benefits and feasibility of integrating remote monitoring from smartphones into the electronic health record. *Rheumatology (United Kingdom)*. https://doi.org/10.1093/rheumatology/kez207

79. Zuidema, R., van Dulmen, S., Sanden, M.N.V.-D., (...), Van Gaal, B. (2019) Efficacy of a web-based self-management enhancing program for patients with rheumatoid arthritis: Explorative randomized controlled trial. *Journal of Medical Internet Research*. https://doi.org/10.2196/12463

80. Weiner, Joan, Lui, Geyanne, Brown, Mackenzie, (...), Navarro-Millán, Iris (2022) *Pilot and Feasibility Studies*. https://doi.org/10.1186/s40814-022-01041-z

81. Seljelid, Berit, Varsi, Cecilie, Solberg Nes, Lise, (...), Børøsund, Elin (2020) Content and system development of a digital patient-provider communication tool to support shared decision making in chronic health care: InvolveMe. *BMC Medical Informatics and Decision Making*. https://doi.org/10.1186/s12911-020-1065-8

82. Ali, Lilas, Wallström, Sara, Barenfeld, Emmelie, (...), Ekman, Inger (2020) Person-centred care by a combined digital platform and structured telephone support for people with chronic obstructive pulmonary disease and/or chronic heart failure: study protocol for the PROTECT randomised controlled trial. *BMJ Open*. https://doi.org/10.1136/bmjopen-2019-036356

83. Diaz-Martinez, J., Leon Morris, S., Delgado-Enciso, I., (...), Hospital, M. (2026) An Integrated, Evidence-Guided Framework for Chronic Kidney Disease Self-Management: Behavioral Mechanisms, Implementation Strategies, and Real-World Application. *Kidney and Blood Pressure Research*. https://doi.org/10.1159/000552085

84. Madanian, S., Nakarada-Kordic, I., Reay, S., Chetty, T. (2023) Patients' perspectives on digital health tools. *PEC Innovation*. https://doi.org/10.1016/j.pecinn.2023.100171

85. Sumner, J., Ng, C.W.T., Teo, K.E.L., (...), Lim, Y.W. (2024) Co-designing care for multimorbidity: a systematic review. *BMC Medicine*. https://doi.org/10.1186/s12916-024-03263-9

86. https://www.sciencedirect.com/science/article/pii/S2056752926001364

87. Bakker, N.F., van Weely, S.F.E., Hutting, N., (...), Knoop, J. (2024) Development of a Multimodal, Physiotherapist-Led, Vocational Intervention for People with Inflammatory Arthritis and Reduced Work Ability: A Mixed-Methods Design Study. *Journal of Occupational Rehabilitation*. https://doi.org/10.1007/s10926-023-10170-y

88. Lopez-Olivo, M.A., Ingleshwar, A., Volk, R.J., (...), Suarez-Almazor, M.E. (2018) Development and Pilot Testing of Multimedia Patient Education Tools for Patients With Knee Osteoarthritis, Osteoporosis, and Rheumatoid Arthritis. *Arthritis Care and Research*. https://doi.org/10.1002/acr.23271

89. Braithwaite, F.A., Arnold, J., Davis, A., (...), Stanton, T.R. (2023) Osteoarthritis consumers as co-researchers: identifying consumer insights to improve osteoarthritis management by co-designing translational research solutions. *Osteoarthritis and Cartilage*. https://doi.org/10.1016/j.joca.2023.03.004

90. Pouliopoulou, D.V., MacDermid, J.C., Billias, N., (...), Bobos, P. (2025) Understanding patient needs and preferences for joint protection in hand osteoarthritis through mixed methods patient co-design. *Scientific Reports*. https://doi.org/10.1038/s41598-025-13734-5

91. Joshi, S.G., Valen, J. (2018) Co-exploring interaction opportunities for enabling technologies for people with rheumatic disorder. *Lecture Notes in Computer Science (including subseries Lecture Notes in Artificial Intelligence and Lecture Notes in Bioinformatics)*. https://doi.org/10.1007/978-3-319-94277-3_65

92. Williams, A.D., O'Brien, J., Mulford, J., (...), Bird, M.-L. (2022) Effect of combined exercise training and behaviour change counselling versus usual care on physical activity in patients awaiting hip and knee arthroplasty: A randomised controlled trial. *Osteoarthritis and Cartilage Open*. https://doi.org/10.1016/j.ocarto.2022.100308

93. Li, T., Wells, G., Westhovens, R., Tugwell, P. (2009) Validation of a simple activity participation measure for rheumatoid arthritis clinical trials. *Rheumatology*. https://doi.org/10.1093/rheumatology/ken448

94. https://www.sciencedirect.com/science/article/pii/S2198658422000245

95. Daste, C., Rannou, F., Mouthon, L., (...), Nguyen, C. (2019) Patient acceptable symptom state and minimal clinically important difference for patient-reported outcomes in systemic sclerosis: A secondary analysis of a randomized controlled trial comparing personalized physical therapy to usual care. *Seminars in Arthritis and Rheumatism*. https://doi.org/10.1016/j.semarthrit.2018.03.013

96. Ward, M.M., Alba, M.I. (2022) Estimates of minimal clinically important improvments vary with the responsiveness of the sample. *Journal of Clinical Epidemiology*. https://doi.org/10.1016/j.jclinepi.2021.11.002

97. de Windt, Tommy S., Vonk, Lucienne A., Slaper-Cortenbach, Ineke C. M., (...), Saris, Daniel B. F. (2017) Allogeneic Mesenchymal Stem Cells Stimulate Cartilage Regeneration and Are Safe for Single-Stage Cartilage Repair in Humans upon Mixture with Recycled Autologous Chondrons. *Stem Cells*. https://doi.org/10.1002/stem.2475

98. Negrini, Francesco, De Lucia, Francesco, Negrini, Stefano, (...), de Girolamo, Laura (2021) Case Report: Rehabilitation After Platelet-Rich Growth Factors' Intra-Articular Injections for Knee Osteoarthritis: Two Case Reports of a Home-Based Protocol. *Frontiers in Pharmacology*. https://doi.org/10.3389/fphar.2021.718060

99. Allin, S., Eckel, E., Markham, H., Brewer, B.R. (2010) Recent Trends in the Development and Evaluation of Assistive Robotic Manipulation Devices. *Physical Medicine and Rehabilitation Clinics of North America*. https://doi.org/10.1016/j.pmr.2009.09.001

100. Nordenskiüld, U. (1994) Evaluation of assistive devices after a course in joint protection. *International Journal of Technology Assessment in Health Care*. https://doi.org/10.1017/S0266462300006255

101. Leese, M.I., Dorociak, K.E., Noland, M., (...), Seelye, A. (2021) Use of in-home activity monitoring technologies in older adult veterans with mild cognitive impairment: The impact of attitudes andcognition. *Gerontechnology*. https://doi.org/10.4017/GT.2021.20.2.10.06

102. Zhang, S., Wu, M., Sun, R., (...), Gong, N. (2025) Exploring the Discontinuous Usage Behavior of Digital Cognitive Training Among Older Adults With Mild Cognitive Impairment and Their Family Members: Qualitative Study Using the Extended Model of IT Continuance. *Journal of Medical Internet Research*. https://doi.org/10.2196/66393

103. Schmitter-Edgecombe, M., Luna, C., Chilton, R.C., (...), Cook, D.J. (2026) Promoting digital memory aid use in older adults with cognitive concerns: A pilot randomized controlled trial of adaptive web-based training. *Neuropsychology*. https://doi.org/10.1037/neu0001071