


# Prediction of Cardiac Arrest From Physiological Signals in the Pediatric ICU

S. Tonekaboni, M. Brudno, A. Goldenberg (University of Toronto)


M. Mazwi, P. Laussen, D. Eytan, R. Greer, S. D. Goodfellow, A. Goodwin (Sickkids hospital)

# Cardiac Arrest in Pediatric ICU

- Many patients in the ICU are at risk of a sudden cardiac arrest
- In-hospital cardiac arrest is strongly associated with mortality
- Only a limited number of interventions are available to deal with this issue, that do not guarantee disability-free survival




# Learning the Existing Pattern in Pre-arrest Physiological Signals



# Challenges:

1. Clinical data is usually corrupted by noise or missing values
2. Time-series physiological recording is densely packed with information.
3. Typical ICU practice is to record vital signs based on perceived patient needs, so not all patients have a consistent set of measurements in their records




# 1) Missing Signal Imputation

Major benefits:


- It can be online
- Smoothens signal and reduces frequency variation

## Low-pass filtering method missing value imputation



## 2) Risk Predictor Model Architecture:


Combining Convolutional and Recurrent networks for the analysis of longitudinal physiological signals



- CNN: Extracts a **compact latent representation** of the window of recording.

## 2) Risk Predictor Model Architecture


- LSTM: Learn the existing temporal dependencies to predict the probability of an arrest in the future



$r$  is feed into a simple regression model along with the **age** of a patient, to generate **risk**, that is the risk score associated to a time window of a certain physiological signal.

$f$  is the condensed latent representation of the original physiological signal extracted by the CNN

## 2) Risk Predictor Model Comparison:



Signal Type	CNN	LSTM	CNN+LSTM
Heart rate (HR)	$0.753 \pm 0.075$	$0.666 \pm 0.118$	$0.816 \pm 0.034$
Respiratory rate (Resp)	$0.739 \pm 0.043$	N/A	$0.67 \pm 0.05$
Pulse rate (Pulse)	$0.720 \pm 0.093$	$0.698 \pm 0.06$	$0.746 \pm 0.092$
Oxygen Saturation level (SpO2)	$0.714 \pm 0.078$	$0.617 \pm 0.101$	$0.723 \pm 0.055$
Ambulatory Blood Pressure (ABP)	$0.675 \pm 0.17$	$0.647 \pm 0.17$	$0.76 \pm 0.13$
SDNN Heart Rate Variability Metric (SDNN)	$0.663 \pm 0.080$	N/A	$0.692 \pm 0.07$


\*\* Numbers reported are the F1 scores



### 3) The Ensemble Structure

Major Benefits:

1. Robust to missing measurements
2. Identifying pathways resulting in an arrest




$$Riskscore = \frac{\sum_{i=1}^n (W_i * risk_i) \mathbb{1}_i}{\sum_{i=1}^n (W_i) \mathbb{1}_i}$$

Where  $\mathbb{1}_i = 0$  if signal  $i$  is missing and  $\mathbb{1}_i = 1$  otherwise



# Future direction: Towards Predictive Personalized Medicine

Approaches that offer promising future for **predictive** rather than **reactive** medicine, in a personalized framework where clinical decision support can be provided at the level of individual patients.



# Major Takeaways:

- The low-pass filtering method is a robust technique for missing value imputation in temporal signals
- CNNs are powerful tools for extracting features from high-dimensional space data and in combination with RNNs, it can also model temporal dependencies
- Ensemble models are helpful tools in cases where samples might have inconsistent set of features.



# Thank you!



Prediction of Cardiac Arrest From physiological Signals in the Pediatric ICU

S. Tonekaboni, M. Brudno, A. Goldenberg (UofT)

M. Mazwi, P. Laussen, D. Eytan, R. Greer,

S. D. Goodfellow, A. Goodwin (Sickkids hospital)