

## Evaluating a Hypothesis

Once we have done some trouble shooting for errors in our predictions by:

- Getting more training examples
- Trying smaller sets of features
- Trying additional features
- Trying polynomial features

hypothesis, given a dataset of training examples, we can split up the data into two sets: a **training set** and a **test set**. Typically, the training set consists of 70 % of your data and the test set is the remaining 30 %.

The new procedure using these two sets is then:

1. Learn  $\Theta$  and minimize  $J_{train}(\Theta)$  using the training set
2. Compute the test set error  $J_{test}(\Theta)$

### The test set error

1. For linear regression:  $J_{test}(\Theta) = \frac{1}{2m_{test}} \sum_{i=1}^{m_{test}} (h_{\Theta}(x_{test}^{(i)}) - y_{test}^{(i)})^2$
2. For classification ~ Misclassification error (aka 0/1 misclassification error):

$$err(h_{\Theta}(x), y) = \begin{cases} 1 & \text{if } h_{\Theta}(x) \geq 0.5 \text{ and } y = 0 \text{ or } h_{\Theta}(x) < 0.5 \text{ and } y = 1 \\ 0 & \text{otherwise} \end{cases}$$

This gives us a binary 0 or 1 error result based on a misclassification. The average test error for the test set is:

$$\text{Test Error} = \frac{1}{m_{test}} \sum_{i=1}^{m_{test}} err(h_{\Theta}(x_{test}^{(i)}), y_{test}^{(i)})$$

This gives us the proportion of the test data that was misclassified.