DESKTOP VERSION

To: <address@email.com>
From: PDQ Communications <info@pdqcom.com>
Initial Subject Line: Ready to prescribe BRINSUPRI™ (brensocatib)?
Preheader: Learn about getting started

Echo1 Subject Line: Prescribing BRINSUPRI™ (brensocatib)
Preheader: A few helpful tips to get started

Echo2 Subject Line: BRINSUPRI™ (brensocatib) Rx: Start here
Preheader: Open for dosing, indication, and pharmacy info

A https://ad.doubleclick.net/ddm/clk/621568076;428493034;n;gdpr=${GDPR};gdpr_consent=${GDPR_CONSENT_755}

B https://ad.doubleclick.net/ddm/clk/621883231;429065142;c;gdpr=${GDPR};gdpr_consent=${GDPR_CONSENT_755}

C https://ad.doubleclick.net/ddm/clk/621883234;429065145;i;gdpr=${GDPR};gdpr_consent=${GDPR_CONSENT_755}

D https://ad.doubleclick.net/ddm/clk/621883240;429065151;c;gdpr=${GDPR};gdpr_consent=${GDPR_CONSENT_755}

E https://ad.doubleclick.net/ddm/clk/621883243;429065154;i;gdpr=${GDPR};gdpr_consent=${GDPR_CONSENT_755}

F https://ad.doubleclick.net/ddm/clk/621883246;429065157;o;gdpr=${GDPR};gdpr_consent=${GDPR_CONSENT_755}

G https://ad.doubleclick.net/ddm/clk/621883249;429065160;l;gdpr=${GDPR};gdpr_consent=${GDPR_CONSENT_755}

H https://ad.doubleclick.net/ddm/clk/621883252;429065163;i;gdpr=${GDPR};gdpr_consent=${GDPR_CONSENT_755}

I https://ad.doubleclick.net/ddm/clk/621883255;429065166;o;gdpr=${GDPR};gdpr_consent=${GDPR_CONSENT_755}

J https://insmed.com/state-disclosure-information/

K Links to PDQ Unsubscribe

HEALTHCARE PROFESSIONAL'S FLASH

Get your patients started

Dear Dr [variable],

Here are some prescribing details to keep in mind[1]:

BRINSUPRI (brensocatib) is indicated for the treatment of non-cystic fibrosis bronchiectasis in adult and pediatric patients 12 years of age and older

Available in dosage strengths of 10 mg or 25 mg

Oral, once-daily dosing

Learn how to prescribe B

How to prescribe

Initiate patients on BRINSUPRI using the BRINSUPRI Prescription and *inLighten*™ Enrollment Form.

or

E-prescribe via an EHR.

BRINSUPRI is a limited-distribution product and can only be dispensed by[a]:

Amber Specialty Pharmacy
(**888-370-1724**) (NCPDP code: 2815338)

Maxor Specialty Pharmacy, A VytlOne Company
(**800-658-6046**) (NCPDP code: 4575518)

PANTHERx Rare Pharmacy
(**855-726-8479**) (NCPDP code: 6008002)

aSeparate distribution is available for 340B-eligible entities and patients insured by the US Department of Veterans Affairs.

Get patients started C

Want to learn more? Visit BRINSUPRIhcp.com to request a visit from a representative

Contact us D

IMPORTANT SAFETY INFORMATION

WARNINGS AND PRECAUTIONS

Dermatologic Adverse Reactions
Treatment with BRINSUPRI is associated with an increase in dermatologic adverse reactions, including rash, dry skin, and hyperkeratosis. Monitor patients for development of new rashes or skin conditions and refer patients to a dermatologist for evaluation of new dermatologic findings.

Gingival and Periodontal Adverse Reactions
Treatment with BRINSUPRI is associated with an increase in gingival and periodontal adverse reactions. Refer patients to dental care services for regular dental checkups while taking BRINSUPRI. Advise patients to perform routine dental hygiene.

Live Attenuated Vaccines
It is unknown whether administration of live attenuated vaccines during BRINSUPRI treatment will affect the safety or effectiveness of these vaccines. The use of live attenuated vaccines should be avoided in patients receiving BRINSUPRI.

ADVERSE REACTIONS

The most common adverse reactions ≥2% in the ASPEN trial included upper respiratory tract infection, headache, rash, dry skin, hyperkeratosis, and hypertension. The safety profile for adult patients with NCFB in WILLOW was generally similar to ASPEN, except for a higher incidence of gingival and periodontal adverse reactions.

Less Common Adverse Reactions

Liver Function Test Elevations
In ASPEN, there was an increase from baseline in average ALT, AST, and alkaline phosphatase levels at all time points from Week 4 through Week 56 in both BRINSUPRI 10 mg and 25 mg arms compared to placebo. The incidence of ALT >3X upper limit of normal (ULN) was 0%, 1.2%, and 0.9%; the incidence of AST >3X ULN was 0.2%, 0.3%, and 0.5%; and the incidence of alkaline phosphatase >1.5X ULN was 2.5%, 4.1%, and 4.0% in patients treated with placebo and BRINSUPRI 10 mg and 25 mg, respectively.

Skin Cancers
In ASPEN, the incidence of skin cancers among patients treated with BRINSUPRI 10 mg and 25 mg was 0.5% and 1.9%, respectively, compared to 1.1% in placebo-treated patients.

Alopecia
In ASPEN, the incidence of alopecia among patients treated with BRINSUPRI 10 mg and 25 mg was 1.5% and 1.6% respectively, compared to 0.4% in placebo-treated patients.

USE IN SPECIFIC POPULATIONS

Pregnancy: There are no clinical data on the use of BRINSUPRI in pregnant women.

Lactation: There is no information regarding the presence of BRINSUPRI and/or its metabolite(s) in human milk, the effects on the breastfed infant, or the effects on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for BRINSUPRI and any potential adverse effects on the breastfed child from BRINSUPRI or from the underlying maternal condition.

Pediatric use: The safety and effectiveness of BRINSUPRI for the treatment of NCFB have been established in pediatric patients aged 12 years and older. Common adverse reactions in pediatric patients aged 12 years and older enrolled in ASPEN were consistent with those in adults. The safety and effectiveness of BRINSUPRI have not been established in pediatric patients younger than 12 years of age.

INDICATION

BRINSUPRI is indicated for the treatment of non-cystic fibrosis bronchiectasis (NCFB) in adult and pediatric patients 12 years of age and older.

Please see full Prescribing Information. E

EHR=electronic health record; NCPDP=National Council for Prescription Drug Programs.

Reference: 1. BRINSUPRI [package insert]. Bridgewater, NJ: Insmed Incorporated; 2025.

F insmed G

This email was sent to pdqcom.8ldd8@litmusemail.com. If you wish to discontinue receiving messages from Insmed Incorporated, click here to unsubscribe.

H Terms of use | Privacy policy I

This email was sent by: **Insmed Incorporated**
700 US Highway 202/206, Bridgewater, NJ 08807

For more information on the price of BRINSUPRI™ (brensocatib) 10 mg and 25 mg as well as information on the variation in efficacy of BRINSUPRI among different racial and ethnic groups, please visit insmed.com/state-disclosure-information/. Insmed has not identified any generic prescription drugs in the same therapeutic class with the same or similar mechanism of action.

J © 2025 Insmed Incorporated. All Rights Reserved. Insmed, BRINSUPRI, and inLighten are trademarks of Insmed Incorporated. All other trademarks are property of their respective owner. PP-BRIN-US-00383

Healthcare Professional's Flash is a Product of PDQ Communications - Healthcare Information in a *Flash!*

If you wish to discontinue receiving messages from PDQ Communications, Click here to unsubscribe.
PDQ Communications | PO Box 191 | Carle Place, NY 11514 K

MOBILE VERSION

To: <address@email.com>
From: PDQ Communications <info@pdqcom.com>
Initial Subject Line: Ready to prescribe BRINSUPRI™ (brensocatib)?
Preheader: Learn about getting started

Echo1 Subject Line: Prescribing BRINSUPRI™ (brensocatib)
Preheader: A few helpful tips to get started

Echo2 Subject Line: BRINSUPRI™ (brensocatib) Rx: Start here
Preheader: Open for dosing, indication, and pharmacy info

A https://ad.doubleclick.net/ddm/clk/621568076;428493034;n;gdpr=${GDPR};gdpr_consent=${GDPR_CONSENT_755}

B https://ad.doubleclick.net/ddm/clk/621883231;429065142;c;gdpr=${GDPR};gdpr_consent=${GDPR_CONSENT_755}

C https://ad.doubleclick.net/ddm/clk/621883234;429065145;i;gdpr=${GDPR};gdpr_consent=${GDPR_CONSENT_755}

D https://ad.doubleclick.net/ddm/clk/621883240;429065151;c;gdpr=${GDPR};gdpr_consent=${GDPR_CONSENT_755}

E https://ad.doubleclick.net/ddm/clk/621883243;429065154;i;gdpr=${GDPR};gdpr_consent=${GDPR_CONSENT_755}

F https://ad.doubleclick.net/ddm/clk/621883246;429065157;o;gdpr=${GDPR};gdpr_consent=${GDPR_CONSENT_755}

G https://ad.doubleclick.net/ddm/clk/621883249;429065160;l;gdpr=${GDPR};gdpr_consent=${GDPR_CONSENT_755}

H https://ad.doubleclick.net/ddm/clk/621883252;429065163;i;gdpr=${GDPR};gdpr_consent=${GDPR_CONSENT_755}

I https://ad.doubleclick.net/ddm/clk/621883255;429065166;o;gdpr=${GDPR};gdpr_consent=${GDPR_CONSENT_755}

J https://insmed.com/state-disclosure-information/

K Links to PDQ Unsubscribe

HEALTHCARE PROFESSIONAL'S FLASH

Get your patients started

Dear Dr [variable],

Here are some prescribing details to keep in mind[1]:

BRINSUPRI (brensocatib) is indicated for the treatment of non-cystic fibrosis bronchiectasis in adult and pediatric patients 12 years of age and older

Available in dosage strengths of 10 mg or 25 mg

Oral, once-daily dosing

Learn how to prescribe B

How to prescribe

Initiate patients on BRINSUPRI using the BRINSUPRI Prescription and *inLighten*™ Enrollment Form.

or

E-prescribe via an EHR.

BRINSUPRI is a limited-distribution product and can only be dispensed by[a]:

Amber Specialty Pharmacy
(**888-370-1724**) (NCPDP code: 2815338)

Maxor Specialty Pharmacy, A VytlOne Company
(**800-658-6046**) (NCPDP code: 4575518)

PANTHERx Rare Pharmacy
(**855-726-8479**) (NCPDP code: 6008002)

[a]Separate distribution is available for 340B-eligible entities and patients insured by the US Department of Veterans Affairs.

Get patients started C

Want to learn more? Visit BRINSUPRIhcp.com to request a visit from a representative

Contact us D

IMPORTANT SAFETY INFORMATION

WARNINGS AND PRECAUTIONS

Dermatologic Adverse Reactions
Treatment with BRINSUPRI is associated with an increase in dermatologic adverse reactions, including rash, dry skin, and hyperkeratosis. Monitor patients for development of new rashes or skin conditions and refer patients to a dermatologist for evaluation of new dermatologic findings.

Gingival and Periodontal Adverse Reactions
Treatment with BRINSUPRI is associated with an increase in gingival and periodontal adverse reactions. Refer patients to dental care services for regular dental checkups while taking BRINSUPRI. Advise patients to perform routine dental hygiene.

Live Attenuated Vaccines
It is unknown whether administration of live attenuated vaccines during BRINSUPRI treatment will affect the safety or effectiveness of these vaccines. The use of live attenuated vaccines should be avoided in patients receiving BRINSUPRI.

ADVERSE REACTIONS

The most common adverse reactions ≥2% in the ASPEN trial included upper respiratory tract infection, headache, rash, dry skin, hyperkeratosis, and hypertension. The safety profile for adult patients with NCFB in WILLOW was generally similar to ASPEN, except for a higher incidence of gingival and periodontal adverse reactions.

Less Common Adverse Reactions

Liver Function Test Elevations
In ASPEN, there was an increase from baseline in average ALT, AST, and alkaline phosphatase levels at all time points from Week 4 through Week 56 in both BRINSUPRI 10 mg and 25 mg arms compared to placebo. The incidence of ALT >3X upper limit of normal (ULN) was 0%, 1.2%, and 0.9%; the incidence of AST >3X ULN was 0.2%, 0.3%, and 0.5%; and the incidence of alkaline phosphatase >1.5X ULN was 2.5%, 4.1%, and 4.0% in patients treated with placebo and BRINSUPRI 10 mg and 25 mg, respectively.

Skin Cancers
In ASPEN, the incidence of skin cancers among patients treated with BRINSUPRI 10 mg and 25 mg was 0.5% and 1.9%, respectively, compared to 1.1% in placebo-treated patients.

Alopecia
In ASPEN, the incidence of alopecia among patients treated with BRINSUPRI 10 mg and 25 mg was 1.5% and 1.6% respectively, compared to 0.4% in placebo-treated patients.

USE IN SPECIFIC POPULATIONS

Pregnancy: There are no clinical data on the use of BRINSUPRI in pregnant women.

Lactation: There is no information regarding the presence of BRINSUPRI and/or its metabolite(s) in human milk, the effects on the breastfed infant, or the effects on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for BRINSUPRI and any potential adverse effects on the breastfed child from BRINSUPRI or from the underlying maternal condition.

Pediatric use: The safety and effectiveness of BRINSUPRI for the treatment of NCFB have been established in pediatric patients aged 12 years and older. Common adverse reactions in pediatric patients aged 12 years and older enrolled in ASPEN were consistent with those in adults. The safety and effectiveness of BRINSUPRI have not been established in pediatric patients younger than 12 years of age.

INDICATION

BRINSUPRI is indicated for the treatment of non-cystic fibrosis bronchiectasis (NCFB) in adult and pediatric patients 12 years of age and older.

Please see full Prescribing Information. E

EHR=electronic health record; NCPDP=National Council for Prescription Drug Programs.

Reference: 1. BRINSUPRI [package insert]. Bridgewater, NJ: Insmed Incorporated; 2025.

F insmed G
This email was sent to pdqcom.89d8@btmusemail.com. If you wish to discontinue receiving messages from Insmed Incorporated, click here to unsubscribe.
H Terms of use Privacy policy I
This email was sent by: **Insmed Incorporated**
700 US Highway 202/206, Bridgewater, NJ 08807

For more information on the price of BRINSUPRI™ (brensocatib) 10 mg and 25 mg as well as information on the variation in efficacy of BRINSUPRI among different racial and ethnic groups, please visit insmed.com/state-disclosure-information/. Insmed has not identified any generic prescription drugs in the same therapeutic class with the same or similar mechanism of action.

© 2025 Insmed Incorporated. All Rights Reserved. Insmed, BRINSUPRI, and inLighten are trademarks of Insmed Incorporated. All other trademarks are property of their respective owner.
PP-BRIN-US-00383
J

Healthcare Professional's Flash is a Product of PDQ Communications - Healthcare Information in a *Flash!*

If you wish to discontinue receiving messages from PDQ Communications, Click here to unsubscribe.
PDQ Communications | PO Box 191 | Carle Place, NY 11514 K